

SUBJECT INDEX

3

3WayPack, xvii, xxi, 63, 73, 431, 488

A

additive adjustment, *see* centering

additive and multiplicative modeling, 128

additive modeling

two-way, 111

agriculture, xviii, 13, 17, 23

AMMI model, 135

biadditive model, 135

double centering, 134

genotype by environment interaction, 111, 135

multiattribute, 129

preprocessing, 129

missing data, 173

multiway factorial data, 36

nonstochastic multiway data, 32, 146

peanut example, 5-6

plant adaptation example, 6

plant breeding example, 5, 111, 474

plant breeding

missing data, 145

phenotypic correlation, 129

preprocessing factorial designs, 129

preprocessing, 130

research questions, 18

three-mode mixture method of clustering, 432

three-way profile data, 403

triadditive modeling, 139

variety trials, 173

air pollution, multiway modeling, 480

algorithms, *see* TUCKALS, Tucker, Parafac, etc

algorithms

definition, 79

reasons for discussing, 77

all-components plot, *see* plotting

ALSCOMP3, 42, 74, 96, 321

alternating least-squares, 21, 80

analysis of dependence, 32, 283

analysis of interdependence, 32, 283

analytical chemistry, 13, 17, 22

application Paratuck2 model, 64

chromatography time-shift example, 475

analytical chemistry (*cont'd*)
 component uniqueness, 61, 322
 explicit Parafac models, 61, 344
 fluorescence spectroscopy, 472
 four-way data, 472
 multiway block models, 24
 nonnegative components, 96
 Parafac2 applications, 63
 problems with split-half procedures, 134
 spectroscopy REM example, 472
 spectroscopy protein example, 472
 applications, standard format, xix
 aspect ratio, 260, 497
 association models, 42, 434
 autoregressive model, 380, 385
 external variables, 380–381
 structural part, 380–381
 transition matrix, 381
 autoregressive process, 379, 384, 402
 core array, 384
 estimation parameters, 380, 383, 391
 first-order, 380, 384, 391–392
 homogeneity, 384
 lag-one regression analysis, 391
 latent covariance matrix, 384
 simulation, 383
 parameter estimation, 386
 second-order, 384, 392
 simplex, 380
 examples, 391
 stability, 384, 386, 392
 state-like, 384
 stationarity, 384, 386
 three-mode PCA, 379
 simulation, 379
 trait-like, 384
 average subject's profile, 130, 271, 277, 321, 332,
 395

B

B-splines
 compression, 103
 smoothing, 71, 377
 bad leverage points, 295
 basic data relation matrix=BDRM, 469
 basic form
 component models, 215
 Parafac model, 60, 215
 PCA, 47
 Tucker models, 215, 225, 335
 Tucker2 model, 66
 Tucker3 model, 65, 140
 basic structure of a matrix, *see* singular value
 decomposition

batch processes
 monitoring, 52
 between-model selection, 175
 deviance plot, 182
 examples, 162, 197, 199, 202, 206
 model hierarchy, 186
 examples, 205
 multiway score plot, 182
 biadditive model, 135–136
 additive plus multiplicative model, 135
 completely multiplicative model, 137
 degrees of freedom, 137
 factorial design, 135
 hypothesis testing, 137
 robust variant, 136
 bibliography
 annotated, 21
 web-based, 21
 bilinear model, 19, 217
 biplot, 332, 492
 aspect ratio, 497
 biplot axis, 495
 calibration, 496
 column markers, 495
 description, 492, 494
 Euclidean distance, 493
 geometry, 499
 graphic, 495
 inner product, 497
 interpretation, 497
 least-squares approximation, 493
 low-dimensional approximation, 493
 principal component analysis, 499
 principal coordinate scaling, 493, 496–498
 row markers, 494
 symmetric scaling, 493, 496–497, 499
 total variability, 493
 variability accounted for, 494
 bipolar scales, 8
 antonym, 9
 centering, 131, 360
 double centering, 131
 multivariate multiway factorial data, 131
 three-way rating scales, 360
 triadditive modelling, 139
 blind beamforming, 22
 blindfolded mountaineer
 model for iterative algorithms, 80–81
 Boolean algebra, 42, 460
 bootstrap, 233, 312, 322, 338, 449, 478
 basics, 188
 dimensionality selection, 176
 examples, 190, 207
 fit measures, 188

multiway analysis, 188
 nonstochastic data, 147
 optimizing congruences, 234
 sample size, 176
 sampling residuals, 233
 stability estimation, 7, 18, 32, 176, 188, 362
 stochastics, 233
 target rotation, 234

C

CANDELINC, 95, 234
 canonical correlation analysis, 401
 canonical decomposition, CANDECOMP, *see*
 also Parafac
 canonical decomposition, CANDECOMP
 individual differences scaling, 22
 linear constraints, CANDELINC, 94
 seven-way algorithm, 24, 487
 Carroll, citations, 23
 centering, *see also* preprocessing
 centering
 additive adjustment, 111
 additive modeling, 111
 appropriate, 112, 119, 142
 average subject's profile, 131, 258
 biadditive modeling, 135
 bipolar scales, 122, 131, 360
 centers components, 120, 321
 centers errors, 120
 combinations of, 125
 comparing models, 129
 congruence coefficient, 196
 definition, 112
 degrees of freedom, 190
 double, 119, 131, 134, 142
 examples, 134
 similarities, 134
 effects of, 111, 120
 examples, 111, 115, 118, 196, 203, 314
 fiber, 113, 117, 119, 121, 127–128, 130, 132,
 138, 142, 222, 224, 298, 321, 386, 392
 advantages, 130
 appropriate, 118, 131
 combining centerings, 121
 content-based arguments, 118
 double, 119
 examples, 133, 314, 364, 395
 model-based arguments, 112
 model equivalence, 118
 moving centroid, 121
 removing offsets, 118, 119
 triple, 119
 growth data, 115
 incompatible normalization, 127

individual differences, 132
 interval scales, 117
 missing data, 153, 159
 model comparison, 121, 142, 204
 monopolar scales, 131, 360
 necessity in PCA, 114
 neutral midpoint, 113, 122
 normalization combined, 127
 offsets, 117
 removing, 118
 single overall, 122
 PCA, 110, 114, 121, 196
 precedes normalization, 127
 proposals, 119
 rating scales, 160, 194, 196, 353
 relationships, 119
 removing offsets, 321
 robust alternatives, 113, 123, 298
 scaling arrangement, 353
 slices, 119, 347
 examples, 386
 three-way analysis of variance models, 137
 three-way factorial designs, 119
 three-way rating scale data, 131
 triple, 119
 two-way analysis of variance, 134
 uncentered components, 120
 examples, 481

chapter structure, xviii

chemistry, xviii, 13
 ballast water example, 184
 chromatography example, 160
 commercial applications, 13
 discoloration of ham example, 473
 early reference three-mode analysis, 21
 enzymatic browning example, 73, 473
 hibiscus flowers example, 184
 improving core consistency, 186
 meat discoloration example, 73
 missing data, 148, 154
 multiway data, 470
 multiway multiblock models, 476
 plotting Parafac components, 267
 potential multiway analysis, 17
 robust PCA, 300
 sugar solution example, 5
 theoretical multiway models, 212
 unimodality constraints, 72
 chemometrics, 12–13, 24, 69
 Child's Report of Parent Behavior Inventory,
 CRPBI, 313, 315, 317, 320
 child studies
 attachment example, 405, 424
 coping example, 133, 241, 346, 354, 355

- child studies (*cont'd*)
 - education example, 4–6
 - intelligence example, 6, 59
 - learning-to-read example, 377
 - parental behavior example, 313
 - perceived-reality example, 36
 - physical growth example, 394
 - playing-with peers example, 454–455
 - problem behavior—CBCL, 356
 - showing respect example, 73
 - three-way questions, 5
- Chopin prelude study
 - content rating scales, 8
 - individual differences, 10
 - judgment technical aspects, 9
 - model selection, 191
 - prelude space, 9
 - research questions, 8
 - semantic space, 8
- chromatography
 - centering and normalization, 160
 - missing data, 154, 159, 168
 - multiple imputation, 161
 - multiway, 470
 - retention data, 160
 - retention rate patterns, 166–167
 - single dependent variable, 160
- circle of fifths, 11
- citations multiway papers, 13
- cluster analysis, 361
 - frontal slices core array, 351
 - grouping individuals, 404
 - k-means, 404
 - algorithm, 24
 - examples, 416
 - starting solutions, 417
 - overview techniques, 23
 - similarity measures, 401
 - simultaneous with component analysis, 404
 - three-way profile data, 404, 412
- columns
 - definition, 29
- combination-mode, 15
- component
 - meaning in Parafac, 212, 232
- component analysis, *see* multiway analysis
- component analysis, *see* principal component analysis
- component constraints, *see* model constraints
- component interpretation, 212, 315, 320, 369
- component matrices
 - comparison across modes, 220
 - comparison coefficients within a mode, 220
 - compensatory scaling core array, 219
 - comparability of centered and uncentered, 124
 - content-based scaling, 220
 - examples, 479, 366, 166, 316, 338, 367–368
 - first-order component scores, 223
 - first-order loading matrix, 222
 - functional constraints, 400
 - interpretation core array, 219–220
 - latent predictors, 222
 - loading interpretation, 219
 - metric for plotting, 220
 - no rotational freedom, 218
 - nonsingular transformations, 218
 - normalized components, 219–220
 - normalized coordinate scaling, 219
 - oblique, 218
 - orthogonal, 222–223
 - orthonormality, 218
 - principal coordinate scaling, 219–220
 - regression coefficients, 221
 - rotations, 220
 - scaling of components, 218
 - second-order component scores, 223
 - second-order loadings, 223
 - sign reversals, 319
 - standard coordinates, 219
 - standardized regression coefficients, 221–222
 - structured component scores, 221
 - structured loading matrix, 224
 - theoretical constructs, 219
 - unit-length components, 219
 - unit mean-square scaling, 219–220
- component uniqueness, 50
- component versus factor models, xvii
- components analysis, *see* principal component analysis
- compression, *see* Parafac algorithm, *compression*
- computer-intensive procedures, 187
- computer programs
 - FastMCD, 301
 - 3WayPack, xvii, xxi, 63, 73, 431, 488
 - ade4 package, 106
 - ALSCOMP3, 42, 74, 96, 321
 - CANDECOMP, 22, 24, 57, 487
 - CANDELINC, 94
 - CuBach, 488
 - GEPCAM, 101, 154, 162, 164
 - Latent Gold, 132
 - LIBRA Toolbox, 299
 - MICL, 174
 - MixClus2, 431
 - MixClus3, 431
 - multilinear engine, 96, 155, 173, 488
 - MultiMix3, 431
 - N-way Toolbox, 63, 73, 473, 488

- NORM, 174
- PTAk, 487–488
- SCA, 62–63
- Tensor Toolbox, 73, 488
- COMSTAT algorithm, *see also* TUCKALS
 - algorithms
- COMSTAT algorithm
 - Tucker3 model, 21
- configurations
 - sets of, 33, 39
- congruence coefficient, 196, 313, 326
 - between-model comparison, 196
 - definition, 196
 - examples, 196, 323–324, 328
 - rule of thumb, 196, 328
- conjugate gradient, 96
- consensus molecular alignment, 40
- constrained model, *see* model constraints
- CORCONDIA, *see* core consistency
- core array
 - anti-diagonal, 246
 - computation, 98
 - constraints, 69
 - diagonal and antidiagonal slices, 247
 - diagonality criteria, 248
 - examples, 253
 - different variants, 64
 - examples, 338–340, 367, 369, 390
 - explained variability of components, 66
 - matricized, 55
 - maximal simplicity, 246–247, 255
 - maximum number of zeroes, 246, 254
 - miniature data box, 228
 - Murakami core, 249, 254–255
 - examples, 255
 - Parafac model, 65
 - Parafac model applied to core array, 256
 - simplicity, 245–247
 - per-component plot, 271
 - slice diagonality, 246, 248
 - superdiagonality, 60, 65, 185, 245–248, 323
 - examples, 328–329, 362
 - orthogonality, 66
 - theoretical simplicity, 254–255
 - three-mode models, 64
 - three-way orthogonality, 56, 65, 179
 - formal proof, 66, 179
 - Tucker2 model, 64
 - Tucker3 model, 64
 - two-mode
 - simplicity, 245
 - two-way, 49
 - types, 246
- core consistency, 185
 - core consistency plot, 186
 - examples, 205–206, 329–330, 481
 - core consistency
 - dimensionality selection Parafac, 185
 - examples, 205, 327–329, 362
 - improving core consistency, 186
 - normalized, 185
 - core covariance matrix, *see* latent covariance matrix
 - correspondence analysis, *see* three-way correspondence analysis
 - correspondence analysis, *see* two-way correspondence analysis
 - covariance matrices
 - sets of, 31, 39, 47
 - Parafac2 model, 62
 - covariances vs correlations in multiway, 37, 124–125
 - covariation chart, 24, 469–470
 - crisscross regression, 148–149
 - comparison iterative OLS, 150
 - convergence, 149
 - deflation, 148–149
 - diverging missing data estimates, 149–150
 - local minima, 149–150
 - Parafac model, 154
 - robustness, 300
 - successive dyadic fitting, 149
 - three-way generalization, 154
 - triadic algorithm, 154
 - weighted, 149
 - cross-sectional data
 - versus repeated measures data, 31
 - cross-validation, 32, 341
 - CuBatch, 488
- D**
- data set
 - academic suitability data, 463–464
 - acrosol particles data, 480, 482–485
 - blue crab data, 409–411, 414–415, 417, 419, 422, 424, 432
 - chair-styles data, 262–263
 - Chopin prelude data, 8–11, 192–195, 197, 199–207
 - chromatography data, xxi, 159–160, 162, 164–168
 - coping data, 132–134, 240, 250–253, 255, 261–262, 355–359
 - electronics industries data, xxi, 282–283, 285, 290–295, 301–307
 - faces data, 296

- data set (*cont'd*)
- girls' growth curves data, 115, 258–259, 261, 264–266, 268, 270, 272, 274–280, 375, 394–398, 400–401
 - happiness data, xxi, 444–447, 450–451, 453–454
 - hospital data, 375, 385–392, 394
 - hostility data, 466–467
 - maize data, 174
 - multiple personality data, xxi, 350, 361, 363–371, 471, 476–479
 - NICHD data, 168–173
 - parental behavior data, 313, 316–319, 323–324, 326–328, 330–332, 334–335, 337–339, 341, 343
 - peer play data, 454–457
 - Strange Situation data, 425–432
 - sugar data, 144, 267, 269
 - university staff data, 461–462
 - world of woman data, 267
- degeneracy. *see* Parafac model
- degrees of freedom, 182
- biadditive model, 137
 - counting free parameters, 177
 - deviance plot, 181
 - maximum-product rule, 177
 - multiway component models, 176
 - multiway models
 - difficulties, 177
 - Parafac model, 177
 - rank of multiway arrays, 178
 - three-way interactions, 178
 - Tucker2 model, 177
 - Tucker3 model, 177
 - two-way models, 177
- demography
- mobility of work force example, 449
- derived data, 31
- deviance plot, 161, 176, 181
- between-model selection, 182
 - convex hull, 181, 183, 333
 - examples, 355, 396, 481
 - degrees of freedom, 181–182
 - examples, 163, 192–194, 333–334, 354–355, 362–363, 395–396
 - four-way examples, 476–477, 481–482
 - u*-criterion, 176, 182–183
 - algorithm, 182
 - comparison DiffFit procedure, 183
 - convex hull, 183
 - examples, 192–194, 355, 362–363, 396
 - four-way examples, 476, 481
 - smallest angle, 183
 - Tucker2 model
 - examples, 192
 - DiffFit procedure, 176, 179
 - comparison *u*-criterion, 183
 - eigenvalue-larger-than-1, 179
 - examples, 192, 194
 - four-way examples, 477
 - multiway scree plot, 179
 - salience value, 179
 - Tucker2 model
 - examples, 193
- dimensionality selection, 175
- bootstrap, 176
 - examples, 314, 327, 362
 - four-way examples, 478
 - jackknife, 176
 - large Parafac models, 184
 - minimum-product rule, 179
 - Parafac model, 184, 204
 - degree of superdiagonality, 185
 - examples, 192, 194–195, 323
 - split-half procedure, 184
 - split-half procedures, 325
 - Tucker model, 184, 203
 - examples, 192
 - Tucker versus Parafac models, 184, 362
- direct fitting, 61
- model hierarchy, 186
- direction cosines, 232, 383
- discrepancy function. *see* loss function
- display modes
- joint biplot, 273, 340
 - examples, 167, 274, 340, 357, 369, 389, 423
- dissimilarity matrices. *see* similarity matrices
- dot product, 500
- double centering. *see* centering
- dynamic factor models, 376, 401
- ## E
- ecology
- preprocessing, 130
 - unimodality constraints, 72
- ERG data
- component uniqueness, 61
 - four-way data, 472
 - multiway, 470
 - six-way data, 472
- Ehrenbreitstein, 470
- EM. *see* expectation-maximization
- environmental sciences, xviii
- aerosol particles data, 96, 471–472
 - air pollution, 471–472
 - environmental stress on blue crabs, 405, 409–410, 424
 - exception modeling example, 480

equal average diagonal normalization, 125
 equal average diagonal standardization, 37
 equilibrium circle, 262
 examples, 396, 398
 Escher's Man with Cuboid, vi
 Euclidean distance, 493
 Eve White–Eve Black, and Jane, *see* psychology
 event-related potentials
 explicit Parafac models, 64
 exchangeable, 17–18, 46, 376
 expectation–maximization algorithms, 80
 estimation, 189
 maximum likelihood, 80
 missing data estimation, 81
 exploratory factor analysis
 rotational indeterminacy, 88
 exploratory versus confirmatory models, xvii
 extended core array, 64
 and Tucker3 model, 55
 binary, 462
 diagonal
 Parafac model, 67
 examples, 339, 369, 371
 explained variability, 54
 nondiagonality, 53
 plotting
 examples, 391
 slice diagonality, 65, 246–247
 Parafac model, 65, 353
 symmetric slices, 232
 Tucker2 model, 53
 two-way orthogonality, 66
 extended three-mode models, *see* triadditive models
 external relationship analysis, *see* analysis of dependence
 external validation, xiv, 234, 322
 examples, 355
 external variables, 234, 312
 factorial design, 234
 handling external information, 234
 using functional information, 234
 using rank order information, 234
 using regression on components, 234
F
 FANOVA, factor analysis of variance, 135
 fiber centering, *see* centering
 fiber normalization, *see* normalization
 fibers
 definition, 29
 types of, 29
 fit/residual ratio, 202, 289, 327
 five-way data, 470

fixed configurations, *see* model constraints,
 external constraints
 flattening three-way data, 7, 12
 fMRI data
 four-way, 472
 four-way data, 473
 multiway, 470
 four-way core array, 231, 475
 four-way data, 16, 481
 aerosol particles data, 96, 472, 480
 agriculture, 5, 471
 analysis techniques, 7
 categorical, 455
 clustering, 404
 EEG example, 472
 excitation–emission matrices example, 472
 factorial data, 471–472
 fMRI example, 472
 longitudinal, 373
 modeling, 474
 multiple personality data, 350, 361, 471
 paucity, 469
 profile data, 35, 470–472, 474
 psychology, 5, 470
 rating scale data, 345, 471
 stimulus–response data, 470
 Tucker4 example, 476
 from three-way to two-way data, 45
 from two-mode to three-mode PCA, 44
 frontal slices, 29
 full core array, *see* core array
 fully crossed designs, *see also* three-way data,
 fully crossed
 fully crossed designs, 28, 31, 33, 35, 38, 43, 308,
 476
 binary data, 459
 common three-mode models for, 45
 compared to multiset designs, 43
 longitudinal data, 373
 fuzzy three-mode clustering, 23, 25

G
 GEMANOVA, 73, 141, 473
 general Euclidean models, 26
 general linear model, 32
 generalizability, 18, 32
 generalized Procrustes analysis, 25, 39–40, 51,
 196, 224
 multiple imputation, 171
 generalized subjective metrics model, *see*
 INDSCAL model
 GEPCAM
 description algorithm, 102
 missing data, 101, 154

GEPCAM (*cont'd*)

- quality solutions, 164
- regression-based algorithm, 101, 154
- robust regression, 101

Gojoechea on multiway analysis, 17

Gower on multiway analysis, 17

Gram–Schmidt orthonormalization, 100

H

Harris–Kaiser independent cluster rotation, *see* rotation

HICLAS, *see* hierarchical classes models

hierarchical classes models, 25, 42, 459, 467

- algorithms, 465

- association rule, 462

- conjunctive, 465

- disjunctive, 463

- binary component matrices, 463

- binary data, 459, 467

- binary extended core array, 463

- Boolean algebra, 460

- component matrices

- examples, 464

- equivalence relationships

- examples, 462–463

- examples, 460

- hierarchical relationships

- examples, 461, 463

- identifiability HICLAS models, 465

- model comparisons, 465

- Parafac–HICLAS model, 465–466

- references, 459

- simulation studies, 465

- Tucker2–HICLAS model, 460, 462

- examples, 461

- Tucker3–HICLAS model, 460, 464, 466

- examples, 466–467

- Tucker3 vs. Tucker2–HICLAS model, 464

higher-order component models, 22, 231

higher-order singular value decomposition, 50

higher-way analysis

- references, 487

higher-way data, 5, 16, 469

- analysis techniques, 7

- categorical, 473

- examples, 471

- factorial designs, 473

- interactions, 473

- interpretation, 470

- longitudinal, 373

- multiway clustering, 404

- origin, 469

- paucity, 469

- rating scales, 345

Tuckers models, 487

history

- book Smilde et al., 12

- Carroll, 12

- chemistry, 12

- Chrysler, 12

- citation analysis, 12

- chemometricians, 12

- founding fathers, 12

- psychometricians, 12

- early Tucker3 and Parafac models, 21

- expansion multiway analysis, 12

- factor analysis terminology, 21

- Harshman, 12

- independent proposal Tucker model, 103

- individual differences, 12

- Jordan, Hitchcock, and Oldenburger, 20

- linear algebra focus, 20–21

- Osgood, 12

- three-mode analysis, 6, 12

- three-way data, 24

- Tucker, 12, 21

horizontal slices, 29

horticulture

- taxonomy of roses example, 106

hypothesis testing, 7, 20, 31

- biadditive model, 137

- complexity multivariate data, 32

- descriptive use, 32

- repeated measures, 375

- structural equation modeling, 385

- three-way interactions, 178

I

idealized subject, *see* interpretation

INDSCAL model, 40

- INDSCAL relative, 23

implicit theories of personality

- three-mode analysis with constraints, 73

INDCLAS, *see* hierarchical classes models

INDCLUS model, *see* individual differences

- clustering

independence of observations, 46

- mixture method of clustering, 408, 432

indirect fitting, 26, 39, 61

- model hierarchy, 186

individual differences clustering, 23, 26, 40, 404

individual differences scaling, 23, 26, 40, 232

- linear constraints, 70

- normalization, 134

- preprocessing, 132

individual differences

- centering, 132

- concept scores, 349

- concept usage, 348
- dissimilarity data, 40
- examples, 355
- extended core array, 351–352
- extent of, 349
- in three-way analysis, 12
- many subjects, 361
- model selection, 193, 208
- multiway models, 18, 312
- number of components, 349
- Parafac model, 353
- preference rank order data, 40
- preprocessing, 133
- rating scales, 350
- raw versus derived data, 31
- response style, 347
- scale loadings, 349
- scale usage, 348
- scales and loadings, 348–350
- similarity data, 40
- subjective judgements, 350
- three-mode models, 350–351
- three-way rating scale data, 346
- Tucker2 model, 351
- Tucker3 model, 352
- types, 346
 - versus sampling from populations, 17, 211
- INDSCAL model, *see* individual differences
 - scaling
- influential points, 284–285
- inner product, 494, 500
- instrumental variables, 361
- intelligence tests
 - different age groups, 172
 - nonnegative components, 70, 95
- inter-battery method of factor analysis, 313
- interactive coding, 276–278, 434, 441, 449, 452, 455, 471, 475
- inferred relationship analysis, *see* analysis of interdependence
- internal validation, xiv, 322
 - assessing fit, 233
 - examples, 342, 487
 - stability parameters, 233
- interpretation
 - aids for subject mode, 234
 - angles of vectors, 9
 - asymmetric treatment of modes, 211
 - complexity, 470
 - component models, 209
 - component uniqueness, 212
 - components and core array, 229
 - conditional statements, 231
 - normalized coordinates, 231
 - content-based, 210
 - core array, 213
 - complexity, 470
 - examples, 367, 390, 484–485
 - latent covariances, 225
 - percentages of explained variation, 225, 227
 - regression weights, 225–227
 - scaling components, 225
 - sources of idealized or latent elements, 225
 - size elements, 238
 - strength link between components, 225
 - three way interaction measures, 225
 - weight component combination, 225–226
 - Euclidean distances, 243
 - extended core array, 225, 231
 - direction cosines, 225, 232
 - full core array, 225
 - graphical displays, 210
 - idealized subject, 213, 228–229
 - keeping track of signs, 230–231
 - lack of rotational freedom, 214
 - latent variables, 213
 - meaning for all components, 229–230
 - model based, 210
 - multidimensional point of view, 210–211
 - low-dimensional subspaces, 211
 - multivariate point of view, 210
 - role random samples, 211
 - Tucker versus Parafac models, 211
- Murakami form
 - extended core array, 231
- Parafac component weights, 225
- Parafac components, 212
- Parafac model, 212–214, 315
 - lack of solution, 214
- prototype conditions, 213, 228
- rotations, 210, 213
 - effect on core array, 213,
 - scaling of components, 210
 - scaling of core arrays, 210
- statistical meaning parameters, 209
- subspace interpretation, 211
- symmetric treatment of modes, 211
- time trends, 213
- transformations, 210
- Tucker model, 212–213, 315
 - availability solution, 214
- Tucker versus Parafac components, 210, 214
- Tucker3 model, 224
- ultrinsic axis property, 58, 212
- invariant factor analysis, 26
- isometric, 272, 274

iterative algorithms

- convergence, 81
- divergence, 81
- global maximum, 81
- local maxima, 81
- local optima, 82
- monotone convergence, 81
- multiple solutions, 81–82
- nonconvergence, 81
- number of iterations, 82–83
- overrelaxation, 83
- slow convergence, 82–83
- starting solutions, 82
- step size, 82–83
- stopping criterion, 81–82

iterative ordinary least squares, 150

iterative proportional fitting, 125

J

jackknife, 18, 176

- basics, 188
- examples, 207
- expectation-maximization estimation, 189
- implementation methods, 189
- multiway analysis, 189
- outliers, 189
- predictive power, 188–189
- stability estimation, 189
- stochastics, 188

joint biplot, *see* plottingjoint plot, *see* joint biplot**L**

latent class model, 25

- nonnegativity components, 70

latent covariance matrix, 378, 381–382, 384, 401

- assessing change, 392
- assessment stability, 393
- assessment stationarity, 384, 393
- autoregressive process, 383–384
- direction cosines, 393
- examples, 393–394
- homogeneity, 384
- identity matrix, 385
- interpretation, 383
- normalization, 383, 393
- null model, 385
- structural equation modeling, 382
- Tucker2 model, 381, 383
- uniform autocorrelation, 385

latent three-mode path models, 401

lateral slices, 29

least-squares loss function, 79–80

level-fit plot, *see* sums-of-squares plot

levels, definition, 28

LIBRA robustness toolbox, 299

Likert scales, 346, 360

linked-mode analysis, 173

loadings, *see* principal component analysis

loadings

- usage term, 48

loglinear analysis, 42, 434

- logits, 440

loss function, 19

- minimization, 79

M

marketing

- multiway rating scale data, 346
- preference rank orders, 40
- repetitive advertising, 24

mathematics, 13

MATLAB, xvii, 63, 73, 299, 473, 488

matricization, 7

- Tucker Method I, 98

matrix-conditional, 42

maximum-product rule, 98, 103, 177

maximum likelihood clustering, 23

maximum likelihood Parafac model, 159

maximum likelihood PCA, 151–152

- missing data, 151
- probabilistic PCA, 151
- two-step procedure, 151

maximum variance components, 211–212, 228,

321, 332

- interpretation, 211

- preprocessing, 118

means plot, *see* mixture method of clustering

medicine, 13

- EEG example, 344

- fMRI example, 5, 344

methods and models chart, 25–26

minimization

- difference structural image and data, 79
- loss function, 80

minimum product rule, 56, 179

- orthogonality core array, 66

- PRESS, 190

minimum spanning tree, *see* plotting

missing data, 143

- analysis for estimation, 144, 157

- analysis in spite of, 144, 157

- analytical chemistry, 144

- at random, 145

- bootstrap procedures, 153, 159

- comparative Parafac analyses, 154

- comparison EM and crisscross, 155
- comparison Tucker3 algorithms, 159, 161, 163–165
- completely at random, 145
- completely missing fibers, 146
- completely missing slices, 146
 - smoothness constraints, 146
- computer programs, 154
- confidence intervals, 150, 164
- crisscross regression, 148, 150
- equivalence EM and iterative OLS, 150
- estimation, 104, 159
 - examples, 163, 165
 - model dependence, 158
 - multiplicative model, 157
 - triadditive model, 157
 - with preprocessing, 157
 - without preprocessing, 157
- expectation–maximization methods, 147–148, 150, 158, 170
 - reduction variability, 147
- full-information methods, 147
- imputing nonstochastic data, 147
- iterative OLS and crisscross algorithms, 150
- Levenberg–Marquardt algorithm, 155
- linked mode data, 174
- maximum likelihood methods, 147–148, 159
- maximum likelihood PCA, 153
- missing at random, 147
- missing fibers, 146
- model comparison, 158
- multiple imputation, 147–148, 152, 157–158
 - error reduction, 147
 - examples, 171–173
 - limited experience multiway, 148
 - nonstochastic data, 147, 177
- multivariable–multioccasion matrix, 146
- multiway data, 154, 157
- necessity of one-step procedure, 153
- nonconvergence, 161, 164, 168
 - examples, 481
- not at random, 145
- optimal scaling, 148, 152
- outliers, 144
- planned missing, 145, 173
 - examples, 145
- preprocessing, 148
 - PCA, 148
- random variables, 146
- richness multiway data, 145
- single imputation, 150, 158
 - overfitting, 152, 157
- starting values, 146–147, 149–150, 481
- test equating, 173
- two-mode PCA, 147–148
- two and three-way patterns, 145
- types of missing data
 - examples, 174
- mixed multiplicative model, 140
- complexity interpretation, 140
- constant components, 141
- estimation, 141
- identification constraints, 141
- Parafac model, 141
- Tucker3 model, 140
- mixture method of clustering
 - allocation to groups, 419
 - categorical response variables, 405
 - common correlation matrix, 421
 - examples, 421–422
 - common covariance matrix, 406, 412
 - diagonal, 413
 - examples, 410, 413, 416, 426–427
- computer programs
 - Latent Gold, 432
 - MixClus2, 431
 - MixClus3, 431
 - MultiMix3, 431
- correct allocation rate, 419
 - examples, 420, 426
- covariances, 405
- distributional assumptions, 404, 408
- examples, 410, 424, 432
- exploratory use, 408
- group-specific covariance matrices, 406, 412–413, 421, 430
 - examples, 413, 416, 421, 425, 427–428
- independence assumption, 408, 432
- lack of hierarchy, 418
- likelihood function, 407, 410, 415–416
- likelihood ratio test
 - examples, 426
- local independence, 413, 431
- local maxima, 407, 415–416
- maximum likelihood estimation, 407
- means, 405
 - examples, 410
 - three-way array, 420
- means plot, 420
 - examples, 411, 420–421, 427
 - standard errors, 421
- missing data, 430
- mixing proportions, 407
- multivariate multinomial, 405, 414, 430
 - examples, 414–415
- multivariate normality, 405–406, 413–414, 432
- nesting of solutions, 418
 - examples, 419, 426

- mixture method of clustering (*cont'd*)
 - normality, 407
 - number of groups, 407, 413, 415
 - cross-classification, 418, 426
 - examples, 416, 418
 - likelihood ratio test, 415–417
 - measures of agreement, 418,
 - posterior probabilities, 418
 - number of parameters, 412–413
 - examples, 413, 417
 - posterior probabilities, 408, 419
 - examples, 420, 426
 - references, 432
 - sample size, 413
 - singular covariance matrices, 412
 - starting allocations, 410, 416
 - examples, 416, 425
 - hierarchical clustering, 407
 - k-means, 416
 - two-mode clustering, 407
 - two-mode mixture method, 407
 - Ward's method, 407, 416
 - starting solutions, 407
 - statistical model, 20
 - theoretical background, 432
 - three-mode, 23, 25, 35, 404–405, 411, 415, 424–425, 432
 - distributions, 406
 - group structure, 407
 - two-mode, 404, 425
 - validation
 - discriminant analysis, 422–423, 429
 - external information, 424–430
 - linear discriminant analysis, 422–423
 - ordination, 408, 422–423, 432
 - quadratic discriminant analysis, 422–423
- mode, xvii, 16
 - definition, 28
- model, use of term, 18
- model and dimensionality selection, 161, 170, 192, 314, 323, 333, 354, 362, 387, 395, 446, 449, 455, 466, 476, 481
- model and number-of-clusters selection, 410, 415, 425
- model constraints
 - constant components, 72
 - equality constraints, 74
 - external configurations, 73
 - examples, 366–367
 - external variables, 72
 - factorial regression, 73
 - functional constraints, 72
 - linear constraints, 70
 - merging incompatible data sets, 73
 - missing data, 74
 - nonnegativity, 70
 - optimal scaling, 74
 - order constraints and autocorrelated data, 73
 - order constraints and clustering, 74
 - orthogonality, 69
 - overview, 68–69
 - smoothness, 71
 - splines, 71
 - unimodality, 72
- model hierarchy, 21, 186
 - between-model selection, 66, 186–187
 - direct fitting, 186
 - examples, 206
 - French proposals, 186
 - fully crossed data, 67
 - indirect fitting, 186
 - mathematical details, 186
 - three-mode models, 67–68
 - Tucker2 model, 186
 - Tucker3 model, 186
- model matrix, *see* structural image
- model selection, *see also* between-model selection
- model selection, *see also* model hierarchy
- model selection, 66, 175
 - chemistry, 175
 - content-based arguments, 208, 333
 - criteria, 175
 - example and model hierarchy, 68
 - examples, 190, 208, 333–334
 - jackknife, 189
 - PRESS criterion, 190–191
 - restrictions on core arrays, 67
 - Tucker versus Parafac, 334
- monopolar scales
 - centering, 131, 360
 - multiway factorial data, 131
 - preprocessing, 129
 - three-way rating scale data, 346, 360
 - triadditive modeling, 139
- more-sets multiway data, 487
- multiblock multiple regression, 25
- multidimensional point of view, 214, 217, 221, 235
 - rotation, 239
- multidimensional scaling, 361
 - preference rank orders, 40
 - types of designs, 33
- multilevel latent class model, 405
- multilinear, 16
- multilinear engine, 96, 155, 173, 488
- multimode, 16
- multimode analysis, *see* multiway analysis
- multimode covariance matrix, 22, 26, 33, 37, 47, 376–377, 382, 385

- algorithm, 88, 99–100, 299
- confirmatory, 238
- exploratory analysis, 37
- latent covariance matrix, 382
- missing data, 155
- size for SEM, 379
- structural equation modeling, 385
- three-mode methods, 385
- multioccasion–multisubject data, 38
- multiple imputation, *see also* missing data
- multiple imputation, 152, 156, 168
 - absence stochastic mode, 152
 - between-imputation standard errors, 153
 - between-imputation variance, 152, 171
 - choice of matrixization, 156
 - comparing PCA bootstrap solutions, 154
 - comparing PCA solutions, 153
 - computer programs, 174
 - MICE, 174
 - NORM, 174
 - error estimates, 157
 - estimate out of range, 157, 169
 - estimates missing data, 157
 - expectation–maximization methods, 172
 - generalized Procrustes analysis, 171
 - matrixizing three-way data, 156
 - multiway analysis, 158
- NICHD
 - comparison of solutions, 171–172
 - fit measures, 170
- PCA, 153
- tall combination-mode matrix, 156
- wide combination-mode matrix, 156, 169
- within-imputation standard errors, 153
- within-imputation variance, 152, 171
- multiplicative adjustment, *see* normalization
- multiplicative modeling
 - two-way, 111
- multisample data, 33, 38
- multiset canonical correlation analysis, 25, 27
- multiset data, 33, 38–39
 - compared to fully crossed data, 43
 - conversion to three-way data, 43
 - Parafac2 model, 62
 - residuals, 286
- multisubject–multisample data, 38
- multivariable–multioccasion matrix, 37, 47, 376
- multivariate data
 - heteroscedasticity, 286
- multivariate kurtosis, 414
- multivariate longitudinal analysis, *see also* repeated measures analysis
- multivariate longitudinal analysis, 25
 - autocorrelation, 378
 - autoregression, 377
 - autoregressive models, 376
 - constraints on components, 377
 - descriptive modeling, 374
 - descriptive multimode covariance modeling, 377
 - dynamic factor analysis, 376
 - exploratory modeling, 375
 - functional constraints, 377
 - latent growth modeling, 377
 - latent structures, 374
 - multiway models, 374
 - role of time, 374
 - rotation of components, 378
 - smoothness constraints, 377
 - splines, 377
 - stochastic modeling, 374
 - structural equation modeling, 376
 - three-mode component analysis, 378
 - time as interpretational device, 374, 378
 - trajectories, 378
 - variable and serial dependence
 - confounding, 378
 - without latent structures, 374
- multivariate longitudinal data, 33–34, 36, 373, 385, 401
 - serial dependence, 378
 - variable and serial dependence
 - three-mode PCA, 378
 - variable dependence, 378
- multivariate multiway analysis of variance
 - profile data, 133
- multivariate one-way design, 135
- multivariate point of view, 214–215, 221, 224, 235
 - Parafac model, 212
 - PCA loadings and scores, 211
 - rotation, 239
- multivariate two-way factorial data, 138
- multiway, *see also* three-way
- multiway algorithms, 24
 - alternating least squares, 80
 - explicit algebraic solutions, 84
 - general characteristics, 84
 - lack of nested solutions, 84
 - number of components, 84
 - simultaneous solution for components, 84
- multiway analysis, xviii, 3, 16
 - a scientific necessity, 3
 - algorithms, 82
 - analysis of residuals, 283
 - bootstrap, 233
 - stochastics, 233
 - chemometrics, 24
 - choice of models and dimensionality, 20
 - component analysis, 25

- multiway analysis (*cont'd*)
 - confirmatory, 233
 - counting ways, 4, 488
 - data-analytic models, 20
 - dealing with complex questions, 6
 - design on subject mode, 70
 - direct and indirect fitting, 61
 - distortion component space, 283
 - effects of outliers, 283
 - exposure, 281, 283
 - facet designs, 70
 - factorial designs, 70
 - fitting systematic variability, 184
 - generalizing three-way, 24
 - higher-way data, 471
 - history, 6, 13
 - hyperboxes, 5
 - increase publications, 13
 - index-based notation, 488
 - individual differences, 17, 31
 - individual differences scaling, 23
 - interpretation, 18
 - iterative algorithms, 80
 - jackknife, 189
 - lack of sampling framework, 17
 - models versus methods, 20
 - notation, 487
 - outliers, 283
 - overview data types, 27
 - preprocessing, 113
 - presentation of results, 321
 - quality of solution, 287
 - rank, 21
 - reduction in parameters, 18
 - research questions, 5, 17
 - robustness, 282
 - role of physical models, 19
 - sample size, 176
 - summarization, 281, 283
- multiway arrays, 16
 - rank, 178
- multiway binary data, 33, 41
- multiway block models, 24
- multiway categorical data, 33, 433
- multiway contingency tables, 33, 41–42, 434
 - interactions, 434
- multiway data, xvii–xviii, 1, 3, 16–17, 33
- multiway data designs, 33
- multiway data
 - designs, 17
 - fully crossed, 470
 - nonstochastic, 161
 - origin, 469
- multiway factorial data, 33, 35
 - agricultural example, 36
 - chromatography, 160
 - coping data, 133
 - cross-cultural example, 471
 - discoloration of ham example, 473
 - enzymatic browning example, 473
 - five-way, 473
 - perceived-reality example, 36
 - preprocessing, 129–130
 - preprocessing recommendations, 142
 - shifted multiplicative model, 122
 - single observation per cell, 111
 - six-way, 473
 - spectroscopy example, 472
- multiway loglinear modeling, 42
- multiway longitudinal data, 168
- multiway multiblock models, 25, 27, 32, 476, 487
- multiway ordinal data, 33
- multiway partial least-squares models, 25
- multiway profile data, 35, 133, 168, 312, 320, 331
 - analysis methods, 25
 - degrees of freedom, 177
 - examples, 258
 - individual differences, 132, 311
 - modeling, 311, 320
 - preprocessing, 312, 321
 - preprocessing recommendation, 142
 - statistical models, 404
 - stochastics, 177, 184, 311–312, 404
 - triadditive modeling, 139
- multiway rating scale data, *see also* three-way rating scale data
- multiway rating scale data, 33, 312, 345
 - analysis methods, 25
 - degrees of freedom, 177
 - fiber normalization, 131
 - frontal slice normalization, 131–132
 - lateral slice normalization, 131–132
 - normalization, 131
 - preprocessing, 312
 - profile arrangement, 35
 - triadditive modeling, 139
- multiway scree plot, 176, 179–180, 355
 - alternating least squares, 180
 - between-model selection, 182
 - compared to deviance plot, 181
 - convex hull, 180
 - examples, 180, 192–193, 195, 333, 335, 362
 - four-way examples, 476–477
 - Parafac model, 182
 - examples, 326
 - Hucker Method I, 180
- Murakami form
 - extended core array, 231

Tucker2 model, 222–223

Tucker3 model, 224, 228

music

classical, 8, 191

judging Chopin's preludes, 7, 191, 346

N

N-way Toolbox, 63, 73, 473, 488

nested designs, 33, 38

nested-mode biplot, *see* plotting

nested-mode per-component plot, *see* plotting

Nuschwanstein, 470

NICHD data

background black families, 168

preprocessing, 169

nonlinear canonical correlation analysis, 25

nonsymmetric correspondence analysis, 433

norm of a matrix, 494

normalization, *see also* preprocessing

normalization, 115–116, 124

appropriate, 112

centering combined, 127

normalization/components, *see* component

matrices

definition, 112

degrees of freedom, 190

effects of, 111

equal average diagonals, 125

equalizing range of variables, 113, 127

examples, 116

fiber, 112, 124–126, 129–132, 142, 348, 392–393

arguments in favor, 126

model incompatibility, 124

standard scores, 124

four-way, 474

homoscedasticity of error terms, 116

incompatible centering, 127

latent covariance matrix, 383

missing data, 153

model-based arguments, 116, 123–124

multiplicative adjustment, 111

multiway rating scales, 131

necessity in PCA, 115

none, advantage, 126

overall, 129

PCA, 110

preceded by centering, 127

ratio scales, 160

recommendations, 142

response styles, 123

robust alternatives, 113, 298

slice, 112–113, 117, 124–125, 127–128, 142, 169, 298, 321, 392

advantage double normalization, 125

examples, 314, 386, 395, 481

incompatibility double normalization, 125

model compatibility, 124

orthogonal, 125

per scale, 131–132

per subject, 131–132

per variable, 133

similarity matrices, 134

simultaneous normalizations, 125

sums-of-squares plot, 292, 336

variables same ranges, 362

normalized coordinates, 441

notation, 27, 490

Feynman-like diagrams, 487

index-based, 438

multiway Kronecker, 487

standardization of terminology, 488

numerical convex-hull-based selection method, *see* deviance plot

nursing

relational communication example, 134

O

object variation, 60, 332

oblique-components plot, *see* plotting

offset(s), *see* centering

optical physics, xviii

optimal scaling, 42

measurement levels, 42, 321

multiway analysis, 74, 434

optimization algorithms, 80

ordinal multiway data, 42

ordination, *see also* mixture method of clustering

ordination, *see also* principal component analysis

ordination, 35, 129–130

cluster means, 420

cluster validation, 412, 422, 432

examples, 423

clustering, 404, 408, 422

outliers, 281–283

accommodation, 282

cause of three-way interactions, 123

component analysis, 284

preprocessing, 283

probability plot, 280

Rao distance, 285

residual sum of squares, 285

testing for discordance, 282

P

pairings

multiway analysis, 3

paired-components plot, *see* plotting

Parafac algorithms

Aitken extrapolation, 86

ALS/COMP3, 96

alternating least squares, 84

basic alternating least squares, 86–87

compression with Tucker model, 101, 103

convergence, 94

examples, 325–327

crisscross algorithm, 86

evaluation of algorithms, 86, 88

example acceleration, 92

examples, 93

linear constraints, 94

local minima, 83

multimode covariance matrix algorithm, 87

multiple solutions, 92

nonconvergence, 61, 83

nonnegativity constraints, 96

optimal scaling, 96

overrelaxation, 86

sensitivity to local optima, 85

simultaneous matrix diagonalization, 88

slow convergence, 86

starting solutions, 86–87

successive linear regressions, 86

triadic crisscross algorithm, 86

uniqueness, 88

uniqueness and applications, 88

Parafac core array, 65, 185, 232

calculation, 65

examples, 205, 329, 332

Tucker structure, 338

Parafac model, 22, 59–60, 85–86

a posteriori clusters, 71

a priori clusters, 70

aerosol particles example, 472

as Tucker2 model, 54, 353

basic form, 60, 215, 218

chemistry, 22

chromatography example, 475

common factor model, 62

component interpretation, 322

component uniqueness, 50, 61

constant components and ANOVA, 73

constant correlations, 58, 313

constraints, 58, 94

contrast components, 95

definition, 57

degeneracy, 59, 61, 83, 89, 214, 232

advice, 92

bounded, 90, 94, 362

constraints, 90

divergent, 90, 204

examples, 91, 194, 204, 325, 362

indicators, 61, 90–91, 185, 204, 362

interpretation, 90

temporary, 90–91

three-way rating scale data, 353

triple cosine product, 90, 204

Tucker variation, 90–91

degrees of freedom, 177

dimensionality selection, 184, 204

core consistency, 185

degeneracy, 185

local minima, 185

early reference, 21

EEG example, 344

equivalence Tucker3 model, 256

evaluating fit, 233

explanatory, 85

explicit algebraic solution, 86

explicit physical models, 61

fitting maximal dimensionality, 184

fitting raw data, 62

fMRI example, 344, 473

four-way, 472, 474, 481

algorithms, 96

component interpretation, 474

fMRI example, 473

orthogonality, 474

uniqueness, 474

functional constraints, 72

generalization singular value decomposition, 61

Harshman's diagnostics, 184

HICLAS variant, 460

INDSCAL relative, 23

interdependence technique, 25

intrinsic axis property, 61, 213

lack of rotational freedom, 58, 214

less terms than Tucker3 model, 59

matrix notation, 59

missing data, 148

mixed multiplicative model, 141

model hierarchy, 68

modeling interactions multiplicatively, 73

modes equal number of components, 60

more components than levels, 473, 481

more restricted than Tucker model, 59, 84

examples, 481

multiway profile data, 313

multiway rating scale data, 196

multiway uniqueness, 475

nonnegative components, 70

nonorthogonality of components, 60

not a decomposition model, 59

not a one-component Tucker3 model, 57

number of components, 58, 213, 353

- examples, 324
- object variation, 60
 - indirect fitting, 61–62
- optimal scaling, 42, 74, 86, 96, 321
- orthogonality, 66, 69, 95, 206, 232
- orthogonality and degeneracy, 70
- orthogonality and model hierarchy, 68
- orthogonality and partitioning variability, 66
- orthogonality constraints, 62, 196, 206
 - examples, 194, 314, 323–324, 329
- Parafac core with Tucker variation, 65
- parallel proportional profiles, 57
- parallel solution, 89, 323
 - examples, 323–324
- partitioning of variability, 215, 227, 233
- per-component plot, 270
- practical aspects, 322
- precursors, 21
- preprocessing, 196
 - examples, 194
- restrictive nature, 84, 194, 212, 214, 218, 323, 472
- role of physical models, 22
- sets of covariance matrices, 62
- seven-way, 487
- single interpretation per component, 59
- smoothness constraints, 71
- speed of convergence
 - examples, 204
- structured two-mode PCA, 221
- suitability multiway rating scale data, 194, 196
- sum notation, 60
- sum of rank-one arrays, 59, 69
- superdiagonal core array, 60, 66, 245
 - three way orthogonality, 246
- swamps, 80
- symmetric model, 60
- trilinear, 221, 328
- Tucker structure, 214, 329, 338, 353
 - examples, 329
- tutorial, 22
- two-component degeneracy, 91
- two-component equivalence Tucker3 model, 158, 197, 334
- uniqueness, 13, 212
- uniqueness and degeneracy, 88
- uniqueness and interpretation, 212, 320
- uniqueness
 - examples, 324
 - necessary conditions, 89
 - strong, 89, 323
 - sufficient conditions, 89
 - weak, 88, 323
- unit mean square components, 220
- univariate three way ANOVA, 73
- unsuitability multiway rating scale data, 198
- versus CANDECOMP, 22
- versus Tucker3 model, 213
- Parafac2 model, 39–40, 62–63, 232
 - component uniqueness, 63
 - constant correlations, 62
 - constraint variants, 63
 - definition, 62
 - equal covariances simplification, 63
 - extension classical PCA, 62
 - four-way, 475
 - fully crossed data, 62
 - less restrictive than Parafac model, 62
 - longitudinal multiset data, 62
 - orthogonality constraints, 63
 - relation with Parafac model, 64
 - three-way, 475
- parallel factor analysis, *see* Parafac model
- parallel factor model, *see* Parafac model
- parallel proportional profiles, 61–62, 84, 184, 203, 214, 270, 313, 322–324, 329, 474
 - intuitive appeal, 58
 - orthogonality constraint, 69
 - reality of components, 58
 - simplicity of components, 58
- Paratuck2 model, 63–64
- partitioning variability
 - core array, 54, 56, 215, 226–227, 233, 390
 - examples, 336, 397
 - examples, 198, 364
 - orthogonality, 69
 - per component, 54, 56, 215, 227, 233, 387, 428
 - examples, 170, 335, 363, 397–398, 481–484
 - four-way examples, 478
 - per individual data point, 243
 - per joint biplot axis, 273
 - examples, 357
 - per level of a mode, 54, 56, 198, 200–204, 215, 233, 363–364
 - examples, 315, 336
 - residuals, 285
- PCASup, 68
- Pearson χ^2 statistic, 435
- Pearson mean-square contingency coefficient, 435
- per-component plot, *see* plotting
- pharmacology
 - cocaine molecules example, 5
 - consensus molecular alignment example, 39
- physical models, 13, 212
- pipes, definition, 29
- plotting
 - across-modes comparisons, 271

plotting (*cont'd*)

- adding explanatory information, 262

- examples, 261–263

- adding external variable, 262

- adding group information, 262

- all-components plot, 266, 322

- examples, 266–269, 387, 398

- auxiliary orthonormal basis, 269

- coordinate spaces, 258

- different modes together, 270

- distance form PCA, 259

- extended core array

- examples, 391

- higher-dimensional spaces, 262

- anaglyphs, 266

- anchoring of points, 265

- dynamic graphics, 262, 266

- examples, 262, 265–268

- minimum spanning tree, 262, 265

- paired-components plot, 262

- sizes of points, 265

- stereopairs, 266

- three-dimensional graph, 265

- joint biplot, 273, 340, 423

- advantages, 274

- and nested-mode biplot, 443, 278

- construction, 273–274

- correspondence analysis, 435

- examples, 9, 167, 274–276, 336, 340–341, 356–359, 369–370, 389, 416, 423–424, 428–431

- joint biplot axes, 273

- three-mode C/A, 440

- unequal spread, 454

- minimum spanning tree

- examples, 264, 396

- nested-mode biplot, 276–278, 399, 402

- construction, 276

- correspondence analysis, 435

- covariance form, 277

- distance form, 277

- examples, 277–279, 399, 447

- origin, 277

- three-mode C/A, 440–441

- total dependence, 442

- trajectories, 277–278, 441–442, 447, 457

- nested-mode per-component plot, 278, 280, 399, 402

- examples, 399–400

- normalized versus principal coordinates, 260, 482

- oblique components plot, 269

- examples, 270, 329, 331

- paired-components plot, 133, 167, 260, 321–322

- examples, 10–11, 166, 264, 329, 355–356, 364–365, 367, 456, 483–484

- origin, 133, 258, 278, 357, 364, 400

- Parafac components, 269

- per-component plot, 312, 220, 271, 322

- examples, 271–272, 316–319

- principal coordinates, 482

- principal versus normalized coordinates, 260, 482

- examples, 261

- single modes, 260

- three-dimensional spaces

- minimum spanning tree, 263

- two-mode biplot, 271, 273

- polyadics, 20

- predictive power, 188

- basics, 190

- examples, 191, 207–208

- jackknife, 188

- model selection, 190

- preprocessing, 190

- PRESS, 190

- W-statistic, 190

- predictive validity, 176

- preference data

- analysis methods, 26

- preference mapping, 361

- preference rank orders, 33, 40

- preprocessing, *see also* centering

- preprocessing, *see also* normalization

- preprocessing

- arbitrary means and variances, 128

- arbitrary variances, 128

- centering plus normalization, 127

- centering precedes normalization, 127

- comparable means and variances per condition, 129

- comparing different preprocessing, 129

- comparison between data sets, 116

- content-based arguments, 116–117

- definition, 109

- fiber normalization, 129

- growth curves, 115

- incomparable means and variances, 129

- interpretable means and variances, 130

- interval scales, 111, 128

- iterative centering and normalization, 127

- longitudinal example, 116

- missing data, 128

- model-based arguments, 116–117

- monopolar scales, 129

- multiway data, 112

- multiway factorial data, 111, 129

- necessity, 114

- neutral midpoints scale, 128
 - one-step procedure, 118, 128
 - overall normalization, 129
 - perpendicular centering and normalization, 127
 - profile preprocessing, 113, 130, 142, 283, 312, 314, 321, 332
 - recommendations, 141
 - similarity matrices, 134
 - slice centering, 129
 - summary recommendations, 112, 117
 - three-way data
 - number of means, 112
 - number of normalizations, 112
 - three-way profile data, 130
 - two-step procedure, 117–118, 128
 - two-way data, 110
 - two-way profile data, 110
 - type of arguments, 112
 - PRESS, *see* predictive power
 - principal component analysis, xviii–19
 - bilinear model, 217
 - centering, 110
 - coefficients as generic term, 217
 - component loadings, 216
 - component scores, 216
 - coordinates as generic term, 217
 - covariance form, 215–216, 222, 241, 259
 - distance form, 216, 259
 - examples, 410
 - Hotelling's geometrical view, 44
 - loadings, 19
 - multiplicative vs additive terms, 48
 - normalization, 110
 - normalized coordinates, 216, 259
 - Pearson's statistical view, 44
 - preprocessing + model fitting, 110
 - preservation Euclidean distances, 260
 - principal coordinates, 216, 259
 - principal component analysis
 - principal coordinates, 411
 - principal component analysis
 - projecting into a lower-dimensional space, 44
 - reduction to a limited number of composites, 44
 - robustness, 282
 - rotational indeterminacy, 88
 - scores, 19
 - standardized regression coefficients, 216
 - sum of rank-one matrices, 49
 - symmetric view of coefficients, 217
 - variable-component correlations, 216
 - principal component residuals, *see* residuals
 - principal coordinates, *see* principal component analysis
 - probabilistic PCA, 151
 - profile arrangement, *see* multiway rating scale data
 - profile preprocessing, *see* preprocessing
 - profile preprocessing, 392
 - examples, 395
 - projection, 500
 - proportional profiles principle, 22
 - psychology, 21
 - Eve White, Eve Black, and Jane, 361, 471, 476, 479
 - happiness example, 444
 - hostile reactions to frustration, 466
 - implicit theory of personality example, 73, 199
 - multiway rating data, 35
 - perceived reality study, 36
 - showing respect example, 471
 - stimulus-response data, 345
 - unimodality constraints, 72
 - psychometrics, 12
 - PTAK, 487–488
- ## Q
- quadrilinear, 16, 229–230
 - quasi three-mode PCA, 26, 37, 100
- ## R
- R, xvii, 106, 487–488
 - rank, 49
 - core array, 178
 - core slice, 273
 - covariance matrix, 299
 - degrees of freedom, 178
 - matrix, 436
 - MCD estimator, 298
 - minimum-product rule, 179
 - multiway arrays, 21, 178–179, 240, 244
 - Parafac model, 255
 - triple cosine matrix, 90
 - rank-one array, 55–56, 59, 69, 141, 245
 - rank-one matrices, *see* rank
 - Rao distance
 - see* residuals
 - RC-association modeling, 42
 - RC(M)-association modeling, 42
 - receptor modeling, 96, 480
 - reference mode, 186, 274
 - biplot, 167
 - examples, 167
 - joint biplot, 273, 340, 443
 - examples, 340, 356, 369, 389, 423, 454
 - nested-mode biplot, 276, 399, 441–442
 - nested-mode per-component plot, 399
 - relative fit, 227, 342
 - relative residual
 - definition, 199

- relative residual (*cont'd*)
 - plot size versus levels, 201
 - size and overall variability, 199, 203
 - Tucker2 versus Tucker3 models, 200–201
- relative residual sum of squares, 227, 289
- repeated measures analysis, *see also* multivariate longitudinal analysis
- repeated measures analysis, 26, 31, 37, 46
 - between-subjects design, 375
 - doubly multivariate, 375
 - individual differences, 376
 - multivariate, 25, 312, 403
 - multivariate analysis of variance, 375
- repeated measures data
 - versus cross-sectional data, 31
- replicated PCA, 56
- resampling, *see* bootstrap
- residuals
 - distribution standardized residuals
 - examples, 342–343
 - gamma probability plot, 286
 - examples, 285–286
 - goals for examination, 283
 - inspection distributions, 288
 - least squares fitting, 285
 - multway, 286
 - principal component analysis, 284
 - probability plot, 286, 288–289
 - Rao distance, 284–288
 - residual plot, 289, 295
 - examples, 293–294, 342–343
 - structured approach, 287
 - structured sample, 286
 - decision scheme, 287–288
 - partitioning of variability, 287
 - supernormality, 286, 293
 - examples, 293–294
 - unmodeled systematic trends, 284
 - unstructured sample, 286, 292
 - decision scheme, 287–289
- response style, 123, 134, 318–320, 333, 340, 347
 - individual differences, 348
 - scale usage, 347
 - social desirability, 347
 - three-way rating scale data, 346
- restrictions on time components, 39
- robustness
 - accommodation, 284, 299
 - bad leverage fibers, 308
 - basics, 295
 - creation of outliers, 301–302
 - crisscross regression, 300
 - decontamination, 297
 - diagnostic distance plot, 305
 - examples, 303, 305
 - examples, 301
 - full-rank covariance matrix, 298
 - gamma Q-Q plot
 - examples, 305–306
 - goals, 284
 - good leverage points, 295
 - identification, 284
 - minimum covariance determinant estimator, 298–299
 - examples, 301–303
 - missing data, 301, 307
 - multimode covariance matrix, 298–299, 301
 - multway analysis, 297
 - orthogonal outliers, 295
 - outliers
 - examples, 296
 - outlying fibers, 295, 297
 - examples, 296
 - outlying individual data points, 295, 298, 300
 - examples, 296
 - outlying slices, 296–297
 - examples, 296
 - PCA per slice, 297
 - penalty function, 300
 - penalty functions, 297
 - per-slice analysis, 301
 - principal component analysis, 295, 297
 - projection pursuit
 - component analysis, 297, 299
 - multway analysis, 300
 - quick-and-dirty, 300
 - RaPCA, 299, 305
 - examples, 303–304
 - regular points, 295
 - RobPCA, 299, 305–306
 - examples, 303–305
 - robust covariance matrices, 297
 - robust PCA
 - examples, 304, 307
 - robust preprocessing, 297–298
 - robust regression algorithms, 297, 300
 - role of different modes, 296
 - three-mode common factor analysis, 299
 - two-mode procedures, 301
- rotation, 249
 - components and core array, 237, 249, 256, 339
 - examples, 250–254, 339
 - components
 - simplification, 238
 - content-based arguments, 239, 241, 249
 - core array, 244,
 - maximal slice diagonality, 252–253
 - maximal superdiagonality, 252–253

- maximum variance, 248, 253
 - oblique, 246, 248
 - orthogonal, 248
 - reducing model complexity, 244
 - simplicity, 244, 248,
 - simplification, 238
 - simplimax, 248
 - slice diagonality, 248
 - superdiagonality, 247–248
 - target rotation, 247
 - counterrotation, 238, 242
 - effect on model fit, 240
 - enabling substantive development, 240
 - equalizing variances, 239
 - finding useful directions for interpretation, 239
 - Harris–Kaiser independent cluster rotation, 240, 243–244
 - examples, 367, 484
 - joint biplot
 - examples, 423
 - model-based arguments, 239–240
 - multidimensional point of view, 243
 - nonsingular transformation, 238
 - normalized versus principal coordinates, 242, 249–250
 - oblimin, 241
 - oblique, 238, 243
 - oblique via orthogonal rotation, 243
 - orthogonal and nonorthogonal axis, 238
 - orthogonal polynomials, 241
 - examples, 267, 387
 - orthomax family, 243, 249–250
 - orthomax simplicity function, 249
 - examples, 250–252
 - natural weights, 250
 - weights, 249
 - orthonormal, 248
 - Parafac model, 237
 - promax, 241, 243
 - quartimax, 249
 - references, 240
 - simplicity components and core array, 237
 - simplicity components versus complexity core array, 239, 254, 338
 - examples, 368
 - simplicity core array versus complexity components, 239
 - simplicity versus complexity, 238
 - time-mode components, 241
 - examples, 267, 387
 - transformations, 238
 - Tucker models, 237, 243
 - Tucker toward Parafac, 247
 - Tucker2 model, 242
 - Tucker3 model, 242
 - two-mode core array, complexity, 245
 - two-way not multiway procedure, 239
 - varimax, 240–241, 243–244, 249–250
 - versus zeroing parameters, 237
 - rotational indeterminacy, PCA, 22, 88
 - rows
 - definition, 29
- ## S
- sample size
 - considerations, 176
 - outliers, 176
 - stability estimates, 176
 - stochastics, 176
 - SAS, 30
 - wide data arrangement, 46
 - scalar product, 494–500
 - scaling arrangement, *see* multiway rating scale data
 - scores, *see* principal component analysis scores
 - usage term, 48
 - second-order component models, *see* higher-order component models
 - semantic differential scales, 8–9, 72, 231, 346–347, 360–361
 - bipolar, 362
 - centering, 362
 - examples, 191
 - individual differences, 352
 - music appreciation, 191
 - normalization, 362
 - three-mode component analysis, 350
 - sensory perception, 39–40
 - example, 106
 - serial dependence, 384
 - shifted multiplicative model, 122–123
 - advantages, 122
 - disadvantages, 122
 - signal processing, 13, 22, 96
 - cell phone example, 5
 - component uniqueness, 61
 - explicit Parafac models, 61
 - theoretical multiway models, 217
 - significance tests, *see* hypothesis testing
 - similarity
 - judgement of, 40
 - similarity data
 - analysis methods, 26
 - similarity matrices, 23, 40
 - double centering, 134
 - frontal-slice normalization, 134
 - preprocessing, 134

- similarity matrices (*cont'd*)
 sets of, 33
- simplex
 correlation matrix, 379–380
 level and gain components, 380
 Markov structure, 379
 origin of, 379
 quasi, 379
 standard PCA, 379
 Wiener, 380
- simultaneous clustering and data reduction, 25
- simultaneous component analysis, 26, 35, 37, 39, 400
 Parafac2 model, 62
- simultaneous factor analysis, 26, 39
- singular value decomposition, 48, 136, 148–149, 186, 196, 227, 245, 271, 273, 436, 492, 494, 499
 basis for multiway generalizations, 50
 generalized
 two-way correspondence analysis, 436
 in triadditive modeling, 139
 modeling interaction, 135
 multiway analogue, 227
 PCA, 49
 per frontal slice, 51
 per slice of three-way array, 51
 separate and generalized Procrustes analysis, 51
 subspace uniqueness, 50
- six-way data, 470, 472
- slabs, *see* slices
- slice normalization, *see* normalization
- slices
 notation, 29
 types of, 29
- social and behavioral sciences, xviii, 13, 18–19, 238–239, 242, 311, 331, 344
 centering, 120
 cohort studies
 missing data, 145
 interpretation components, 219
 interval data, 128
 missing values, 161
 Parafac2 applications, 63
- sociology
 organizational example, 385
 Sampson monastery example, 471
- spatial evolution analysis, 25
- spectroscopy
 chromating nonlinear Rayleigh scatter, 155
 fluorescence excitation–emission data, 144, 155
 multiway, 370
 nonnegative components, 70
 parameter estimation, 281
 protein example, 472
 undesired signals as missing data, 144
- split-half procedures, *see also* dimensionality selection
- split-half procedures
 examples, 325
 orthogonal splits, 185, 325
 Parafac model, 184
 examples, 326, 328
 sample size, 184
 stochastic, 177, 184, 325
- SPSS, 30
 Categories, 42
 wide data arrangement, 46
- st-criterion, *see* deviance plot
- stages in examples
 Stage 1: Objectives, xix
 Stage 2: Data description and design, xix
 Stage 3: Model and dimensionality selection, xix
 Stage 4: Results and their interpretation, xix
 Stage 5: Validation, xix
- standard reduction equation, 138
- standardization, *see* centering
- standardization, *see* normalization
- standardization
 definition, 113
 standardized component weights, 204, 387
- STATIS, 20, 25, 37, 39, 78, 195–196, 375
 ade4 package, 106
 compromise, 106
 interstructure, 195, 399
 intrastructure, 106, 187
 R, 106
 sensory perception example, 106
 taxonomy of roses, 106
- statistical multiway models, 20, 22, 25
- statistical stability, 233, 322, 478
 fit measures, 187
 nonparametric estimation, 187
 parameter estimates, 187–188
 profile data, 312
- stimulus–response data, 470
- stimulus–response three-way data, 35
- stochastic longitudinal three-mode modeling, 36
- Strange Situation
 attachment research, 425
 clinical classification, 425, 430
 data set, 425
 procedure, 425
- striking out three-way arrays, 7
- structural base, *see* structural image
- structural equation modeling, 25, 37, 39, 47, 100, 124, 224, 376, 382, 385, 404

structural image, 19, 48, 56, 79, 213, 219, 226,
229–230, 238, 315, 317, 368, 460, 462,
483, 493–494

structure matrix, 130, 315, 321

structured component scores, 221–222, 399

structured components, 382–383, 393, 401

subspace interpretation, 212, 321

 examples, 369

subspace uniqueness, 50, 243

SumPCA model, 53, 186

 examples, 207

 model hierarchy, 68

sums-of-squares plot, 287–290

 examples, 202–203, 290–293, 335, 337

 fit/residual ratio, 202

 influential points, 291

 normalization, 292

superdiagonality, *see* core array

superidentity array, 185

supermatrix, *see* tall combination-mode matrix

system variation, 60, 332

T

tall combination-mode matrix, 7, 15–16, 30, 68,
98, 277, 297, 301, 304, 378

tensor, 16, 20, 56

Tensor Toolbox, 73, 488

terminology, basic reference, 29

test-retest reliability, 362

test equating

 missing data, 173

The Three-Mode Company, xxi, 258, 394, 431

third-order component models, *see* higher-order
 component models

three-dimensional block, *see* three-way array

three-dimensional graphs, *see* plotting

three-mode, *see also* three-way

three-mode

 preferred over three-way terminology, xviii

three-mode analysis

 birth of, 7

 dawn of, 21

 exploratory, 7

 history, 12

 leading case for multiway analysis, 44

 multiway analysis, 16

 not necessary, 7

 when necessary, 7

three-mode books, 75

three-mode common factor analysis, 21–22, 26,
37, 223, 377, 404

 interpretation, 224

 longitudinal models, 377

 missing data, 155

 multivariate longitudinal data, 385

 multiway profile data, 312

three-mode component analysis, *see also* Tucker2
 model

three-mode component analysis, *see also* Tucker3
 model

three-mode component analysis

 categorical data, 433

three-mode component models

 generalization of two-mode PCA, 44

 generalizations of SVD, 45

 IBCLAS variants, 460

 model hierarchy, 67–68

three-mode components analysis, *see* three-mode
 component analysis

three-mode core array

 notation, 226

three-mode distances, 33

three-mode mixture method of clustering, *see*
 mixture method of clustering

three-mode model tree, 66–68

three-mode models

 multiway models, 16

three-mode partitioning, 23, 25, 35

three-mode path models, 379

three-mode principal component analysis, *see*
 three-mode component analysis

three-mode rank-order data, 41

three-mode scaling, 134

three-mode unfolding, 41

three-way analysis of variance, *see* univariate
 three-way analysis of variance

three-way binary data, *see* hierarchical classes
 models

three-way conjoint analysis, 70

three-way contingency tables, *see also* three-way
 correspondence analysis

three-way contingency tables, 220, 433–434, 437,
 458

 examples, 444

 independence model, 434

 inertia, 437

 interactions, 437

 marginal dependence, 437–438

 marginal totals, 438

 measuring dependence, 434, 437

 one-way marginal totals, 438

 partial dependence, 437

 partitioning dependence, 438–439

 degrees of freedom, 450

 examples, 446, 449–450

 partitioning inertia, 458

 examples, 446, 455–456

- three-way contingency tables (*cont'd*)
 - three-way dependence, 437
 - three-way independence model, 437
 - three-way interaction, 444
 - structure plus error, 450
 - total dependence, 434, 437
 - measuring, 437
 - two-way design interaction, 444
 - examples, 446
- three-way correspondence analysis, *see also* three-way contingency tables
- three way correspondence analysis, 25, 42, 433–435, 437, 458
 - calibration axes, 452
 - design variables, 449
 - distributional equivalence, 449
 - eliminating design margins, 440
 - examples, 455
 - examples, 444
 - goals, 448
 - inertia, 439
 - interactive coding, 449
 - interpretation as two-mode CA, 442
 - interpretation
 - examples, 447, 456–457
 - joint biplot, 440, 443, 452
 - examples, 454
 - marginal dependence, 443
 - model selection, 451
 - examples, 451
 - modeling dependence, 435, 439, 458
 - examples, 446, 455
 - modeling marginal dependence, 439–440
 - modeling partial dependence, 440
 - modeling total dependence, 439
 - nested-mode biplot, 440–442, 452
 - examples, 446, 452, 456
 - marginal dependence, 442, 452
 - nonfrequency data, 449
 - partitioning dependence
 - examples, 451
 - plotting all dependences, 442, 452
 - plotting dependence, 435, 440, 458
 - plotting marginal dependence, 441
 - examples, 452–454
 - reference mode, 453
 - plotting partial dependence, 442
 - plotting total dependence, 441
 - sample size, 449
 - Tucker3 model, 439
 - examples, 446, 450
 - types of data, 448
 - weighted least squares, 439
- three way, *see also* multiway
 - three-way, *see also* three-mode
 - three-way data, xvii, 3–4, 15, 16
 - three way data array, 16, 28, 74
 - definition, 28
 - fibers, 29
 - notation, 29
 - slices, 29
 - two-mode, 16
 - type of modes, 16
 - three-way data
 - analysis of interdependence, 32
 - analyzed with two-mode methods, 18
 - bootstrap, 32
 - classification, 31
 - converting to two-way, 30
 - designs, 43
 - distributional knowledge, 7
 - fully crossed, xvii, 62, 470
 - into two-way data, 6
 - nature of, 4
 - nested, 38
 - preprocessing, 499
 - reduced to averages, 46
 - two-mode modeling, 45
 - three-way interactions, 220
 - due to outliers, 123
 - lack of stochastics, 146
 - outliers, 144
 - three-way multidimensional scaling, 26, 40
 - three-way orthogonality, *see* core array, three-way orthogonality
 - three-way profile data, 33, 34
 - parental behavior data, 314
 - preprocessing, 130
 - profile arrangement, 34, 35
 - three way rating scale data, *see also* multiway rating scale data
 - three-way rating scale data, 34, 36, 346, 348, 360, 370
 - centering, 131
 - examples, 354, 361
 - external validation, 360–361
 - factorial arrangement, 35, 131, 133
 - model comparison, 346, 353, 361
 - no individual differences, 360
 - normalization, 348
 - objectives analysis, 360
 - Parafac model, 353
 - preprocessing, 346, 348, 353, 357
 - preprocessing recommendations, 142
 - profile arrangement, 35, 131, 133, 314
 - scaling arrangement, 35, 346, 353
 - Tucker2 model, 351
 - Tucker3 model, 352

- examples, 354
- three-way rating scales
 - model comparison, 369
 - no individual differences, 347
- three-way similarity data, 41
- three-way ultrametric tree models, 23, 26
- three-way unfolding, 26
- time as an interpretational device, 36
- time series data, 17
- topographic component model, *see* Parafac model
- topographic component model
 - explicit Parafac models, 61
- trajectories, 46, 402
 - nested-mode biplot
 - examples, 441, 457
 - nested-mode biplots, 277, 442, 447
- transformation, *see* rotation
- tradditive model, 112, 137–138
 - ANOVA decompositions, 138
 - common decomposition interactions, 140
 - complete Parafac model, 141
 - complete Tucker model, 140
 - constant components, 140
 - Cowier proposals, 138, 140
 - individual differences, 140
 - missing data, 157
 - offsets, 118
 - separate decompositions interactions, 140
 - types of, 138
- trilinear, 16, 58, 155, 221
- tubes
 - definition, 29
- TUCKALS algorithms, 97–98
 - accuracy, 333
 - alternating least squares, 84
 - best approximate decomposition, 99
 - compression with B-splines, 103
 - convergence properties, 99, 103
 - description, 100
 - FM algorithm, 99
 - global optimum, 85
 - guaranteed solution, 85
 - missing data, 99
 - missing data example, 103–104
 - multimode covariance matrix, 99
 - missing data, 100
 - quasi three-mode PCA, 100
 - structural equation models, 100
 - quality solutions, 164
 - Tucker Method 1 as starting solution, 86, 304
 - Tucker2 model, 99
- Tucker
 - citations, 21
 - interview, 12
- Tucker Method 1 algorithm, 97
 - dimensionality selection, 98
 - starting solution TUCKALS, 98–99
- Tucker model
 - multiway analogue singular value decomposition, 227
 - basic form, 215
 - cross-linking of components, 90
 - decomposition model, 84
 - dimensionality selection, 179
 - estimation, 101
 - evaluating fit, 233, 334–336
 - exploratory, 85, 333
 - fitting systematic variability, 184
 - flexibility, 215
 - four-way, 474
 - functional constraints, 72
 - higher-order component models, 223
 - maximum-product rule, 98, 103
 - multiway, 66
 - multiway profile data, 313
 - orthogonality, 69
 - overidentification, 99
 - partitioning of variability, 215
 - practical aspects, 331
 - projection into lower-dimensional space, 69, 211
 - rotational freedom, 213, 331
 - separate interpretation modes, 213
 - smoothness constraints, 71
 - subspace interpretation, 322
 - subspace uniqueness, 70, 211
 - transformational freedom, 99
- Tucker structure, *see* Parafac model
- Tucker0 model, *see* SumFCA model
- Tucker1 model, 52, 186, 350
 - longitudinal data, 52
 - model hierarchy, 68
 - solved by two-mode PCA, 52
- Tucker2 model, *see also* extended core array
- Tucker2 model, 53–54, 97, 99–102
 - asymmetric, 99
 - average correlation matrix, 223
 - basic form, 217
 - changes in scores and loadings, 223
 - decomposition model, 54
 - definition, 53
 - degrees of freedom, 177
 - extended core array, 231, 247
 - HICLAS variant, 460
 - higher-order scores and loadings, 223
 - higher-order structure matrix, 224
 - INDSCAL relative, 23
 - joint biplot, 273–274, 276
 - lack of component uniqueness, 54

- Tucker2 model (*cont'd*)
 - lack of nesting, 54
 - linear constraints, 95
 - longitudinal example, 401
 - model hierarchy, 68, 186
 - Murakami form, 222–223
 - preprocessing, 224
 - orthogonality core array, 66
 - Parafac variant, 231
 - partitioning of variability, 54
 - stochastic formulation, 382
 - subspace uniqueness, 54
 - tall combination-mode matrix, 278
 - Tucker3 algorithms
 - comparison, 161
 - GEPCAM and TUCKALS, 161
 - starting values, 161, 164
 - missing data estimates, 164
 - missing data
 - convergence, 161, 164, 168
 - Tucker3 model, *see also* Core array
 - Tucker3 model, 55, 97
 - Appello's reinvention, 21
 - basic form, 218
 - complexity, 56
 - component uniqueness, 56
 - covariance form, 224
 - decomposition model, 56
 - definition, 54
 - degrees of freedom, 177
 - DiffFit procedure, 180
 - distance form, 224, 228
 - equivalence Parafac model, 158, 256
 - examples, 8, 258, 409, 427
 - external information, 73, 234
 - generalization of SVD, 56
 - HICLAS variant, 460
 - higher-order components, 223
 - higher-order SVD, 56
 - joint biplot, 273–274
 - lack of nesting, 56
 - linear constraints, 95
 - loadings, principal coordinates, 55
 - minimum-product rule, 56
 - missing data estimation, 159
 - mixed multiplicative model, 140
 - model hierarchy, 68, 186
 - more terms than Parafac model, 59
 - Murakami form, 224–225, 228
 - nested linear combinations, 223, 228
 - partitioning of variability, 56
 - per-component plot, 271
 - precursor, 21
 - rank-one array, 55
 - restrictions on number of components, 56
 - scores and loadings, 224
 - scores, standard coordinates, 55
 - single component in a mode, 56, 170, 333
 - split-half procedures, 185
 - structural image, 230
 - symmetric, 99
 - tensors, 56
 - three-mode singular value decomposition, 439
 - three-way orthogonal core array, 65
 - two-component equivalence Parafac model, 197, 334
 - used for compression, 103
 - versus Parafac model, 213
 - Tucker4 model, 475, 487
 - compared to Tucker3 model, 480, 487
 - examples, 481
 - minimum-product rule, 66, 179
 - model symmetry, 475
 - multiple personality data, 471
 - vec notation, 475
 - Tucker6 model
 - fMRI example, 472
 - Tuckers model, 475, 487
 - algorithm, 24
 - two-way and three-way data, 4, 30
 - two-way contingency tables, *see also* two-way
 - correspondence analysis
 - two-way contingency tables
 - chi-term, 435
 - examples, 452–453
 - dependence, 435
 - independence model, 435
 - inertia, 435
 - measuring dependence, 435
 - standardized residual, 435
 - two-way correspondence analysis, *see also*
 - two-way contingency tables
 - two-way correspondence analysis, 435
 - attraction between categories, 436
 - biplot, 437
 - generalized singular value decomposition, 436, 439
 - modeling dependence, 436
 - partitioning inertia, 436
 - plotting dependence, 437, 440
 - principal coordinates, 220
 - repulsion between categories, 436
- U**
- unfolding
 - used for matricization in chemistry, 46
 - uniqueness, *see* Parafac model

univariate multiway analysis of variance
 profile data, 133
 univariate three-way analysis of variance, 36, 220
 examples, 133
 univariate three-way factorial data, 138
 decomposition, 138
 univariate two-way analysis of variance, 111
 univariate two-way factorial data, 135
 Mandel contribution, 136
 Tucker-Gillott discussion, 136–137

V

validating multiway analyses, 233, 486
 validation, *see also* external validation
 validation, *see also* internal validation
 validation, 207, 322, 342, 422, 427, 486
 variability, variance, sum of squares, 226
 variable-component correlations, 48, 55, 130, 210,
 216, 221–222, 225, 241–242, 244, 315,

320–321, 474
 variable dependence, 383
 varimax rotation, 240, 241, 243–244, 249–250
 examples, 196, 249, 254, 274, 338–339,
 364–367, 396–397

vector

angle, 500
 collinear, 500
 cosines and correlations, 500
 length, 500
 orthogonal, 500
 projection, 500

vectors, 499

W

way, xvii, 16
 definition, 28
 weighted PCA, 57
 wide combination-mode matrix, 7, 15–16, 30, 55,
 98, 297–298, 301, 303, 378