

SUBJECT INDEX

- Absorption bands, mid-infrared pulse-shaping technology, ground-state vibrational dynamics, coherent control, 11–16
- Absorption coefficient:
 - flat well, special configuration and permittivity, libration spectral functions, 487–489
 - ice spectrum:
 - complex permittivity, 424
 - high-temperature resonance calculations, 396–402
 - research background, 395–396
 - rotary elasticity, dimensionless absorption, 453–456
 - T- and V-band calculations, 404–407
 - temperature influence, water spectrum, 372–373
- Acoustic Casimir effect, ultrathin solid-fluid-solid films, 222–223
- “Action spectrum,” dissociative double photoionization, 141–143
- Adiabatic ground state transfer, local control theory, 89–90
- Adiabatic rapid passage, mid-infrared pulse shaping, 2–3
- Adsorption rate constant, interface between two immiscible electrolyte solutions, ion transfer kinetics, 187–195
- Analytical theory, spectral calculations, 328–330
- Angel’s strong-fragile classification, thermotropic liquid crystals, shear viscosity temperature dependence, 261–262
- Angular correlations, double photoionization, complex atoms, 109
- Angular integration, double photoionization, 110–111
- Angular vibrations, ice spectrum, rotary force constant, 457
- Anisotropic corrugations, ultrathin solid-fluid-solid films, 237–241
- Applicability criterion, harmonic longitudinal vibrations, 370–371
- Aqueous spectra. *See* Ice spectra; Water spectra
- Argon, double photoionization, valence DPI, deep inner-shell thresholds, 122–124
- Arrhenius behavior:
 - calamitic liquid crystal fragility, 296–297
 - thermotropic liquid crystals:
 - energy landscapes, 301–303
 - shear viscosity temperature dependence, 261–262
- Asymmetric interaction parameters, interface between two immiscible electrolyte solutions, 177–180
- Atomic structures. *See also* Complex atoms
 - thermotropic liquid crystals, energy landscapes, 301–303
- Atomistic mesogenic models, thermotropic liquid crystals, 253–255
- supercooled liquids, heterogeneous dynamics, 294
- Auger effect, double photoionization:
 - inner-shell DPI, 126–128, 146
 - molecules, 134–140
 - resonant Auger phenomena, 143
- Autocorrelation functions (ACFs):
 - dielectric relaxation, spectral functions, 496–499
 - extended simple point charge (SPC/E) model, 506–509
 - ice spectrum, nonrigid dipole elastic longitudinal vibration and rigid dipole elastic reorientation, 432–436
 - ice/water spectral calculations, 329–330
 - nonrigid dipoles, transverse susceptibility, 460–463
- Autoionization, dissociative double photoionization, 140–143, 147

- Automated two-dimensional infrared spectroscopy, mid-infrared pulse-shaping technology, 16–25
- Average perturbation theorem, dielectric relaxation, complex susceptibility, Gross collision model, 500–502
- Azimuthal velocity, hat well, special configuration and permittivity, precession approximation, planar libration, 478
- Back-to-back double layer ion distribution, interface between two immiscible electrolyte solutions, 165–180
- Bandwidth estimation, ice and water spectra, 424–426
- Bending elastic forces:
 - temperature influence, water spectra, 388–389
 - ultrathin solid-fluid-solid films, 232–237
- Biaxiality parameter, thermotropic liquid crystals:
 - Landau-de Gennes (LdG) theory, 269–270
 - mesophase models, 254–255
- Boltzmann energy distribution:
 - dielectric relaxation:
 - Gross collision model, 502–505
 - qualitative analysis, 493–495
 - hat well, special configuration and permittivity, planar spectral functions, 483–485
 - interface between two immiscible electrolyte solutions, 179–180
 - nonrigid dipole elastic longitudinal vibration and rigid dipole elastic reorientation, 426–432
- Born-Oppenheimer approximation, local control theory, 32
 - coupled electronic/nuclear motion, 86–89
 - electronic excitation, 60–65
 - optical paralysis, 66–69
- Bose-Einstein (BE) equation, ice/water spectral analysis, overall permittivity, Raman spectrum, 350–351
- Boundary-layer equations, ultrathin solid-fluid-solid films, 225–228
- Bragg deflection angle, germanium acoustooptic modulator, 7
- Butler-Volmer formalism, interface between two immiscible electrolyte solutions, electron transfer kinetics, 196–205
- Butler-Volmer potential dependence, interface between two immiscible electrolyte solutions, ion transfer kinetics, 182–195
- Cadmium, double photoionization, valence DPI, deep inner-shell thresholds, 124–125
- Cage structures, thermotropic liquid crystals, supercooled liquids, heterogeneous dynamics, 292–294
- Calamitic liquid crystals:
 - energy landscapes, 298–303
 - fragility, 294–297
 - future research issues, 312–313
 - inherent structure orientation, 303–306
 - liquid crystal-supercooled liquid orientational dynamics, 264–266
 - thermotropic mesophases and order parameters, 252–255
- Capacitance, interface between two immiscible electrolyte solutions, enhancement of, 167–180
- Capillary wave persistence, interface between two immiscible electrolyte solutions, 158–180
 - ion transfer kinetics, 189–195
- Carboxy hemoglobin, local control theory, selective mode excitation, 45–47
- Center-of-mass (COM) motion, extended simple point charge (SPC/E) model, 506–509
- Chemical reactions, mid-infrared pulse-shaping technology:
 - ground-state vibrational dynamics, coherent control, 10–16
 - two-dimensional infrared spectroscopy, 17–25
- Cholesteric mesophase, thermotropic liquid crystals, 253–255
- Chu-Moroi (CM) model, thermotropic liquid crystals, orientational and translational order, 307–311
- Classical mechanics:
 - harmonic longitudinal vibrations, 369–371
 - ice and water spectra, minimal action and phase transition estimation, 424–426
 - local control theory, 33–34
 - photoassociation and, 51

- Coherent control, mid-infrared pulse-shaping technology, ground-state vibrational dynamics, 7–16
- Coincidence rates, double photoionization, molecular quantum yields, 131–133
- Cold target recoil momentum spectroscopy (COLTRIMS), double photoionization, 111
- Cole-Cole plots:
 - ice spectra, high-temperature resonance spectra calculations, 396–402
 - wideband water spectrum, room temperature conditions, ordinary and heavy water, 365–367
- Collective orientational dynamics, thermotropic liquid crystals, isotropic-nematic phase transition, 279–287
- Columnar phases, thermotropic liquid crystals, 253–256
- Common-ion mechanism, interface between two immiscible electrolyte solutions, electron transfer kinetics, 197–205
- Compact layer structure, interface between two immiscible electrolyte solutions, 166–180
- Complete experiments, double photoionization, indirect processes, 110
- Complex atoms, double photoionization, 120–128
 - angular correlations, 109
 - deep inner-shell thresholds, valence DPI, 121–124
 - inner-shell DPI, 124–128
- Complex susceptibility, dielectric relaxation:
 - Gross collision model, 499–505
 - perturbed trajectories, 495–496
 - qualitative analysis, 491–495
 - spectral function, autocorrelation function, 496–499
- Complex topography, ultrathin solid-fluid-solid films, surface undulation control, 224
- Compressive strains, ultrathin solid-fluid-solid films, 237–241
- Computer simulations:
 - thermotropic liquid crystals, 273–297
 - calamitic liquid crystals, 303–306
 - energy landscapes, 297–311
 - isotropic-nematic transition dynamics, 275–287
 - model systems, 274–275
 - orientational and dynamical signatures, 306–311
 - supercooled liquids, 288–297
 - ultrathin solid-fluid-solid films, 229–231
- Concavity, direct double photoionization, helium, 106–109
- Continuity breaks, temperature influence, water spectra, 389–393
- Continuum theory, ultrathin solid-fluid-solid films, 225–228
- Contour plots, local control theory, coupled electronic/nuclear motion, 87–89
- Control systems:
 - coherent control, mid-infrared pulse-shaping technology, ground-state vibrational dynamics, 7–16
- local control:
 - analytical examples, 34–36
 - basic formulation, 32–33
 - coupled electronic/nuclear motion, 86–94
 - adiabatic ground state transfer, 89–90
 - combined electron-nuclear dynamics, 91–94
 - model parameters, 86–89
 - two-state model transfer, 90–91
- electronic excitation, 60–65
 - photodissociation and, 69–72
- Fermi's Golden Rule, 35–36
- free particle deceleration, 34
- future research issues, 94–95
- multiple potential wells, 51–60
 - double well potentials, 51–57
 - triple-well potentials, 57–60
- optical paralysis, 65–69
- photoassociation, 47–51
 - orientation role in, 83–86
- photodissociation, 36–44
 - direct photodissociation, 36–39
 - electronic transitions and, 69–72
 - predissociation dynamics, 40–42
 - selective mode excitation vs., 42–44
- quantum vs. classical mechanics, 33–34
- rotational motion, 72–86
 - directional motion control, 73–78
 - excited-state fragmentation, 78–83
 - photoassociation orientation, 83–86
- selective mode excitation, 44–47
 - photodissociation vs., 42–44
- theoretical background, 30–32

- Convergent close coupling (CCC), direct double photoionization, helium, 108–109
- Cooling process, local control theory, 32
- Core-hole formation:
 direct double photoionization, molecular Auger effect, 136–140
 double photoionization, inner-shell DPI, 126–128
- Core polarization potentials (CCP), local control theory, optical paralysis, 66–69
- Core-shell structures, ultrathin solid-fluid-solid films, surface undulation control, 224
- Correlation energy, double photoionization, 144–146
- Coulomb interactions, interface between two immiscible electrolyte solutions, 176–180
 ion transfer kinetics, 193–195
- Coupled electronic/nuclear motion, local control theory, 86–94
 adiabatic ground state transfer, 89–90
 combined electron-nuclear dynamics, 91–94
 model parameters, 86–89
 two-state model transfer, 90–91
- Coupled oscillator systems, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 24–25
- Coupled two-state model, local control theory, 90–91
- Coupling reactions, thermotropic liquid crystals, orientational and translational order, 306–311
- Cross-peak formation, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 18–25
- Cross-sectional diameter, thermotropic liquid crystals, computer simulations, 274–275
- Current-time response, interface between two immiscible electrolyte solutions, interfacial deposition, 205–208
- “Curtain edge” structures, dissociative double photoionization, 141–143
- Data presentation, double photoionization, 116–120
- Data validation, interface between two immiscible electrolyte solutions, 167–180
- Debye relaxation:
 ice spectrum:
 loss gap, 420–422
 nonresonance spectrum, 409–410
 interface between two immiscible electrolyte solutions, 168–180
 nonrigid dipole elastic longitudinal vibration and rigid dipole elastic reorientation, 426–432
 temperature influence, water spectrum, 373–380
 steady state and LIB subensemble, 381–384
 thermotropic liquid crystals:
 energy landscapes, 303
 future research issues, 312–313
- Debye-Stokes Einstein (DSE) relationship, thermotropic liquid crystals, dynamic decoupling phenomena, 263–264
- Deep inner-shell thresholds, valence double photoionization, 121–124
- Deep well permanent dipoles, nonharmonic libration, permanent dipoles, 337–343
- de Gennes-McMillan coupling, thermotropic liquid crystals, orientational and translational order, 306–311
- Density profile:
 double well potential, local control theory, 55–57
 interface between two immiscible electrolyte solutions, 178–180
 L-L interface, 157–158
 local control theory, excited-state fragmentation, 81–83
 temperature-dependent water spectra, 382–384
 thermotropic liquid crystals, Onsager theory, 266–267
- Depolarized light scattering (DLS), thermotropic liquid crystals, liquid crystal-supercooled liquid orientational dynamics, 265–266
- Deuterium, double photoionization, valence DPI, deep inner-shell thresholds, 122–124
- Diabatization, coupled electronic/nuclear motion, local control theory, 86–89
- Diagonal peaks, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 18–25

- Diatomic molecules, direct photodissociation, 36–39
- 1,4-Diazobicyclo[2,2,2]octane (DABCO), double photoionization, molecular Auger effect, 138–140
- Dication structures, double photoionization: Auger effect, 138–140
TOEsCO and TOF-PEPECO technique, 128–130
- Dielectric loss spectra: ice spectrum: high-temperature resonance spectra, 396–402
loss gap, 420–422
nonresonance spectrum, 407–410
research background, 395–396
transverse vibration and, 406–407
- ice/water spectral analysis: nonharmonic libration, 340–343, 344–346
overall permittivity, 350–351
- nonrigid dipoles, nonharmonic vibrations, transverse-vibration loss spectrum, 471–473
- temperature influence, water spectrum, 373–380
extended simple point charge (SPC/E) model, 507–509
- thermotropic liquid crystals, 262–264
- wideband room temperature conditions, ordinary and heavy water, 358–367
- Dielectric relaxation: Gross collision model complex susceptibility, 499–505
spectral function, 502–505
undisturbed variable transitions, 499–502
- ice/water spectra, 335–351
hydrogen bond dipolar harmonic vibration, 343–346
librational state, 335–336
low-frequency Raman spectrum, 350–351
molecular models, 334
nonharmonic libration, deep well permanent dipole, 337–343
nonrigid dipole, nonharmonic transverse vibration, 346–348
overall permittivity, 348–350
Teixeira's data analysis, 352–353
vibrational state, 336–337
perturbed trajectories, power consumption, 495–496
qualitative features, 491–495
spectral function and autocorrelation function spectrum, 496–499
temperature influence, water spectra, 373–380
- Diffusional jumps, ice/water spectral models, 352–353
- Diffusion description, thermotropic liquid crystals, isotropic-nematic phase transition, 258–260
- Dimensionless absorption: dielectric relaxation, spectral function, autocorrelation function, 497–499
hat well, special configuration and permittivity, libration spectral functions, 487–489
ice spectrum: autocorrelation functions, 435–436
integral absorption, 449
inverse temperatures, 414–417
rotary elasticity, 453–456
nonrigid dipoles, nonharmonic vibrations, autocorrelation, transverse susceptibility, 462–463
- Dimensionless capacitance measurement, interface between two immiscible electrolyte solutions, 173–180
- Dimensionless strong-collision frequency, ordinary and heavy water spectra, static permittivity, isotropic independence, 356–357
- Dimer formation: ice spectrum, kinetic energy, 438–440
ice/water spectral calculations, molecular modeling, 331–333
nonrigid dipole elastic longitudinal vibration and rigid dipole elastic reorientation, two-fraction model, 426–432
- Dipolar interactions: far-infrared ice spectra, 509–510
hat well, special configuration and permittivity, dipole confinement, hydrogen bond, 479–481
ice/water spectral calculations: hydrogen bond harmonic vibrations, 343–346
nonharmonic libration, 337–343
nonharmonic transverse vibrations, nonrigid dipoles, 346–348

- Dipolar interactions: (*Continued*)
 interface between two immiscible electrolyte solutions, 159
 dipolar width, 164–165
 temperature influence, water spectra, 383–384
- Direct double photoionization:
 helium, 105–109
 molecules, 133–134, 146
- Direct knockout mechanism, direct double photoionization, helium, 108–109
- Direct photodissociation, local control theory, 36–39
- Discotic liquid crystals:
 future research issues, 314
 orientational time correlation function, 281–287
 thermotropic mesophases and order parameters, 252–255
- Dissociative double photoionization, molecules, 140–143, 147
- Distribution potential, interface between two immiscible electrolyte solutions, 156–157
- Double-Debye formula:
 temperature influence, water spectrum, steady state and LIB subensemble, 381–384
 wideband water spectrum, room temperature conditions, 354–355
 ordinary and heavy water, 363–367
- Double layer structure, interface between two immiscible electrolyte solutions, 167–180
- Double-Lorentz representation, temperature-dependent water spectra, 381–384
- Double photoionization (DPI):
 angular integration, 110–111
 complex atoms, 120–128
 angular correlations, 109
 deep inner-shell thresholds, valence DPI, 121–124
 inner-shell DPI, 124–128
 correlation and, 144–146
 data presentation, 116–120
 direct *vs.* indirect processes, 104–105
 energetic electron pair analysis:
 magnetic bottle time-of-flight methods, 113–116
 molecules, 128–130
 helium, 105–109
 indirect atomic processes, 109–110
 molecules:
 Auger effects, 134–140
 direct DPI, 133–134
 dissociative DPI, 140–141
 mapping and electron distributions, 133
 quantum yields, 131–133
 resonant Auger phenomena, 144
 TPEsCO and TOF-PEPECO spectra, 128–130
 research background, 104–105
 summary and future research, 146–148
 threshold photoelectrons coincidence, 111–113, 116
 molecules, 128–130
- Double well potentials, local control theory, 51–57
- Driving instability mechanisms, ultrathin solid-fluid-solid films, 222–223
- Droplet size dependence, interface between two immiscible electrolyte solutions, ion transfer kinetics, 187–195
- “Dutcher films,” research background, 217–220
- Dynamic correlation, double photoionization, 144–146
- Dynamic decoupling phenomena, thermotropic liquid crystals, 262–264
- Effective core potentials (ECPs):
 hat well, special configuration and permittivity, precession approximation, planar libration, 477–479
 local control theory, optical paralysis, 66–69
- Ehrenfest theorems:
 local control theory, quantum *vs.* classical mechanics, 33–34
 thermotropic liquid crystals, Onsager theory, 266–267
- Eigenstates:
 local control theory, photoassociation, 48–51
 mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 18–25
- Elasticity theory, ultrathin solid-fluid-solid films, 225–228
- Elastic longitudinal vibration:
 nonrigid dipoles, ice spectra, 426–458
 autocorrelator transformation, 432–436
 dimer kinetic energy, 438–440

- equations of motion and spectral function, 443–448
- frozen translation librations, 451–452
- frozen translation rotations, 450–451
- heavy water, 452–453
- hydrogen bond strained-state energy, 436–437
- hydrogen bond translation/reorientation independency, 449–450
- integral absorption, 448–449
- rotary elasticity, 453–458
- translational vibration dielectric response, 440–443
- two-fraction dimer model, 426–432
- temperature influence, water spectra, 376–380
- Elastic reorientation, rigid dipoles, ice spectra, 426–458
 - autocorrelator transformation, 432–436
 - dimer kinetic energy, 438–440
 - equations of motion and spectral function, 443–448
 - frozen translation:
 - librations, 451–452
 - rotations, 450–451
 - heavy water, 452–453
 - hydrogen bond:
 - strained-state energy, 436–437
 - translation/reorientation independency, 449–450
 - integral absorption, 448–449
 - rotary elasticity, 453–458
 - translational vibration dielectric response, 440–443
 - two-fraction dimer model, 426–432
- Electrified liquid-liquid interface:
 - electron transfer kinetics, 195–205
 - future research issues, 208–209
 - interfacial deposition, 205–208
 - ion transfer kinetics, 180–195
 - research background, 153–157
 - structure, 157–180
- Electrocapillary curves, interface between two immiscible electrolyte solutions, interfacial deposition, 207–208
- Electrochemical techniques, interface
 - between two immiscible electrolyte solutions, electron transfer kinetics, 199–205
 - Electrolyte-free phases, interface between two immiscible electrolyte solutions, 179–180
- Electron distribution, double photoionization, 133
- Electron dynamics, local control theory, 91–94
- Electronic excitation, local control theory, 60–65
 - photodissociation and, 69–72
- Electronic transitions, local control theory, photodissociation and, 69–72
- Electron localization function (ELF), local control theory, coupled electronic/nuclear motion, 86–89
- Electron transfer (ET), interface between two immiscible electrolyte solutions, 195–205
- Energetic electron pair analysis, double photoionization, magnetic bottle time-of-flight methods, 113–116
- Energy landscapes, thermotropic liquid crystals, 297–311
- Energy loss function, ice spectrum, 407
 - research background, 395–396
- Enthalpy, thermotropic liquid crystals, supercooled regimes and glass transition, 256–258
- Entropy, thermotropic liquid crystals, supercooled regimes and glass transition, 256–258
- Equations of motion:
 - extended simple point charge (SPC/E) model, 506–509
 - ice spectra:
 - Hamilton equation transformation, 443–444
 - translational vibration, 441–443
 - ice/water nonharmonic libration, 342–343
 - ice/water spectral models, 334
 - nonrigid dipoles, nonharmonic vibrations:
 - autocorrelation, transverse susceptibility, 461–463
 - nonlinear equations, 463–464
- Equilibrium conditions:
 - dielectric relaxation, complex susceptibility, Gross collision model, 499–502
 - ice spectrum, nonrigid dipole elastic longitudinal vibration and rigid dipole elastic reorientation, 431–432

- Etalon generation, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 21–25
- Euler angles, thermotropic liquid crystals, Mode coupling theory (MCT), 270–273
- Evolutionary models, ultrathin solid-fluid-solid films, 231–237, 245–246
- Excess protons, ice/water spectral calculations, vibrational state, 336–337
- Excited-state populations:
 double photoionization, complex atoms, 120–124
 local control theory:
 electronic excitation, 61–65
 fragmentation, 78–83
- Expansion coefficients, thermotropic liquid crystals, isotropic-nematic transition, 278–287
- Expectation values, local control theory, combined electron-nuclear dynamics, 93–94
- Extended simple point charge (SPC/E) model, molecular dynamics simulations, 505–509
- Extra-thermodynamic techniques, interface between two immiscible electrolyte solutions, 155–157
- Fano profiles, double photoionization, indirect processes, 110
- Faraday's law, interface between two immiscible electrolyte solutions, ion transfer kinetics, 183–195
- Far infrared (FIR) spectroscopy:
 aqueous spectral calculations, 328–330
 nonharmonic libration, permanent dipoles, 337–343
 dielectric relaxation, spectral function, autocorrelation function, 498–499
 far-infrared ice spectra, 509–510
 ice spectrum, loss gap, 420–422
 nonresonance ice spectra, 510
- Fast Fourier transform, ultrathin solid-fluid-solid films, evolutionary models, 234–237
- Fast-pulse discharge lamp design, double photoionization, time-of-flight photoelectron-photoelectron coincidence (TOF-PEPECO) technique, 114–116
- Femtosecond pulse shaping:
 double photoionization, complex atoms, 121–124
 research background, 1–4
- Fermi correlation, double photoionization, 145–146
- Fermi's Golden Rule, local control theory, 35–36
- Field-dressed potentials, local control theory, predissociation dynamics, 41–42
- Field-induced photodissociation, local control theory, 38–39
 excited-state fragmentation, 80–83
- Final-state correlation, direct double photoionization, 107–109
- Finite difference approximation, ultrathin solid-fluid-solid films, 229–231
- First-order perturbation, local control theory, 35–36
- Fitted model parameters, temperature influence, water spectra, 376–380
 harmonically vibrating hydrogen bond molecules, 384–385
 steady state and LIB subensemble, 380–384
- Fitting coefficients, ice spectrum, nonresonance spectrum, 408–410
- FORCE theorem, dielectric relaxation, complex susceptibility, Gross collision model, 500–502
- Formaldehyde, double photoionization, molecular Auger effect, 137–139
- Form factor, ice/water nonharmonic libration, 341–343
- Fourier transform:
 hat well, special configuration and permittivity, planar spectral functions, 483–485
 interface between two immiscible electrolyte solutions, 168–180
 local control theory, electronic excitation, 64–65
 mid-infrared pulse-shaping technology:
 ground-state vibrational dynamics, coherent control, 13–16
 two-dimensional infrared spectroscopy, 19–25
- Four-wave mixing geometry, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 25

- Fragility index:
 calamitic liquid crystals, 312–313
 future research issues, 314
- Fragmentation mechanisms, local control theory:
 direct photodissociation, 36–39
 excited-state fragmentation, 78–83
- Franck-Condon region:
 double photoionization:
 Auger effect, 138–140
 maps and electron distributions, 133
 TOESCO and TOF-PEPECO technique, 128–130
 local control theory, electronic excitation, 62–65
- Free-energy functional method, interface between two immiscible electrolyte solutions, 170–180
- Free induction decay, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 19–25
- Free particle deceleration, local control theory, 34
- Frequency comb spectrum, germanium acoustooptic modulator, 7
- Frequency dependence:
 hydrogen bonds, longitudinal/rotational vibration coupling, 369
- ice spectra:
 high-temperature resonance calculations, 396–402
 low-temperature spectrum, 414–417
 rotary elasticity, dimensionless absorption, 454–456
- ice spectrum:
 loss gap, 420–422
 rotary elasticity, dimensionless absorption, 453–456
 T- and V-band calculations, 404–407
- ice/water nonharmonic libration, 339–343
 nonharmonic transverse vibrations, nonrigid dipoles, 347–348
 nonrigid dipoles, nonharmonic vibrations, transverse-vibration loss spectrum, 471–473
 thermotropic liquid crystals, orientational time correlation function, 285–287
- Frequency factor, nonrigid dipoles, nonharmonic vibrations, transverse-vibration loss spectrum, 471–473
- Friction coefficient, ice/water spectral calculations, 329–330
- “Frozen” rotation, ice spectra, translational vibrations, 450–451
- “Frozen” translations, ice spectrum librations, 451–452
 hydrogen bond length restrictions, 458
- Frumkin technique, interface between two immiscible electrolyte solutions, electron transfer kinetics, 198–205
- Functional derivatives, ultrathin solid-fluid-solid films, 244–245
 pressure behavior, 227–228
- Galvani potential difference, interface between two immiscible electrolyte solutions, 155–157
 electron transfer kinetics, 196–205
 ion distribution, 165–180
- Gaussian models, thermotropic liquid crystals, energy landscapes, 300–303
- Gaussian-shaped pulses:
 local control theory, adiabatic ground state transfer, 89–90
 mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 20–25
- Gay-Bern (GB) pair potential:
 orientational time correlation function, 281–287
 thermotropic liquid crystals:
 computer simulations, 274–275
 energy landscapes, 298–303
 isotropic-nematic transition, 276–287
 orientational and translational order, 306–311
 supercooled liquids:
 dynamic similarity, 288–289
 heterogeneous dynamics, 290–294
- Germanium acoustooptic modulator (AOM), mid-infrared (mid-IR) pulse-shaping technology:
 basic principles, 4–5
 performance evaluation, 5–7
- Germanium compounds, double photoionization, molecular Auger effect, 136–140

- Gibbs energy, interface between two immiscible electrolyte solutions, 155–157
- dipolar and Yukawa interactions, 159
- interfacial deposition, 207–208
- interfacial roughness, 171–180
- ion transfer kinetics, 193–195
- Glass-transition temperature:
- calamitic liquid crystals, 312–313
- substrate-fluid-solid films, 220–221
- thermotropic liquid crystals, supercooled liquid regimes, 255–258
- Global deformation, ultrathin solid-fluid-solid films, 237–241
- Global optimization, local control theory, 31–32
- Gouy-Chapman capacitance, interface between two immiscible electrolyte solutions, 165–180
- electron transfer kinetics, 199–205
- Grayscale imaging, double photoionization, 117–120
- Gross collision model:
- dielectric relaxation, complex susceptibility: spectral function, 502–505
- undisturbed variables, 499–502
- ice/water nonharmonic libration, 338–343
- Ground-state populations, local control theory, electronic excitation, 61–65
- Ground-state vibrational dynamics, mid-infrared pulse-shaping technology, coherent control, 7–16
- Hamaker constant, ultrathin solid-fluid-solid films:
- evolutionary models, 234–237
- pressure behavior, 226–228
- Hamiltonian equations:
- ice spectrum equations of motion, 443–444
- local control theory, 32–33
- quantum vs. classical mechanics, 33–34
- Harmonic oscillator, nonrigid dipoles, nonharmonic vibrations, transverse complex susceptibility, 468–471
- Harmonic vibrations:
- classical theory, 369–371
- hydrogen bonds, longitudinal/rotational vibration coupling, 368–369
- ice/water spectral models, 336–337
- hydrogen bond dipoles, 343–346
- temperature influence, water spectrum, 373–380
- hydrogen bonded molecules, 384–385
- wideband water spectrum, room temperature conditions, ordinary and heavy water, 360–367
- Hartree-Fock limit, double photoionization, correlation energy, 144–146
- Hat potential:
- far-infrared ice spectra, 509–510
- ice spectra:
- high-temperature resonance spectra calculations, 396–402
- hydrogen bond bending and breakage, 403–407
- ice/water nonharmonic libration, 337–343
- spatial configuration, summary permittivity: hat potential evolution, 474–479
- planar libration, 479–482
- spectral function, 482–489
- water molecules, 489–491
- temperature influence, water spectrum, 373–380
- steady state and LIB subensemble, 381–384
- Heating process, local control theory, 32
- predissociation dynamics, 40–42
- Heavy water:
- ice spectra:
- dimer kinetics, 439–440
- kinetic energies, 452–453
- ice/water nonharmonic libration, 340–343
- wideband water spectrum, room temperature conditions, 354
- low-density and high-density concepts, 354–355
- Raman spectra, 358–367
- static permittivity isotropic independence, 355–357
- Helium, direct double photoionization, 105–109
- Herringbone formation, ultrathin solid-fluid-solid films, surface undulation control, 224
- Hess-Frenkel-Allen model, thermotropic liquid crystals, orientational and translational order, 307–311
- Heterogeneous dynamics, thermotropic liquid crystals, supercooled liquids, dynamic similarity, 289–294

- High-density-liquid (HDL), wideband water spectrum, room temperature conditions, 354–355
- High-frequency approximation:
dielectric relaxation, spectral function, autocorrelation function, 498–499
- ice spectrum, nonrigid dipole elastic longitudinal vibration and rigid dipole elastic reorientation, 432–436
- nonrigid dipoles, nonharmonic vibrations:
autocorrelation, transverse susceptibility, 461–463
transverse complex susceptibility, 471
- High-temperature resonance spectra, ice spectra, 396–407
calculations, 396–402
lattice mode spectroscopy, 402–407
- Hole burning, mid-infrared pulse-shaping technology:
ground-state vibrational dynamics, coherent control, 11–16
two-dimensional infrared spectroscopy, 17–25
- Homeotropically aligned nematic liquid crystal, thermotropic liquid crystals, isotropic-nematic phase transition, 260
- Hydrides, direct double photoionization, 134–135
- Hydrodynamics, interface between two immiscible electrolyte solutions, ion transfer kinetics, 191–195
- Hydrogen bonds:
aqueous spectral calculations, 328–330
molecular modeling, 330–333
far-infrared ice spectra, 509–510
harmonic longitudinal vibrations, classical theory, 369–371
- hat well, special configuration and permittivity:
dipole precession confinement, 479–482
libration spectral functions, 485–489
planar libration approximation, 478–479
water molecules, 489–491
- ice spectrum:
bending and breakage, 403–407
complex permittivity, 423–424
integral absorption, 448–449
length restrictions, 458
nonrigid dipole elastic longitudinal vibration and rigid dipole elastic reorientation, 427–432
research background, 394–396
rotary elasticity, dimensionless absorption, 456
strained-state energy, 436–437
translational vibration, dielectric response, 441–443
translation-reorientation independency, 449–450
- ice/water spectral models, 333–334
dipolar harmonic vibration, 343–346
librational state, 335–336
nonharmonic transverse vibrations, nonrigid dipoles, 346–348
overall permittivity, 348–350
Teixeira interpretation, 352–353
vibrational state, 336–337
- longitudinal/rotational vibration coupling, 368–369
- temperature influence, water spectra, 373–380
harmonic vibrations, 384–385
polarization, 388
two-fractional model, 387–388
- wideband water spectrum, room temperature conditions, 353–354
ordinary and heavy water, 360–367
- Hydrophilic ions, interface between two immiscible electrolyte solutions, transfer kinetics, 181–195
- Hydrophobic cation penetration, interface between two immiscible electrolyte solutions, 171–180
- Hyperspherical R-matrix with semiclassical outgoing waves (HRM-SOW), direct double photoionization, helium, 108–109
- Ice spectra. *See also* Water spectra
analytical calculations, 328–330
classical minimal action and phase transition estimations, 424–426
complex permittivity, 422–424
dielectric relaxation, molecular mechanisms, 335–351
hydrogen bond dipolar harmonic vibration, 343–346
librational state, 335–336

Ice spectra (*Continued*)

- low-frequency Raman spectrum, 350–351
- nonharmonic libration, deep well
 - permanent dipole, 337–343
- nonrigid dipole, nonharmonic transverse vibration, 346–348
- overall permittivity, 348–350
- Teixeira's data analysis, 352–353
- vibrational state, 336–337
- dimensionless inverse temperature, 414–417
- extended simple point charge (SPC/E) model, 505–509
- loss gap, 417–422
- low-temperature conditions, 410–414
- nonresonance spectrum, 407–410
- nonrigid dipoles, elastic longitudinal vibration, 426–458
 - autocorrelator transformation, 432–436
 - dimer kinetic energy, 438–440
 - equations of motion and spectral function, 443–448
- frozen translation:
 - librations, 451–452
 - rotations, 450–451
- heavy water, 452–453
- hydrogen bond:
 - strained-state energy, 436–437
 - translation/reorientation independency, 449–450
- integral absorption, 448–449
- rotary elasticity, 453–458
- translational vibration dielectric response, 440–443
- two-fraction dimer model, 426–432
- research background, 394–396
- resonance spectrum, high-temperature conditions, 396–407
 - calculations, 396–402
 - lattice mode spectroscopy, 402–407
- rigid dipoles, elastic reorientation, 426–458
 - autocorrelator transformation, 432–436
 - dimer kinetic energy, 438–440
 - equations of motion and spectral function, 443–448
- frozen translation librations, 451–452
- frozen translation rotations, 450–451
- heavy water, 452–453
- hydrogen bond strained-state energy, 436–437

- hydrogen bond translation/reorientation
 - independency, 449–450
- integral absorption, 448–449
- rotary elasticity, 453–458
- translational vibration dielectric response, 440–443
- two-fraction dimer model, 426–432
- transverse vibrations, water/ice spectra, 473–474
- two-fraction model, 333–334
- water spectra comparisons, 509–512
 - future research issues, 510–512
 - low-frequency transitions, 510
 - nonresonance far-IR ice spectrum, 510
 - resonance far-IR ice spectrum, 509–510
- Indirect processes, double photoionization:
 - complex atoms, 109–110
 - research background, 104–105
- Inertial moment, temperature influence, water spectra, 382–384
- Infinite temperature relaxation, calamitic liquid crystal fragility, 295–297
- Infrared (IR) array detector, mid-infrared pulse-shaping technology, ground-state vibrational dynamics, coherent control, 11–16
- Inherent structures:
 - calamitic liquid crystals, 303–306
 - thermotropic liquid crystals, energy landscapes, 297
- Initial instant of time, nonrigid dipoles,
 - nonharmonic vibrations, transverse spectral function, 466–471
- Inner-shell double photoionization, complex atoms, 124–128, 146
- Inner valence (IV) states, double photoionization, molecular Auger effect, 137–140, 147
- Integral absorption:
 - ice spectrum, 448–449
 - nonrigid dipoles, nonharmonic vibrations, transverse complex susceptibility, 468–471
- Intensity-modulated photocurrent modulation (IMPS), interface between two immiscible electrolyte solutions, electron transfer kinetics, 195–205
- Intensity parameter, nonrigid dipoles,
 - nonharmonic vibrations, transverse complex susceptibility, 470–471

- Interface between two immiscible electrolyte solutions (ITIES):
 - electron transfer kinetics, 195–205
 - future research issues, 208–209
 - historical development, 154–157
 - interfacial deposition, 205–208
 - ion transfer kinetics, 180–195
 - physicochemical properties, 154
 - research background, 154–157
 - thermodynamics, 155–157
- Interfacial deposition, interface between two immiscible electrolyte solutions, 205–208
- Interfacial roughness, interface between two immiscible electrolyte solutions, 168–180
- Intermolecular interactions, ice spectrum, high-temperature resonance spectra calculations, 402–407
- Internal field calculations, water summary permittivity, 489–491
- Intrinsic interfacial thickness, interface between two immiscible electrolyte solutions, 158–159
- Ion distribution, interface between two immiscible electrolyte solutions, 165–180
- Ion identity, interface between two immiscible electrolyte solutions, 166–167
- Ion penetration, interface between two immiscible electrolyte solutions, 172–173
- Ion transfer kinetics, interface between two immiscible electrolyte solutions, 180–195
- Isobaric heat capacity, thermotropic liquid crystals, supercooled regimes and glass transition, 256–258
- Isochors, thermotropic liquid crystals, energy landscapes, 298–303
- Isotropic dependence, wideband water spectrum, room temperature conditions, ordinary and heavy water, 361–367
- Isotropic independence:
 - static permittivity, ordinary/heavy water spectra, 355–357
 - wideband water spectrum, room temperature conditions, ordinary and heavy water, 361–367
- Isotropic-columnar (I-C) transition, thermotropic liquid crystals:
 - future research issues, 313
 - orientational time correlation function, 284–287
- Isotropic-nematic (I-N) phase transition:
 - calamitic liquid crystals, inherent structure orientation, 303–306
 - thermotropic liquid crystals:
 - calamitic liquid crystal fragility, 294–297
 - computer simulations, 275–287
 - future research issues, 311–313
 - Landau-de Gennes (LdG) theory, 268–270
 - Maier-Saupe theory, 267–268
 - microscopic view, 294
 - Onsager theory, 266–267
 - phase transition dynamics, 258–260
 - research background, 250–252
 - supercooled liquids, heterogeneous dynamics, 289–294
- Isotropic phases, liquid crystal-supercooled liquid orientational dynamics, 264–266
- Kauzmann temperature, thermotropic liquid crystals, supercooled regimes and glass transition, 257–258
- Kinetic energy, ice spectra dimer formation, 438–440
- Kirkwood correlation factor:
 - ice/water nonharmonic libration, 338–343
 - thermotropic liquid crystals, isotropic-nematic transition, 277–287
 - wideband water spectrum, room temperature conditions, static permittivity, isotropic independence, ordinary and heavy water, 356–357
- Kohlrausch-Williams-Watts (KWW) function, thermotropic liquid crystals, nonexponential relaxation, 262
- Kramers-Cronig rule, ice spectrum loss gap, 418–422
- Kronecker delta, thermotropic liquid crystals:
 - isotropic-nematic transition, 278–287
 - mesophase models, 254–255
- Krypton, double photoionization:
 - inner-shell DPI, 126–128
 - valence DPI, deep inner-shell thresholds, 122–124

- Ladder climbing:
 local control theory:
 photodissociation, 39
 selective mode excitation, 45–47
 mid-infrared pulse shaping, 2
 ground-state vibrational dynamics,
 coherent control, 8–16
- Ladder descending, local control theory,
 photoassociation, 48–51
- Landau approximation, thermotropic liquid
 crystals, Landau-de Gennes (LdG)
 theory, 269–270
- Landau-de Gennes (LdG) theory, thermotropic
 liquid crystals:
 isotropic-nematic phase transition, 258–260,
 268–270
 liquid crystal-supercooled liquid orientational
 dynamics, 264–266
 Mode coupling theory and, 271–273
- Landau-Zener theory:
 interface between two immiscible electrolyte
 solutions, electron transfer kinetics,
 198–205
 local control theory, predissociation
 dynamics, 40–42
- Langevin equations, interface between two
 immiscible electrolyte solutions, ion
 transfer kinetics, 192–195
- Langmuir models, interface between two
 immiscible electrolyte solutions,
 175–180
- Large-scale structure, ultrathin solid-fluid-solid
 films, 237–241
- Lattice gas interaction, interface between two
 immiscible electrolyte solutions,
 159–161
 double layer model, 167–180
- Lattice mode spectroscopy, ice spectrum, high-
 temperature resonance spectra
 calculations, 402–407
- Lebwohl-Lasher (LL) model, thermotropic
 liquid crystals, orientational
 time correlation function,
 281–287
- Legendre polynomial, thermotropic liquid
 crystals:
 isotropic-nematic transition,
 276–287
 orientational time correlation function,
 282–287
- Lennard-Jones interactions:
 interface between two immiscible electrolyte
 solutions, ion transfer kinetics,
 193–195
 thermotropic liquid crystals:
 computer simulations,
 274–275
 energy landscapes, 297–298
 supercooled liquids, dynamic similarity,
 288–289
- Librational band (LIB) fractions. *See also* Planar
 libration
 hydrogen bonds, longitudinal/rotational
 vibration coupling, 368–369
- ice spectrum:
 complex permittivity, 422–424
 “frozen” translations, 451–452
 hat well reorientation, 403–407
 high-temperature resonance spectra
 calculations, 400–402
 nonrigid dipole elastic longitudinal
 vibration and rigid dipole elastic
 reorientation, 429–432
 research background, 394–396
- ice/water spectral calculations, 330
 dielectric relaxation, 335–336
 molecular modeling, 330–333, 334,
 393–394
 nonharmonic permanent dipole libration,
 337–343
 overall permittivity, 348–350
 temperature influence, water spectrum,
 372–373
 hydrogen bond polarization, 388
 subensemble parameter, 380–384
 two-fraction model, 388
- wideband water spectrum, room temperature
 conditions, 353–354
 Raman spectra, 359–367
- Lifetime parameters:
 hydrogen bond harmonic vibrations, 344–346
 ice spectrum, high-temperature resonance
 spectra, 404–407
 temperature influence, water spectrum:
 transverse vibration, hydrogen bond
 molecules, 385–387
 two-fraction model, 387–388
- Light emitting diodes (LEDs), ultrathin
 solid-fluid-solid films,
 223–224

- Limiting displacement, nonrigid dipoles, nonharmonic vibrations, nonlinear equation of motion, 467–471
- Linear cross-correlations, mid-infrared pulse-shaping technology, ground-state vibrational dynamics, coherent control, 13–16
- Lippmann equation, interface between two immiscible electrolyte solutions, 165–180
- L-L interface:
 - electron transfer kinetics, 195–205
 - interface between two immiscible electrolyte solutions, 154–157
 - ion transfer kinetics, 185–195
 - structure, 157–180
- Local control theory (LCT):
 - analytical examples, 34–36
 - basic formulation, 32–33
 - coupled electronic/nuclear motion, 86–94
 - adiabatic ground state transfer, 89–90
 - combined electron-nuclear dynamics, 91–94
 - model parameters, 86–89
 - two-state model transfer, 90–91
- electronic excitation, 60–65
 - photodissociation and, 69–72
- Fermi's Golden Rule, 35–36
- free particle deceleration, 34
- future research issues, 94–95
- multiple potential wells, 51–60
 - double well potentials, 51–57
 - triple-well potentials, 57–60
- optical paralysis, 65–69
- photoassociation, 47–51
 - orientation role in, 83–86
- photodissociation, 36–44
 - direct photodissociation, 36–39
 - electronic transitions and, 69–72
 - predissociation dynamics, 40–42
 - selective mode excitation *vs.*, 42–44
- quantum *vs.* classical mechanics, 33–34
- rotational motion, 72–86
 - directional motion control, 73–78
 - excited-state fragmentation, 78–83
 - photoassociation orientation, 83–86
- selective mode excitation, 44–47
 - photodissociation *vs.*, 42–44
- theoretical background, 30–32
- Local oscillator methods, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 19–25
- Longitudinal vibrations:
 - classical theory applications, 369–371
 - hydrogen bonds, longitudinal/rotational vibration coupling, 368–369
- nonrigid dipoles, ice spectra, 426–458
 - autocorrelator transformation, 432–436
 - dimer kinetic energy, 438–440
 - equations of motion and spectral function, 443–448
 - frozen translation librations, 451–452
 - frozen translation rotations, 450–451
 - heavy water, 452–453
 - hydrogen bond strained-state energy, 436–437
 - hydrogen bond translation/reorientation independency, 449–450
 - integral absorption, 448–449
 - rotary elasticity, 453–458
 - translational vibration dielectric response, 440–443
 - two-fraction dimer model, 426–432
- temperature influence, water spectra, 376–380
- Long-time decay, thermotropic liquid crystals, Mode coupling theory (MCT), 271–273
- Lorentzian-shaped pulses:
 - ice spectrum:
 - autocorrelation functions, 432–436
 - complex permittivity, 423–424
 - nonrigid dipole elastic longitudinal vibration and rigid dipole elastic reorientation, 430–432
 - spectral function reduction, 447–448
 - ice/water spectral calculations:
 - hydrogen bond harmonic vibrations, 344–346
 - vibrational state, 336–337
- mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 17–25
- temperature influence, water spectrum, steady state and LIB subensemble, 381–384

- Loss gap:
ice spectrum, 417–422
nonrigid dipoles, nonharmonic vibrations,
transverse-vibration loss spectrum,
471–473
- Low-density domains (LDD), temperature
influence, water spectra, 389
- Low-density liquid (LDL) model, wideband
water spectrum, room temperature
conditions, 354–355
- Low-frequency Raman spectrum:
dielectric relaxation, spectral function,
autocorrelation function, 498–499
ice spectrum:
high-temperature resonance spectra
calculations, 401–402
nonresonance spectrum, 409–410
transition mechanisms, 510
ice/water spectral analysis, overall
permittivity, 350–351
temperature influence, water spectra,
375–380
wideband water spectrum, room temperature
conditions, ordinary and heavy water,
365–367
- Low-temperature spectrum, ice molecules:
computed absorption and loss factors,
414–417
minus-30°C prediction, 414–417
model parameterization, 410–414
T=100 K data, 410
- Magnetic bottle time-of-flight methods, double
photoionization, 113–116
inner-shell DPI, 127–128
- Magnetic susceptibility tensor, thermotropic
liquid crystals, Landau-de Gennes
(LdG) theory, 268–270
- Maier-Saupe theory, thermotropic liquid
crystals, 267–268
- Marcus model, interface between two
immiscible electrolyte solutions,
electron transfer kinetics, 197–205
- Mass-conserving kinematic equation,
ultrathin solid-fluid-solid films,
225–228
- Mass-transport limited current, ion transfer
kinetics, interface between two
immiscible electrolyte solutions,
184–195
- Maxwell equation, dielectric relaxation:
perturbed trajectories, 495–496
qualitative analysis, 492–495
- Mean-square interfacial displacement:
interface between two immiscible electrolyte
solutions, 158–180
thermotropic liquid crystals, supercooled
liquids, heterogeneous dynamics, 294
- Mechanical heterogeneity, ultrathin solid-fluid-
solid films, 239–241
- Memory function, thermotropic liquid crystals,
Mode coupling theory and, 272–273
- Mercury, double photoionization, valence DPI,
deep inner-shell thresholds, 124
- Mesogens:
calamitic liquid crystals, inherent structure
orientation, 303–306
thermotropic liquid crystals, 252–255
future research issues, 314
isotropic-nematic phase transition,
258–260, 279–287
- Microelectrochemical measurements at
expanding droplets (MEMED),
interface between two immiscible
electrolyte solutions, ion transfer
kinetics, 186–195
- Micron-scale interface between two immiscible
electrolyte solutions, ion transfer
kinetics, 183–195
- Microscopic models:
thermotropic liquid crystals, supercooled
liquids, heterogeneous dynamics,
294
ultrathin solid-fluid-solid films, 225–228
- Mid-infrared (mid-IR) pulse-shaping
technology:
automated two-dimensional infrared
spectroscopy, 16–25
basic principles, 4–5
future research issues, 26
germanium acoustooptic modulator
performance, 5–7
ground-state vibrational dynamics control,
7–16
local control theory, selective mode
excitation, 44–47
parametric transfer comparison, 7
research background, 2–4
- Minimal action estimation, ice and water
spectra, 424–426

- Mixed regions, interface between two immiscible electrolyte solutions, 158–159
- Mixed solvent layers, interface between two immiscible electrolyte solutions, 171–180
- Mode coupling theory (MCT), thermotropic liquid crystals, 270–273
 - supercooled liquids, dynamic similarity, 288–289
- Modified Poisson-Boltzmann (MPB) models, interface between two immiscible electrolyte solutions, 168–180
- Modulator characteristics, mid-infrared pulse shaping, 3–4
- Moed selective chemistry, local control theory, photodissociation via, 42–44
- Molecular dynamics (MDs):
 - extended simple point charge (SPC/E) model, 505–509
 - ice spectrum:
 - high-temperature resonance spectra calculations, 402–407
 - research background, 395–396
 - ice/water spectral calculations, 328–330
 - temperature influence, water spectrum, 372–373
 - interface between two immiscible electrolyte solutions, 158–180
 - thermotropic liquid crystals:
 - isotropic-nematic phase transition, 258–260
 - isotropic-nematic transition, 276–287
- Molecular excited-state photofragmentation, local control theory, 70–72
- Molecular mapping, double photoionization, 133
 - dissociative double photoionization, 140–143
- Molecular modeling:
 - ice spectra, high-temperature resonance spectra calculations, 400–402
 - ice/water spectral calculations, 330–333
 - thermotropic liquid crystals, 253–255
- Molecular motor concept, rotational motion, 73–78
- Molecules, double photoionization:
 - Auger effects, 134–140
 - direct DPI, 133–134
 - dissociative DPI, 140–141
 - mapping and electron distributions, 133
 - quantum yields, 131–133
 - resonant Auger phenomena, 144
 - TPEsCO and TOF-PEPECO spectra, 128–130
- Mole fraction, interface between two immiscible electrolyte solutions, 159–161
- Monte Carlo simulation, interface between two immiscible electrolyte solutions, 176–180
- Morse potential, mid-infrared pulse-shaping technology, ground-state vibrational dynamics, coherent control, 8–16
- Multielectron coincidence, double photoionization, inner-shell DPI, 126–128
- Multiple potential wells, local control theory, 51–60
 - double well potentials, 51–57
 - triple-well potentials, 57–60
- Navier-Stokes equation:
 - interface between two immiscible electrolyte solutions, ion transfer kinetics, 191–195
 - ultrathin solid-fluid-solid films, 225–228
- Near-edge structure, double photoionization:
 - inner-shell DPI, 128
 - resonant Auger phenomena, 143
- Nearly constant loss (NCL), thermotropic liquid crystals, liquid crystal-supercooled liquid orientational dynamics, 265–266
- Nematic glass, future research issues, 314
- Nematic phase:
 - calamitic liquid crystals, inherent structure orientation, 304–306
 - thermotropic liquid crystals:
 - future research issues, 313
 - Maier-Saupe theory, 268
 - thermotropic mesophases and order parameters, 252–255
- Nematic-to-smectic-A (NA) transition, thermotropic liquid crystals, orientational and translational order, 306–311
- Neon, double photoionization, valence DPI, deep inner-shell thresholds, 123–124

- Nernst-Donnan equation, interface between two immiscible electrolyte solutions, 156–157
ion transfer kinetics, 186–195
- Nernst equation, interface between two immiscible electrolyte solutions, electron transfer kinetics, 196–205
- Nernst-Planck equation, interface between two immiscible electrolyte solutions, ion transfer kinetics, 190–195
- Net driving, interface between two immiscible electrolyte solutions, electron transfer kinetics, 200–205
- Neutral molecules, ice/water spectral calculations, molecular modeling, 331–333
- Nitrogen:
direct double photoionization, 134
dissociative double photoionization, 141–143
- Nonergodic behavior, thermotropic liquid crystals, supercooled liquids, heterogeneous dynamics, 294
- Nonexponential relaxation, thermotropic liquid crystals, 262
- Non-Gaussian parameter (NGP), thermotropic liquid crystals, supercooled liquids, heterogeneous dynamics, 290–294
- Nonharmonic libration:
ice/water spectral calculations, deep well permanent dipoles, 337–343
wideband water spectrum, room temperature conditions, 360–367
- Nonharmonic vibrations:
ice/water spectral calculations, nonrigid dipoles, 346–348
temperature influence, water spectrum, hydrogen bond molecules, 385–387
transverse vibration, nonrigid dipoles:
autocorrelator susceptibility, 460–463
frequency factor and loss spectrum, 471–473
nonlinear equation of motion, 463–464
research background, 459–460
spectral function, 464–471
water/ice spectra, 473–474
wideband water spectrum, room temperature conditions, ordinary and heavy water, 360–367
- Nonlinear equation of motion, nonrigid dipoles, nonharmonic vibrations, 463–464
- Nonlinear spectroscopy:
interface between two immiscible electrolyte solutions, 161–180
mid-infrared pulse-shaping technology, 16–25
- Nonresonance spectrum:
far-infrared ice spectra, 510
ice spectra, 407–410
- Nonrigid dipoles:
ice spectra:
elastic longitudinal vibration, 426–458
autocorrelator transformation, 432–436
dimer kinetic energy, 438–440
equations of motion and spectral function, 443–448
frozen translation librations, 451–452
frozen translation rotations, 450–451
heavy water, 452–453
hydrogen bond strained-state energy, 436–437
hydrogen bond translation/reorientation independency, 449–450
integral absorption, 448–449
rotary elasticity, 453–458
translational vibration dielectric response, 440–443
two-fraction dimer model, 426–432
ice/water spectral calculations:
molecular modeling, 331–333
nonharmonic transverse vibrations, 346–348
vibrational state, 336–337
nonharmonic transverse vibration:
autocorrelator susceptibility, 460–463
frequency factor and loss spectrum, 471–473
nonlinear equation of motion, 463–464
research background, 459–460
spectral function, 464–471
water/ice spectra, 473–474
- Normalization constant, thermotropic liquid crystals, Landau-de Gennes (LdG) theory, 269–270
- Nuclear dynamics, local control theory, 91–94
- Nucleation response:
interface between two immiscible electrolyte solutions, interfacial deposition, 205–208
thermotropic liquid crystals, orientational time correlation function, 285–287

- One-fraction model, water molecular modeling, 330–333
- One-particle approximation:
 - ice/water spectral calculations, 329–330
 - temperature influence, water spectrum, 373–380
- Onsager theory, thermotropic liquid crystals, 266–267
- Onset temperatures, thermotropic liquid crystals, energy landscapes, 302–303
- Optical constants, ice spectra, high-temperature resonance spectra calculations, 396–402
- Optical heterodyne-detected optical Kerr effect (OHD-OKE) measurements, thermotropic liquid crystals:
 - isotropic-nematic phase transition, 259–260
- Mode coupling theory, 273
- supercooled isotropic phases, 264–266
- supercooled liquids, dynamic similarity, 289
- Optical Kerr effect (OKE), thermotropic liquid crystals, 250–252
 - future research issues, 311–313
 - isotropic-nematic phase transition, 258–260
 - liquid crystal-supercooled liquid orientational dynamics, 265–266
 - Mode coupling theory (MCT), 271–273
 - orientational time correlation function, 279–287
 - supercooled liquids, dynamic similarity, 289
- Optical paralysis, local control theory, 32
 - applications, 65–69
- Optical parametric amplifier (OPA), mid-infrared (mid-IR) pulse-shaping technology, 4–5
- Order parameters, thermotropic liquid crystals, 252–255
- Organic solvent thin film, interface between two immiscible electrolyte solutions, ion transfer kinetics, 188–195
- Orientational dynamics:
 - calamitic liquid crystals, inherent structure orientation, 303–306
 - ice spectrum, complex permittivity, 423–424
 - thermotropic liquid crystals:
 - energy landscapes, 301–303
 - Maier-Saupe theory, 267–268
 - orientational time correlation function, 279–287
 - supercooled isotropic phases, 264–266
 - supercooled liquids, heterogeneous dynamics, 289–294
 - translational order and, 306–311
- Orientational time correlation function (OTCF), thermotropic liquid crystals:
 - isotropic-nematic transition, 275–287
 - supercooled liquids, dynamic similarity, 289
- Ornstein-Zernike form, thermotropic liquid crystals:
 - isotropic-nematic transition, 278–287
- Landau-de Gennes (LdG) theory, 270
- Overpumping problems, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 23–25
- Oxygen atoms:
 - dissociative double photoionization, 140–143
 - hat well, special configuration and permittivity, dipole precession confinement, 481–482
- Packet waveforms, local control theory, photoassociation, 47–51
- Pairwise electron correlation, double photoionization, 145–146
- Palladium deposition, interface between two immiscible electrolyte solutions, 206–208
- Parametric transfer, mid-infrared pulse-shaping technology, germanium acoustooptic modulator, 7
- Particle transfer, double well potential, local control theory, 51–57
- Pendular state, local control theory, excited-state fragmentation, 82–83
- Periodic function, nonrigid dipoles, nonharmonic vibrations, nonlinear equation of motion, 467–471
- Permanent dipoles, nonharmonic libration, 337–343
- Permittivity:
 - dielectric relaxation, qualitative analysis, 491–495
 - extended simple point charge (SPC/E) model, 506–509
 - hat well, spatial configuration:
 - hat potential evolution, 474–479
 - planar libration, 479–482
 - spectral function, 482–489
 - water molecules, 489–491

- Permittivity: (*Continued*)
 hydrogen bonds, longitudinal/rotational vibration coupling, 368–369
 ice spectra, formulas for, 422–424
 ice/water spectral analysis:
 hydrogen bond harmonic vibrations, 344–346
 low-frequency Raman spectrum, 350–351
 nonharmonic libration, 338–343
 overall vibrational contribution, 348–350
 temperature influence, water spectrum, 372–373
 wideband water spectrum, room temperature conditions, ordinary and heavy water, 364–367
- Perturbed trajectories, dielectric relaxation, power consumption, 495–496
- Phase and intensity masks, mid-infrared pulse-shaping technology, ground-state vibrational dynamics, coherent control, 13–16
- Phase-matching geometries, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 19–25
- Phase-separating blend, ultrathin solid-fluid-solid films, 243–244
- Phase transition dynamics:
 ice and water spectra, 424–426
 thermotropic liquid crystals:
 future research issues, 314
 isotropic-nematic phase transition, 258–260
 Onsager theory, 266–267
 orientational time correlation function, 285–287
- Phase variables, nonrigid dipoles, nonharmonic vibrations, transverse complex susceptibility, 468–471
- Photoassociation, local control theory, 47–51
 orientation role, 83–86
- Photocurrent density, interface between two immiscible electrolyte solutions, electron transfer kinetics, 201–205
- Photodissociation, local control theory, 36–44
 direct photodissociation, 36–39
 electronic transitions and, 69–72
 predissociation dynamics, 40–42
 selective mode excitation vs., 42–44
- Photon echo, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 19–25
- Planar libration, hat well, special configuration and permittivity:
 precession approximation, 475–479
 spectral functions, 482–489
 variants, 479–482
- Poiseuille flow behavior, ultrathin solid-fluid-solid films, 225–228
- Poisson-Boltzmann equation, interface between two immiscible electrolyte solutions, 165–180
- Poisson brackets, local control theory, quantum vs. classical mechanics, 33–34
- Polarity:
 interface between two immiscible electrolyte solutions, 163–180
 temperature influence, water spectra, hydrogen bonds, 388
- Population inversion and selectivity, mid-infrared pulse-shaping technology, ground-state vibrational dynamics, coherent control, 10–16
- Porphyrin complexes, interface between two immiscible electrolyte solutions, electron transfer kinetics, 200–205
- Post-collision interaction (PCI), double photoionization:
 indirect processes, 110
 inner-shell DPI, 127–128
- Potential-dependent adsorption equilibrium, interface between two immiscible electrolyte solutions, 173–180
- Potential energy curves:
 interface between two immiscible electrolyte solutions, 165–180
 local control theory:
 electronic excitation, 60–65
 optical paralysis, 67–69
 photoassociation, 47–51
 rotational motion, 73–78
 thermotropic liquid crystals, energy landscapes, 299–303
- Potential surfaces, double well potential, local control theory, 53–57
- Power absorption coefficient, temperature influence, water spectra, 389–390
- Power law decay, thermotropic liquid crystals: energy landscapes, 303

- future research issues, 313
- isotropic-nematic phase transition, 279–287
- supercooled liquids, dynamic similarity, 288–289
- Precession approximation, hat well, special configuration and permittivity:
 - dipole confinement, hydrogen bond, 479–481
 - planar libration, 475–479
- Predissociation dynamics, local control theory, photodissociation, 40–42
- Pressure behavior, ultrathin solid-fluid-solid films, 225–228
- Probability density function, local control theory, rotational motion, 75–78
- Proportionality relationship, low-temperature spectrum, ice molecules, dimensionless inverse temperatures, 415–417
- Protrusion-extrusion model, interface between two immiscible electrolyte solutions, ion transfer kinetics, 190–195
- Pseudo-first-order models, interface between two immiscible electrolyte solutions, electron transfer kinetics, 197–205
- Pseudonematic domains, thermotropic liquid crystals, supercooled liquids, heterogeneous dynamics, 289–294
- Pulsed laser spectroscopy, interface between two immiscible electrolyte solutions, electron transfer kinetics, 201–205
- Pulse-shaping technology:
 - local control theory, selective mode excitation, 45–47
 - research background, 1–4
- Pump-probe spectrum:
 - local control theory, rotational motion, 76–78
 - mid-infrared pulse-shaping technology:
 - ground-state vibrational dynamics, coherent control, 11–16
 - two-dimensional infrared spectroscopy, 21–25
- Qualitative analysis, dielectric relaxation, 491–495
- Quantitative agreement, temperature influence, water spectra, 377–380
- Quantum-break factor, temperature influence, water spectra, harmonically vibrating hydrogen bond molecules, 384–385
- Quantum mechanics:
 - harmonic longitudinal vibrations, 370–371
 - local control theory, 33–34
 - direct photodissociation, 37–39
 - photoassociation and, 51
 - rotational motion, 75–78
- Quantum yields, double photoionization:
 - direct DPI, 133–134
 - molecules, 131–133
- Quasi-elastic light scattering (QELS), interface between two immiscible electrolyte solutions, 179–180
- Quasi-reversible first-order process, interface between two immiscible electrolyte solutions, transfer kinetics, 181–195
- Rabi oscillations, local control theory:
 - coupled two-state model, 91
 - electronic excitation, 63–65
- Radial distribution function:
 - calamitic liquid crystals, inherent structure orientation, 305–306
 - ice spectrum, nonrigid dipole elastic longitudinal vibration and rigid dipole elastic reorientation, 430–432
- Radiofrequency (RF) signals, mid-infrared (mid-IR) pulse-shaping technology, 5–6
- RAK model, wideband water spectrum, room temperature conditions, 354–355
- Raman spectrum:
 - ice spectrum, high-temperature resonance spectra, 407
 - ice/water spectral analysis, overall permittivity, 350–351
 - temperature influence, water spectra, 375–380
 - transverse vibration and, 388–389
 - wideband room temperature conditions, ordinary and heavy water, 358–367
- Rate constants, interface between two immiscible electrolyte solutions:
 - electron transfer kinetics, 196–205
 - ion transfer kinetics, 189–195
- Reaction coordinate density, double well potential, local control theory, 55–57
- Recurrence equations, nonrigid dipoles, nonharmonic vibrations, transverse spectral function, 464–471

- Redox reactions, interface between two
immiscible electrolyte solutions,
electron transfer kinetics, 197–205
- Refraction index:
ice spectrum loss gap, 419–422
wideband water spectrum, room temperature
conditions, ordinary and heavy water,
361–363
- Relaxation times:
ice spectrum, loss gap, 420–422
ice/water spectra, 335–351
hydrogen bond dipolar harmonic vibration,
343–346
librational state, 335–336
low-frequency Raman spectrum, 350–351
molecular models, 334
nonharmonic libration, deep well
permanent dipole, 337–343
nonrigid dipole, nonharmonic transverse
vibration, 346–348
overall permittivity, 348–350
Teixeira's data analysis, 352–353
vibrational state, 336–337
temperature influence, water spectra,
373–380
steady-state parameters, 382–384
- Reorientation lifetime:
ice spectrum, translation-reorientation
independency, 449–450
temperature influence, water spectrum,
hydrogen bond molecules, 386–387
- Resolution parameters, germanium acoustooptic
modulator, 5–7
- Resonance spectra:
far-infrared ice spectra, 509–510
ice spectra, 396–407
calculations, 396–402
lattice mode spectroscopy, 402–407
nonrigid dipole elastic longitudinal vibration
and rigid dipole elastic reorientation,
426–432
- Resonant stimulated Raman scattering process
(RISRS), local control theory, optical
paralysis, 67–69
- Restricted primitive models, interface between
two immiscible electrolyte solutions,
168–180
- Rigid dipoles:
ice spectra, elastic reorientation, 426–458
autocorrelator transformation, 432–436
dimer kinetic energy, 438–440
equations of motion and spectral function,
443–448
frozen translation librations, 451–452
frozen translation rotations, 450–451
heavy water, 452–453
hydrogen bond strained-state energy,
436–437
hydrogen bond translation/reorientation
independency, 449–450
integral absorption, 448–449
rotary elasticity, 453–458
translational vibration dielectric response,
440–443
two-fraction dimer model, 426–432
- ice/water spectral models:
hydrogen bond harmonic vibrations,
344–346
vibrational state, 336–337
temperature influence, water spectrum,
373–380
- Rotary elasticity, ice spectra, 453–458
angular vibration dependence, 457
dimensionless absorption, 453–456
vibration spectrum approximation, 457–458
- Rotary force constant:
ice spectra:
angular vibration dependence, 457
frozen translation librations, 451–452
ice spectrum:
frozen rotation translations, 450–451
rotary elasticity, dimensionless absorption,
453–456
translation-reorientation independency,
449–450
- Rotational invariant functions, thermotropic
liquid crystals, isotropic-nematic
transition, 278–287
- Rotational vibrations:
hydrogen bonds, longitudinal/rotational
vibration coupling, 368–369
ice spectra, dielectric response, 440–443
ice spectrum, autocorrelation functions, 436
ice/water spectral models, 353
local control theory, 72–86
directional motion control, 73–78
excited-state fragmentation, 78–83
photoassociation orientation, 83–86
temperature influence, water spectrum, two-
fraction model, 388

- Rydberg states, double photoionization:
 - deep inner-shell thresholds, 121–124
 - dissociative double photoionization, 141–143
 - molecular Auger effect, 137–140, 146
- Scanning electrochemical microscope (SECM),
 - interface between two immiscible electrolyte solutions:
 - electron transfer kinetics, 195–205
 - ion transfer kinetics, 185–195
- Scattering phenomena, local control theory,
 - photoassociation, 47–51
- Schrödinger equation, local control theory,
 - 32–33
 - combined electron-nuclear dynamics, 91–94
 - coupled electronic/nuclear motion, 87–89
 - optical paralysis, 66–69
- Second harmonic generation (SHG)
 - spectroscopy, interface between two immiscible electrolyte solutions, 179–180
- Seed pulse characteristics, local control theory:
 - electronic excitation, 62–65
 - optical paralysis, 66–69
- Selective mode excitation, local control theory,
 - 44–47
 - photodissociation via, 42–44
- Self-consistency condition, thermotropic liquid crystals, Maier-Saupe theory, 268
- Self-diffusion coefficients:
 - ice/water spectral models, 353
 - thermotropic liquid crystals, orientational and translational order, 306–311
- Shake-off process, direct double
 - photoionization, helium, 108–109
- Shear viscosity, thermotropic liquid crystals:
 - orientational time correlation function, 285–287
 - supercooled temperature dependence, 261–262
- Short-time power law, thermotropic liquid crystals, orientational time correlation function, 279–287
- Sine theorem, strained-state energy, hydrogen bonds, ice spectrum, 437
- Single-axis spectra, double photoionization,
 - data presentation, 116–120
- Single-laser microdroplet experiment, interface
 - between two immiscible electrolyte solutions, ion transfer kinetics, 186–195
- Single-particle orientational dynamics,
 - thermotropic liquid crystals:
 - isotropic-nematic phase transition, 279–287
 - supercooled liquids, heterogeneous dynamics, 294
- Single penetrating field analyzer, threshold
 - photoelectrons coincidence apparatus, 112–113
- Smectic phases:
 - calamitic liquid crystals, inherent structure orientation, 304–306
 - thermotropic liquid crystals, 253–255
 - orientational and translational order, 309–311
- Soft condensed matter systems, thermotropic liquid crystals, 250–251
- Solid capping layers:
 - substrate-solid-solid films, 221–222
 - ultrathin solid-fluid-solid films, 219–220
 - evolutionary models, 234–237
 - surface undulation control, 223–224
- Spatial configuration:
 - hat well, summary permittivity:
 - hat potential evolution, 474–479
 - planar libration, 479–482
 - spectral function, 482–489
 - water molecules, 489–491
 - ultrathin solid-fluid-solid films, 229–231
- Spectral analysis:
 - calculations, analytical theory, 328–330
 - interface between two immiscible electrolyte solutions, electron transfer kinetics, 201–205
 - mid-infrared (mid-IR) pulse-shaping technology, 5–6
- Spectral functions:
 - dielectric relaxation, autocorrelation function, 496–499
 - dielectric relaxation, complex susceptibility,
 - Gross collision model, 502–505
 - hat well, special configuration and permittivity, 482–489
 - librator approximations, 482–485
 - ice complex permittivity, 422–424
 - ice spectrum:
 - analytical formula, 444–447
 - frozen rotation:
 - libration, 451–452

- Spectral functions: (*Continued*)
 translation, 450–451
 hydrogen bond translation-reorientation
 independency, 449–450
 nonrigid dipole elastic longitudinal
 vibration and rigid dipole elastic
 reorientation, 432–436
 sum of Lorentz line reduction, 447–448
 nonrigid dipoles, nonharmonic vibrations,
 transverse functions, 464–471
- Stark shift, local control theory:
 predissociation dynamics, 41–42
 rotational motion, 76–78
- State-selective photoassociation, local control
 theory, 50–51
- Static correlation, thermotropic liquid crystals,
 isotropic-nematic transition,
 277–287
- Static parameters, temperature influence, water
 spectrum, 373–380
- Static permittivity:
 ice/water nonharmonic libration, 339–343
 temperature-dependent water spectra,
 382–384
 wideband water spectrum, room temperature
 conditions, isotropic independence,
 355–357
- Steady-state parameters:
 ice spectrum, research background, 395–396
 temperature influence, water spectra,
 380–384
- Stiffness properties, ultrathin solid-fluid-solid
 films, 239–241
- Stimulated Raman adiabatic passage (STIRAP)
 methodology:
 local control theory, 31–32
 mid-infrared pulse shaping, 2–3
- Stokes-Einstein (SE) relation, thermotropic
 liquid crystals, dynamic decoupling
 phenomena, 263–264
- Strained-state energy, hydrogen bonds, ice
 spectrum, 436–437
- Stress-guided self assembly, ultrathin solid-
 fluid-solid films, 223–224
- Stretching elastic forces, ultrathin solid-fluid-
 solid films, 232–237
- Subpulse properties, local control theory,
 electronic excitation, 63–65
- Substrate-fluid-solid films, undulation growth,
 220–221
- Substrate-solid-solid films, undulation growth,
 221–222
- Supercooled liquid regimes:
 ice/water spectral analysis, nonharmonic
 transverse vibrations, nonrigid
 dipoles, 348
 temperature influence, water spectra,
 378–380
 thermotropic liquid crystals, 260–264
 calamitic liquid crystal fragility,
 294–297
 dynamic decoupling, 262–264
 dynamic similarity, computer simulations,
 288–297
 glass transition, 255–258
 heterogeneous dynamics, 289–294
 liquid crystal-supercooled liquid
 orientational dynamics, 264–266
 liquids and glass transition, 255–258
 microscopic view, 294
 nonexponential relaxation, 262
 power law relaxation, 288–289
 shear viscosity temperature dependence,
 261–262
- Surface corrugations, ultrathin solid-fluid-solid
 films, 238–241
- Surface undulation control, ultrathin solid-fluid-
 solid films, 223–224
 evolutionary models, 231–237
- s-wave scattering, local control theory,
 photoassociation orientation, 83–86
- Symmetrical distribution function, thermotropic
 liquid crystals, Maier-Saupe theory,
 267–268
- Synchrotron radiation pulses, double
 photoionization:
 time-of-flight photoelectron-photoelectron
 coincidence (TOF-PEPECO)
 technique, 115–116
 valence DPI, deep inner-shell thresholds,
 123–124
- Tafel expression, interface between two
 immiscible electrolyte solutions,
 electron transfer kinetics,
 195–205
- T-band spectrum:
 ice spectrum:
 high-temperature resonance spectra,
 404–407

- nonrigid dipole elastic longitudinal
 - vibration and rigid dipole elastic reorientation, 429–432
- rotary elasticity, dimensionless absorption, 454–456
- temperature influence, water spectra, 375–380
- water molecular modeling, 330–333
- Teixeira interpretation, ice/water spectral analysis, 352–353
- Temperature dependence:
 - ice spectra, minimal action and phase transition estimation, 424–426
 - loss spectra, extended simple point charge (SPC/E) model, 507–509
- thermotropic liquid crystals:
 - Landau-de Gennes (LdG) theory, 269–270
 - Maier-Saupe theory, 267–268
 - supercooled regime, 261–262
- water spectra:
 - calculations, 373–380
 - fitted parameters, 380–387
- hydrogen bond:
 - harmonic vibration, 384–385
 - polarization, 388
 - transverse vibration, 385–387
- librational state, 393–394
- minimal action and phase transition
 - estimation, 424–426
- model parameters, 389–393
- power absorption coefficient, 389
- research background, 372–373
- steady state and librational band
 - subensemble, 380–384
- transverse vibration and, 388–389
- two-fractional model, 387–388
- vibrational state mechanisms, 394
- Tetracyanoquinodimethane (TCNQ)/ferrocyanide, interface between two immiscible electrolyte solutions, electron transfer kinetics, 200–205
- TGN model:
 - hydrogen bonds, longitudinal/rotational vibration coupling, 368–369
- ice/water spectral models, hydrogen bond
 - harmonic vibrations, 344–346
- Thermodynamics:
 - ice spectrum, research background, 395–396
 - interface between two immiscible electrolyte solutions, 155–157
 - interfacial deposition, 206–208
 - ion transfer kinetics, 187–195
- Thermotropic liquid crystals:
 - computer simulations, 273–297
 - calamitic liquid crystals, 303–306
 - energy landscapes, 297–311
 - isotropic-nematic transition dynamics, 275–287
 - model systems, 274–275
 - orientational and dynamical signatures, 306–311
 - supercooled liquids, 288–297
 - future research issues, 311–314
 - isotropic-nematic phase transition, 258–260
 - Landau-de Gennes theory, 268–270
 - Maier-Saupe theory, 267–268
 - mesophases and order parameters, 252–255
 - mode coupling theory analysis, 270–273
 - Onsager theory, 266–267
 - orientational dynamics, isotropic phase and supercooled liquids, 264–266
 - research background, 250–252
 - supercooled regime dynamics, 260–264
 - calamitic liquid crystal fragility, 294–297
 - computational methods, 288–297
 - dynamic decoupling, 262–264
 - heterogeneous dynamics, 289–294
 - liquids and glass transition, 255–258
 - microscopic view, 294
 - nonexponential relaxation, 262
 - power law relaxation, 288–289
 - shear viscosity temperature dependence, 261–262
- Thin film technology, interface between two immiscible electrolyte solutions, ion transfer kinetics, 188–195
- Threshold law, direct double photoionization, helium, 106–109
- Threshold photoelectrons coincidence (TPESCO), double photoionization:
 - basic principles, 111–113
 - characteristics of, 116
 - data presentation, 116–120
 - inner-shell DPI, 127–128
 - molecules, 128–130
 - quantum yields, 131–133
 - valence DPI, deep inner-shell thresholds, 121–124

- THz spectra:
 temperature influence, water spectra,
 376–380
 transverse vibration and, 388–389
 wideband water spectrum, room temperature
 conditions, ordinary and heavy water,
 365–367
- Time dependence, ice spectrum, spectral
 function coordinates, 448
- Time-dependent close coupling (TDCC), direct
 double photoionization, helium,
 108–109
- Time-dependent Schrödinger equation, local
 control theory, 32–33
 combined electron-nuclear dynamics, 91–94
 optical paralysis, 66–69
- Time-of-flight mass spectrometry (TOF-MS),
 double photoionization:
 energetic electron pair analysis, 113–116
 threshold photoelectrons coincidence
 apparatus, 112–113
- Time-of-flight photoelectron-photoelectron
 coincidence (TOF-PEPECO)
 technique, double photoionization:
 characteristics of, 116
 correlation energy, 145–146
 molecules, 128–130
 research background, 114–116
 resonant Auger phenomena, 143
 valence DPI, deep inner-shell thresholds,
 121–124
- Time-reversed etalon formation, mid-infrared
 pulse-shaping technology, two-
 dimensional infrared spectroscopy,
 22–25
- Timescale calculations, ice spectrum, complex
 permittivity, 422–424
- Toluene, double photoionization, molecular
 Auger effect, 137–138
- Total internal reflectance (TIR) fluorescence
 spectroscopy, interface between two
 immiscible electrolyte solutions,
 161–180
 electron transfer kinetics, 201–205
- Tracking control:
 dielectric relaxation, perturbed trajectories,
 495–496
 local control theory, 31–32
- Transfer dynamics, local control theory:
 coupled two-state model, 90–91
 double well potential, 52–57
 triple-well potentials, 57–60
- Translational ban maximum, wideband water
 spectrum, room temperature
 conditions, ordinary and heavy water,
 364–367
- Translational degrees of freedom (TDOF):
 ice spectrum, autocorrelation functions,
 436
 thermotropic liquid crystals:
 orientational dynamics and, 306–311
 supercooled liquids, dynamic similarity,
 288–289
 supercooled liquids, heterogeneous
 dynamics, 290–294
- Translational spectral functions, ice spectrum,
 complex permittivity, 423–424
- Translational vibrations, ice spectra:
 dielectric response, 440–443
 dimer kinetics, 438–440
 frozen rotation, 450–451
 frozen translation librations, 451–452
 hydrogen bond reorientation, 449–450
- Transverse intensity and frequency,
 temperature influence, water spectra,
 383–384
- Transverse optic-longitudinal optic splitting, ice
 spectrum, 407
- Transverse vibrations (TV):
 nonrigid dipoles:
 ice/water spectral calculations, 346–348
 nonharmonic vibrations:
 autocorrelator susceptibility, 460–463
 frequency factor and loss spectrum,
 471–473
 nonlinear equation of motion, 463–464
 research background, 459–460
 spectral function, 464–471
 water/ice spectra, 473–474
 temperature influence, water spectrum,
 373–380
 hydrogen bond molecules, 385–387
 molecular bending motion, 388–389
 wideband water spectrum, room temperature
 conditions, ordinary and heavy water,
 360–367
- Trilayer films, ultrathin solid-fluid-solid films,
 evolutionary models, 235–237
- Triple-well potentials, double well potential,
 57–60

- Triply differential cross section (TDCS), direct
 - double photoionization, helium, 107–109
- Two-channel electron multiplier, threshold photoelectrons coincidence apparatus, double photoionization, 112–113
- Two-dimensional infrared (2D IR) spectroscopy:
 - mid-infrared pulse shaping, research background, 2–4
 - mid-infrared pulse-shaping technology, automated processes, 16–25
- Two-dimensional nuclear magnetic resonance (NMR) spectroscopy, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy *vs.*, 17–25
- Two-fraction model:
 - ice/water spectra:
 - dielectric relaxation, 335
 - librational state, 335–336
 - vibrational state, 336–337
 - nonrigid dipole elastic longitudinal vibration and rigid dipole elastic reorientation, 426–432
 - temperature influence, water spectrum, 387–388
 - water molecular modeling, 330–333
- Two-vibrator systems, mid-infrared pulse-shaping technology, two-dimensional infrared spectroscopy, 18–25
- Ultrashort light pulses, double photoionization, complex atoms, 121–124
- Ultrathin solid-fluid-solid films:
 - computational details, 229–231
 - driving instabilities, alternative mechanisms, 222–223
 - evolution, 231–237, 245–246
 - functional derivatives, 244–245
 - future research issues, 241–244
 - methodology, 224–231
 - research background, 217–220
 - substrate-fluid-solid films, 220–221
 - substrate-solid-solid films, 221–222
 - surface undulation control, 223–224
 - theoretical principles, 225–228
 - undulation manipulation and final structure, 237–241
- Uncoupled processes, ice/water spectral models, 334
- Undisturbed variables, dielectric relaxation, complex susceptibility, Gross collision model, 499–502
- Undulation growth:
 - substrate-fluid-solid films, 220–221
 - substrate-solid-solid films, 221–222
 - ultrathin solid-fluid-solid films, manipulation and final structure, 237–241
- Unperturbed molecular systems, local control theory, electronic excitation, 60–65
- Valence double photoionization, deep inner-shell thresholds, 121–124
- Van der Waals forces:
 - substrate-fluid-solid films, 220–221
 - thermotropic liquid crystals, Maier-Saupe theory, 267–268
 - ultrathin solid-fluid-solid films, 218–220
 - computational analysis, 230–231
 - driving instabilities, 222–223
 - evolutionary models, 231–237
 - pressure behavior, 226–228
 - undulation orientation, 242–244
- Vapor-water phase transition, ice and water spectra, 424–426
- Verlet Leap-frog algorithm, extended simple point charge (SPC/E) model, 506–509
- Verwey-Niessen model, interface between two immiscible electrolyte solutions, 166–180
- Vibrational band (VIB) fractions:
 - hydrogen bonds, longitudinal/rotational vibration coupling, 368–369
- ice spectrum:
 - approximations, 457–458
 - high-temperature resonance spectra calculations, 400–402
 - nonrigid dipole elastic longitudinal vibration and rigid dipole elastic reorientation, 429–432
 - research background, 394–396
 - rotary elasticity, dimensionless absorption, 454–456
 - T- and V-band mechanisms, 404–407
- ice/water spectral calculations, 330
 - dielectric relaxation, 336–337
 - hydrogen bond harmonic vibrations, 343–346

- Vibrational band (*Continued*)
 molecular modeling, 331–333
 overall permittivity, 348–350
 ice/water spectral models, 333–334
 temperature influence, water spectrum, 372–373
 hydrogen bond polarization, 388
 wideband water spectrum, room temperature conditions, 353–354
- Vibrationally mediated photodissociation, local control theory, photodissociation via, 42–44
- Vibrational sum-frequency spectroscopy, interface between two immiscible electrolyte solutions, 162–180
- Virial approximation, thermotropic liquid crystals, Onsager theory, 266–267
- Vogel-Fulcher-Tammann (VFT) equation: calamitic liquid crystal fragility, 295–297
 thermotropic liquid crystals, shear viscosity temperature dependence, 261–262
- Volt-fluorograms, interface between two immiscible electrolyte solutions, ion transfer kinetics, 183–195
- Wannier theory, double photoionization, valence DPL, deep inner-shell thresholds, 122–124
- Water-carbon tetrachloride, interface between two immiscible electrolyte solutions, 159–180
- Water-dichloroethane (DCE), interface between two immiscible electrolyte solutions, 158–180
 interfacial deposition, 205–208
 ion transfer kinetics, 184–195
- Water spectra:
 classical minimal action and phase transition estimations, 424–426
 dielectric relaxation, molecular mechanisms, 335–351
 hydrogen bond dipolar harmonic vibration, 343–346
 librational state, 335–336
 low-frequency Raman spectrum, 350–351
 nonharmonic libration, deep well permanent dipole, 337–343
 nonrigid dipole, nonharmonic transverse vibration, 346–348
 overall permittivity, 348–350
 Teixeira's data analysis, 352–353
 vibrational state, 336–337
 extended simple point charge (SPC/E) model, 505–509
 frequency ranges, 328–330
 hat well, special configuration and permittivity, 489–491
 ice spectra comparisons, 509–512
 future research issues, 510–512
 low-frequency transitions, 510
 nonresonance far-IR ice spectrum, 510
 resonance far-IR ice spectrum, 509–510
 molecular modeling, 330–333
 temperature dependence:
 calculations, 373–380
 fitted parameters, 380–387
 hydrogen bond:
 harmonic vibration, 384–385
 polarization, 388
 transverse vibration, 385–387
 librational state, 393–394
 model parameters, 389–393
 power absorption coefficient, 389
 research background, 372–373
 steady state and librational band subensemble, 380–384
 transverse vibration and, 388–389
 two-fractional model, 387–388
 vibrational state mechanisms, 394
 transverse vibrations, water/ice spectra, 473–474
 two-fraction model, 333–334
 wideband water spectrum, room temperature conditions:
 mixed model (water, LDL, HDL) concepts, 354–355
 ordinary and heavy water, 355–367
 research background, 353–354
 static permittivity, isotropic independence, 355–357
- Water transfer. *See also* Aqueous spectra; Ice/water spectra
 interface between two immiscible electrolyte solutions, ion transfer kinetics, 194–195
- Waveform characteristics, mid-infrared pulse-shaping technology, ground-state vibrational dynamics, coherent control, 10–16

- Wave geometry, mid-infrared pulse-shaping
 technology, two-dimensional infrared
 spectroscopy, 19–25
- Wavenumber:
 aqueous spectral calculations, 328–330
 orientational time correlation function:
 future research issues, 313
 thermotropic liquid crystals, Mode
 coupling theory (MCT), 270–273
- Wavepacket dynamics, local control theory:
 electronic excitation, 62–65
 triple-well potentials, 59–60
- Wideband loss frequency dependence, low-
 temperature spectrum, ice molecules,
 416–417
- Wideband water spectrum, room temperature
 conditions:
 mixed model (water, LDL, HDL) concepts,
 354–355
 ordinary and heavy water, 355–367
 research background, 353–354
 static permittivity, isotropic independence,
 355–357
- Wigner plot, mid-infrared pulse-shaping
 technology, ground-state vibrational
 dynamics, coherent control,
 9–16
- Xenon ions, double photoionization:
 data presentation, 117–120
 deep inner-shell thresholds, 121–124
 inner-shell DPI, 126–128
- Young-Dupré equation, interface between two
 immiscible electrolyte solutions,
 interfacial deposition, 207–208
- Young's modulus, ultrathin solid-fluid-solid
 films, 238–241
 evolutionary models, 234–237
- Yukawa interactions, interface between two
 immiscible electrolyte solutions, 159

