

Subject Index

- Acceptance sampling, 3
- Advantages of sampling, 1-2
- Aligned systematic sample, 228
- Analytical surveys, 4. *See also* Domains of study
- Area sample, 89
- Attributes, *see* Proportions
- Autocorrelated populations, 219

- Balanced repeated replications, 320
- Best linear unbiased estimate, 158
- Bias:
 - definition, 13
 - deliberate use of biased procedures, 12, 15
 - due to errors in stratum weights, 117
 - due to nonresponse, 359-361
 - of interviewer, 378
 - of ratio estimator, 160
 - of regression estimator, 198
 - in small areal units, 233
- Binomial distribution, 59
 - confidence limits, 59
 - erroneous use in cluster sampling, 64-68
 - references to tables, 55
 - for sample proportions, 55
- Boundaries of strata, rule for determining, 127-130

- Call-backs, 365
 - effects on response ratio, 365
 - mathematical model, 367
 - optimum policy, 369-370
 - relative cost, 366-367
- Cauchy-Schwarz inequality, 97
- Census, use of sampling, 2
- Centrally-located systematic samples, 205
- Change, estimates of, 345, 352-353
- Cluster sampling (single-stage):
 - clusters of equal sizes, 233
 - compared with SRS of elements, 242
 - estimation of proportions, 64-65, 246
 - optimum size and shape of cluster, 234-240, 246
 - variance as function of size of cluster, 243
 - variance of mean per element, 240
 - clusters of unequal sizes, 249
 - comparison of $\hat{Y}_u$, $\hat{Y}_R$, $\hat{Y}_{pps}$, 255-258
 - equal-probability selection, unbiased estimator $\hat{Y}_u$, 249
 - ratio estimator $\hat{Y}_R$, 250
- Horvitz-Thompson estimator, 259-261
 - optimum selection probabilities, 255
 - ratio estimators, 271
 - selection:
 - methods related to systematic sampling, 265
 - without replacement (WOR), 258-259
 - selection WOR, Brewer's method, 261
 - comparison of methods, 270
 - Murthy's method, 263
 - Rao, Hartley, Cochran method, 266
 - in stratified sampling, 270
 - unequal-probability selection with replacement (WR), $\hat{Y}_{ppz}$, 251-255
- Cluster sampling (with subsampling), *see* Two-stage sampling
- Coefficient of variation (cv), 54
- Collapsed strata method, 139, 227
- Combined ratio estimate, 155
 - compared with separate ratio estimate, 167
 - estimated variance, 167
 - in estimating domain means, 143
 - optimum allocation for, 172
 - in stratified two-stage sampling, 318
 - upper bound to relative bias, 156
 - variance, 166
- Combined regression estimate, 202
 - compared with separate regression estimate, 203
 - estimated variance, 203
 - variance, 202
- Composite estimate, in repeated sampling, 354
- Conditional distribution, of proportions, 51
- Confidence limits, 12
 - conditional, 60
 - definition for attributes, 57
 - effect of bias on, 14
 - effect of nonresponse on, 361-363

- for means in simple random sampling, 27
- for proportions or percentages, 57-60
- for ratio estimates, 156
- in stratified random sampling, 95
- validity of normal approximation, 39-44
- Consistency:
 - definition, 21
 - of mean of simple random sample, 21
 - of ratio estimate, 153
 - of regression estimate, 190
- Controlled selection, 124
- Correction for continuity, 58, 64
- "Correct value," 377
- Correlation coefficient:
 - in finite populations, 154
 - intracluster, 209
 - within a systematic sample, 209
- Correlogram, 219
- Cost function:
 - for number of strata, 134
 - in determining optimum probabilities of selecting primary units, 311-318
 - in determining optimum sampling fraction for nonrespondents, 371
 - in determining optimum size of cluster unit, 244
 - in determining optimum subsampling fraction, 280, 313-316
 - in determining sample size, 84
 - in double sampling:
 - for analytical comparisons, 335
 - for regression estimates, 341
 - with stratification, 331
 - in stratified random sampling, 96
 - in two-frame sampling, 145
- Covariance of sample means, 25
- Cumulative $\sqrt{f}$ rule, 129
- Degrees of freedom, effective number in stratified sampling, 96
- Descriptive surveys, 4
- Design effect (Deff), 21
- Domains of study (subpopulations), 34
 - comparisons between means, 39
 - comparisons between proportions and totals, 64
 - domains cutting across strata, 142-144
 - estimates of means and totals (continuous variate), 34-35
 - estimates of proportions and totals (0-1 variate), 63
 - sample size needed, 82-83
 - strata as domains, 140
- Double sampling, 327
 - ratio estimates, 343-344
- Double sampling with regression estimates, 335
 - comparison with simple random sampling, 341
 - estimated variance, 343
 - optimum sample sizes, 341
 - variance, 339-340
- Double sampling with stratification, 327
 - comparison with simple random sampling, 332
 - estimated variance, 333
 - estimation of proportions, 330
 - optimum sample sizes, 331
 - variance, 328
- E, average over all possible samples, 11
- E₁, E₂, averages over first and second stages, 275
- Elements, 31
- End corrections, 216
- Errors in surveys, types of, 359
- Errors of measurement, 377
 - effects of constant bias, 379
 - effects of correlation between errors, 383
 - effects of errors uncorrelated within sample, 380
 - mathematical model, 377-379
 - summary of effects, 384
 - techniques for studying, 384-392
 - use of interpenetrating subsamples, 389, 391
- Estimates of population variances:
 - effects of errors in S_h on precision of stratified sampling, 115
 - for determining sample size, 78
- Expansion factor, 21
- Eye estimates, 189
- Finite population correction (fpc), 24
 - rule for ignoring, 25
 - in stratified random sampling, 92
 - in two-stage sampling, 277
- Frame, 6
 - estimates when frame has units not in target population, 37-38
 - sampling from two frames, 144-146
- Geographic stratification, 102
 - construction of strata, 131
 - gains in precision, 102-103
- Hard core in nonresponse, 364
- Horvitz-Thompson estimator, 259-261
- Hypergeometric distribution, 55
 - as conditional distribution, 61

- confidence limits, 57
- reference to charts and tables, 57
- Index of inconsistency, 387
- Inflation factor, 21
- Interpenetrating subsamples, 388-392
- Interviewer bias, 378, 384
- mathematical models, 377-379
- Intraclass correlation, 209
- Inverse sample (Haldane's method), 77
- Item, definition, 20
- Jackknife method, 174
 - for estimating variance of a ratio, 178
 - for variance of nonlinear function, 321
- Keyfitz, short-cut method of variance estimation, 169-171
- Knight's move latin square, 229
- Labels, 44
- Lattice sampling, 228
- Linear regression estimator, *see* Regression estimator
- Listing of primary units, effect of cost on optimum selection probability, 316
- Loss function, 83, 121
- Mail surveys, 145, 360, 370
- Matching in repeated sampling of same population, 364-365
- Mean of sample, optimum properties, 44-45
- Mean square error, definition, 15
 - justification for use of, 15
 - relation to variance and bias, 15
- Measure of homogeneity, 248
- Model-unbiased, 158
- Multivariate ratio estimate, 185
- Neyman allocation, 99
 - best stratum boundaries for, 127
- Noncoverage, 364
- Nonnormality:
 - effect of stratification on, 44
 - effect on confidence limits, 41
 - frequently encountered in sampling practice, 40
- Nonresponse, 359
 - bias produced by, 359-361
 - effect of call-backs, 365-370
 - effect on confidence limits, 361-363
 - effect on variance in stratified sampling, 144
 - optimum sampling fraction among nonrespondents, 370-374
 - Politz and Simmons' method, 374-377
 - reasons for, 364
- Normal distribution, 8
 - as approximation to hypergeometric, 57
 - as limiting distribution of sample means, 39
 - use in surveys, 11
 - validity with means from continuous data, 39-44
- Notation:
 - for errors of measurement, 377-379
 - for proportions, 50, 63
 - for ratio estimates, 151
 - for simple random sampling, 20
 - for stratified sampling, 90
 - for two-stage sampling, 276
 - for variances of estimates, 27
- Optimum allocation in stratified sampling:
 - comparison with proportional allocation, 99, 109
 - comparison with simple random sampling, 99, 109
 - with double sampling, 330
 - effect of deviations from the optimum, 115
 - effect of errors:
 - in S_h , 115-117
 - in stratum sizes, 117
 - estimation from previous data, 102, 131
 - for fixed sample size, 98-99
 - for fixed total cost, 96-98
 - with more than one item, 119-123
 - with ratio estimates, 172
 - requiring more than 100% sampling, 104
 - in sampling for proportions, 108
- Optimum allocation in stratified two-stage sampling, 280
- Optimum per cent matched, in repeated sampling, 347, 350
- Optimum size of subsample:
 - primary units of equal size, 280
 - primary units of unequal size, 313-316
- Percentages, estimation of, *see* Proportions, estimation of
- Periodic variation, effect on systematic sampling, 217
- Pilot survey:
 - use in estimating optimum sampling and subsampling fractions, 283-285
 - use in estimating sample size, 78
- Politz and Simmons' method for handling nonresponse, 374-377

- Population, 5
 sampled, 5
 target, 5
 Poststratification, 134
 compared with proportional allocation, 134
 Precision contrasted with accuracy, 16
 relative, 99
 Pretest, 7
 Primary sampling units (primary units), 274
 Probability proportional to size, pps, (one-stage sampling), 250
 compared with equal probabilities, 255-258
 method of drawing sample, 250
 selection without replacement, 258
 Pps (two-stage sampling), 295, 308
 compared with equal probabilities, 299, 310
 selection without replacement, 308
 Probability sampling, definition and properties, 9
 alternatives to, 10
 Product estimator, 186
 Proportional allocation in stratified sampling, 91
 advice on use, 103, 110
 comparison with optimum allocation, 99, 103, 109
 comparison with poststratification, 134
 comparison with simple random sampling, 99, 109
 in estimating proportions, 109
 sample self-weighting, 91
 variance formulas, 92-95
 Proportions, estimation of, 50
 in cluster sampling, 64-68, 246
 in double sampling, 330
 effect of nonresponse, 361
 effect of population P on precision, 53
 with more than two classes, 60, 61
 in simple random sampling, 50-68
 size of sample for, 74
 in stratified random sampling, 107-111
 in two-stage sampling (units of equal size), 279
 in two-stage sampling (units of unequal size), 311
 Purposive selection, 10

 Quadratic confidence limits for ratio estimate, 156
 Qualitative characteristics, 50. *See also*
 Proportions, estimation of
 Quota sampling, 135

 Raising factor, 21
 Random member of household, method of
 selecting, 365

 Random numbers, 19
 Randomized response method, 392
 other variants, 395
 unrelated second question, 393
 Warner's original method, 392
 Random sampling, *see* Simple random sampling
 Rare items, Haldane's method, 77
 Ratio estimator, 30, 150
 accuracy of approximate variance, 162
 adjustments to decrease bias, 174
 bias, 160-162
 in cluster sampling, 31, 66
 compared with mean per unit, 157
 compared with regression estimate, 195
 compared with stratified sampling, 169
 conditions under which unbiased, 158
 confidence limits, 156
 consistency, 153
 in double sampling, 343
 estimated variance, 32, 155, 178
 Hartley-Ross estimate, 174
 jackknife estimate of variance, 178
 Lahiri's method, 175
 multivariate, 185
 optimum allocation in stratified sampling, 172
 optimum: conditions for, 160
 as special case of regression estimate, 190
 standard error for comparison of two ratios, 180
 in stratified random sampling, 164
 in stratified two-stage sampling, 317
 unbiased ratio-type estimates, 174
 upper bound to relative bias, 162
 variance in large samples, 30, 153
 see also Combined ratio estimate, Separate ratio
 estimate
 Ratio of two ratio estimators, 183
 Record checks, 385
 Regression coefficient in finite populations, 191
 Regression estimator, 189
 accuracy of large-sample variance, 197
 bias, 198
 compared with mean per unit, 195
 compared with ratio estimate, 195
 conditions under which unbiased, 199
 in double sampling, 338
 effect of error in slope, 192
 estimated variance, 195
 with preassigned slope, 190
 in repeated sampling of same population
 346-355
 in stratified random sampling, 201, 202
 uses, 189

- variance in large samples, 194
- see also* Combined regression estimate; Separate regression estimate
- Reinterviews in study of errors of measurement, 385, 386
- Relative precision, 99
 - method of calculating, 103
- Repeated measurements, 386, 391
- Repeated sampling of population, 344
 - composite estimate, 354
 - current estimates, 346-355
 - estimates of change, 345, 352
 - optimum percent matched, 347, 350
 - rotation policies, 354
 - use of nonrespondents from previous surveys, 377
- Response deviation, 379
- Response variance, 379
 - correlated component, 383
 - sample, 383
 - total, 383
- Sampled population, 5
- Sampling fraction, 21
 - first-stage, 277
 - second-stage, 277
- Sampling unit, definition, 6
- Sampling with replacement (WR), 18, 29
- Sampling without replacement (WOR), 18
- Satterthwaite's approximation to number of df, 96
- Self-weighting estimate, 91
 - in two-stage sampling, 303, 304, 307, 317
- Sensitive questions, 392
 - randomized response method, 392-395
- Separate ratio estimate, 164
 - compared with combined ratio estimate, 167
 - estimated variance, 167
 - liability to bias, 165
 - optimum allocation, 172
 - variance, 164
- Separate regression estimate, 201
 - compared with combined regression estimate, 203
 - estimated variance, 202
 - liability to bias, 202
 - variance, 202
- Short-cuts, in computation of variance of ratio estimate, 173
- Simple expansion, 169
- Simple random sampling, 18
 - for classification into more than two classes, 60
 - confidence limits,
 - for sample mean, 27
 - for sample proportion, 57
 - distribution of sample proportion, 55-57
 - estimated variance, 202
 - of sample mean, 26
 - of sample proportion, 52
 - method of drawing, 19
 - optimum linear estimator of population mean, 44
 - sample size:
 - needed for means, 77
 - sample size needed for proportions, 75
- Simple response variance, 383
- Size of sample for specified limits of error:
 - analysis of problem, 73
 - for comparisons between domains, 83
 - with continuous data, 77
 - Cox's method of two-step sampling, 78-80
 - for means over domains, 82
 - by minimizing cost plus loss due to errors, 83
 - with more than one item, 81
 - for normal approximation to confidence limits
 - for continuous data, 41-44
 - for normal approximation to confidence limits of proportions, 58
 - with proportions, 75
 - with rare items, 76
 - in stratified random sampling, 105, 110
- Skewness:
 - coefficient of, 42, 197
 - effect of stratification on, 44
 - effect on confidence limits, 41
- Square grid sample, 227
- Standard deviation in finite population, 23
- Standard error:
 - of difference between domain means, 39
 - of difference between two ratios, 180
 - of domain mean, 33-34
 - in stratified sampling, 145-147
 - of domain total, 35
 - in stratified sampling, 143
 - of estimated population total from simple random sample, 24
 - of mean:
 - in cluster sampling, 240
 - of simple random sample, 24, 25-27
 - in stratified sampling, 91-98
 - of systematic sample, 207-212
 - in three-stage sampling, 286-287
 - of proportion:
 - in cluster sampling, 64-68
 - in simple random sampling, 51-55

- in stratified sampling, 107-109
 - in two-stage sampling, 279
 - over a domain, 62-63
- of ratio estimate, 31-32, 153-156
 - in stratified sampling, 164-167
- of ratio in two-stage sampling, 311
- of regression estimate, 190, 194, 195
- of regression in stratified sampling, 200-203
- of sample standard deviation, 43
- of total in population possessing some attribute, 52
- Standard error (approximate) of nonlinear estimators, 318
 - balanced repeated replications, 320
 - comparison of methods, 322-324
 - jackknife method, 321
 - Taylor series method, 319
- Steps in a sample survey, 4
- Strata, 89
 - construction, 127-131
 - optimum number, 132-134
- Stratification, 89
 - best variable for, 101
 - effect of number of strata on precision, 132
 - effect on normality of variate, 44
 - two-way, 124
- Stratified random sampling, 89
 - compared with simple random sampling, 99-101
 - compared with ratio estimate, 165
 - compared with systematic sampling, 209-223
 - construction of strata, 127-131
 - estimate p_{st} , 107-108
 - estimate $\bar{y}_{st}$, 91-92
 - estimate of gain in precision, 136
 - estimated variance of $\bar{y}_{st}$, 95
 - estimated variance of p_{st} , 108
 - estimates for domains of study, 142-144
 - Neyman allocation, 98-99
 - with one unit per stratum, 138
 - optimum allocation for fixed cost, 96-98
 - with ratio estimates, 164
 - with regression estimates, 200
 - size of sample, 105, 110
 - type of population giving large gains, 101
- Subpopulations, *see* Domains of study
- Subunits (elements), 233
- Superpopulation, 158
- Systematic sampling, 205
 - in autocorrelated populations, 219
 - compared with simple random sampling, 208-223
 - compared with stratified sampling, 209-223
 - effect of periodic variation, 217
 - end corrections, 216
 - estimation of variance, 223-226
 - method of drawing, 206
 - in natural populations, 221
 - in populations:
 - in "random" order, 212
 - with linear trend, 214-217
 - in single-stage cluster sampling, 265
 - stratified systematic sampling, 226
 - in two dimensions, 227
 - in two-stage sampling, 279
 - recommendations about use, 229
 - relation to cluster sampling, 207
 - variance of mean, 207-212
- Target population, 5
- Taylor series method, 319
- Theory, in sample surveys, 8
- Three-stage sampling, 285
 - optimum sampling and subsampling fractions, 288
 - variance of mean per third-stage unit, 286-287
- Total response variance, 383
- Travel costs, formula, 96
- Two-phase sampling, *see* Double sampling
- Two-stage sampling (units of equal size):
 - advantage, 274
 - optimum sampling and subsampling fractions, 280
 - stratified sampling of the primary units, 288-289
 - table for selecting optimum size of subsample, 283
 - use of pilot survey, 283
 - variances:
 - of estimated mean, 276
 - of estimated proportions, 279
- Two-stage sampling (units of unequal size), 292
 - comparison of methods, 310
 - methods with one unit per stratum, 293-299
 - optimum sampling and subsampling fractions, 313-316
 - ratio estimators, 311
 - units chosen:
 - with equal probabilities, 293-295, 303-305
 - with unequal probabilities WOR, 308-310
 - with unequal probabilities WR, 306-308
- Unaligned systematic sample, 228
- Unbiased estimate, definition, 11

- Unit (sampling unit), definition, 6
- United States Census, use of sampling, 2-3
- Uses of sample surveys, 2-4
- by business and industry, 3
 - in decennial censuses, 2-3
 - in market research, 3
 - by members of United Nations, 2
 - in opinion polls, 4
- Variance, definition of S^2 and σ^2 , 22, 25
- Variance function, 243
- Variance of population, advance estimates, 78-81
- Variance of sample estimates, *see* Standard error

