

SUBJECT INDEX

- Aftereffect solution, single-domain
 - ferromagnetic particle relaxation, Brown's model, 232–234
- Alignment term, single-domain
 - ferromagnetic particle relaxation, Brown's model, 226–227, 230–234
- Anisotropic axes:
 - magnetoanisotropic nanoparticles, 10–13
 - Hamiltonian formulas, 67–69
 - linear susceptibility, random distribution, 63, 66–67
 - magnetization, series expansions
 - along, 53–57
 - partition function, series expansion, 29–32
 - powers expanded by free energy, 38–39
 - thermal equilibrium quantities, longitudinal-field energy barriers, 46–49
 - nonlinear susceptibility:
 - random distribution, 75–76
 - temperature dependence, 83–85
 - overbarrier rotations, individual spins, 134–135
 - powers expanded by free energy:
 - field expansion of free energy, 37–38
 - free energy series random distribution, 35–36
 - partition function, 15–16
 - reduced nonlinear susceptibility, 76–77
 - spin ensemble dynamics, 142–144
 - temperature dependence, 136–137
 - quantum electrodynamics (QED),
 - resonance energy transfer, fluorescence depolarization, 380–383
 - single-domain ferromagnetic particles, nonaxially symmetric particles, 246–256
- Anisotropic property, defined, 330
- Anisotropy field, dynamical properties, magnetoanisotropic nanoparticles, deterministic dynamics, Landau–Lifshitz–Gilbert equation, 101–103
- Approximation formulas, nonaxial symmetry:
 - asymptotic formulas:
 - vs. experimental observations, 277–279
 - as function of field angle, 275–277
 - Brown's high-energy approximation, 256–262
 - analytic solutions, 263–265
 - IHD vs. LD formulas, Fokker-Planck equation, 268–275
 - Kramers low-damping and transition-state results, 265–267
 - validity as function of damping parameter, 267–268
- Arrhenius law, magnetoanisotropic nanoparticles:
 - Shliomis-Stepanov model, longitudinal-field energy barriers, 91
 - spin ensemble dynamics, 144
- Asymptotic expansion:
 - approximate formulas, 190–193
 - Fokker-Planck matrix calculations, eigenvalue comparisons, 285–297
 - large negative argument, 188–189
 - large positive argument, 187–188
 - magnetoanisotropic nanoparticles:
 - free energy, strong anisotropy, 40
 - longitudinal response, 148–150
 - magnetization, expansion for strong anisotropy, 57–59

- Asymptotic expansion: (*continued*)
 partition function, strong anisotropy, 32–35
 thermal equilibrium quantities, longitudinal-field regimes, 48–49
 nonaxial symmetry:
 experimental observations, 277–279
 field angles and, 275–277
 future research applications, 308
 one-dimensional Ising spin-lattice model, Ising ring formation, 347–353
- Autocorrelation function, single-domain ferromagnetic particle relaxation, Brown's model, 233–234
- Average squared magnetization (ASM), one-dimensional Ising spin-lattice model:
 Ising chain formation, 353–354
 Ising ring formation, 346–353
 research background, 337–340
 statistical thermodynamics, 342–345
- Averaging equations:
 anisotropy axis, magnetoanisotropic nanoparticles:
 field expansion of free energy, 37–38
 free energy random distribution, 36
 dynamical properties, magnetoanisotropic nanoparticles, magnetic moment averages, 120–122
- Axial symmetry:
 defined, 330
 dynamical properties, magnetoanisotropic nanoparticles, deterministic Landau–Lifshitz equation, 103–104
 Fokker-Planck equation, Sturm-Liouville expression, 118–120
 single-domain ferromagnetic particles:
 Brown's low-energy-barrier approximation, 236
 domain structure, 220
 solutions for, 234–236
- Azimuthal angles:
 anisotropy axis, magnetoanisotropic nanoparticles, 36
 magnetoanisotropic nanoparticles, deterministic dynamics, 105–106
- Backreaction on system:
 Brown-Kubo-Hashitsume model, linear couplings, 162–163
 equations of motion, nonlinear couplings, 203–207
- Barrier height parameter:
 Fokker-Planck matrix calculations:
 correlation times, numerical calculation of, 297–306
 lowest eigenvalues, 284–285
 nonaxial symmetry, vs.
 intermediate-to-high damping (IHD) formula, 268–275
 single-domain ferromagnetic particles, nonaxially symmetric particles, 246–256
- Bath polaritons:
 quantum electrodynamics (QED), resonance energy transfer, 363–364
- Bath polaritons, quantum electrodynamics (QED), resonance energy transfer:
 background, 360–361
 molecular dynamics in dielectric media, 389–400
 nonrate regimes, 397–400
- Beer law, quantum electrodynamics (QED), resonance energy transfer, 360–361
- dielectric media, vibrational structure, 378–379
- Bessel function, magnetoanisotropic nanoparticles, partition function, 16
- Bilinear coupling:
 Brown-Kubo-Hashitsume model:
 phenomenological equations, 156–157
 spin-environment Hamiltonians, 160

- Binomial coefficients,
 - magnetoanisotropic nanoparticles, partition function, field series expansion, 23–24
- Bloch equations, dynamical properties, magnetoanisotropic nanoparticles, averaging equations, magnetic moments, 121–122
- Blocked particles, in-phase linear dynamical susceptibility, magnetic nanoparticle dynamics, 92–94
- Blocking temperature:
 - magnetoanisotropic nanoparticles, 7–8
 - low-frequency Shliomis-Stepanov equation, 94–96
 - Shliomis-Stepanov model, longitudinal-field energy barriers, 89–91
 - spin ensemble, longitudinal dynamical response, 140–141
 - nonaxial symmetry, future research issues, 308
 - single-domain ferromagnetic particle relaxation, Néel relaxation model, 224
- Boltzmann distribution:
 - Brown-Kubo-Hashitsume model, phenomenological equations, 155–156
 - Fokker-Planck equation:
 - Markovian regime, 178–179
 - stationary solution, 113–115
 - magnetoanisotropic nanoparticles, partition function and free energy, 14–15
 - statistical properties, fluctuating magnetic field, decoupling conditions, 167–168
- Boltzmann's constant, single-domain ferromagnetic particles, domain structure, 217–220
- Born-Oppenheimer approximation, quantum electrodynamics (QED), resonance energy transfer, dielectric media, vibrational structure, 376–379
- Bose operator, resonance dipole-dipole interaction (RDDI), dielectric media, 372–376
- Box-Muller algorithm, dynamical properties, magnetoanisotropic nanoparticles, Gaussian stochastic process, 132–133
- Brillouin zone, quantum electrodynamics (QED), resonance energy transfer, displacement field operator, 402–403
- Brown-Aharoni formula, Fokker-Planck matrix calculations, asymptotic estimate vs. numerical calculations, 292–297
- Brownian particle dynamics, Brown-Kubo-Hashitsume model, linear couplings, 163
- Brown-Kubo-Hashitsume model:
 - dynamical properties, magnetoanisotropic nanoparticles, 107–128
 - Fokker-Planck equations, 110–120
 - Langevin equations, 108–110
 - magnetic moment averages, equations for, 120–122
 - relaxation times, 123–128
 - stochastic dynamics, 100
 - magnetoanisotropic nanoparticles, 6
 - Markovian regime, 179–180
 - stochastic dynamical equations, 179–180
 - free dynamics and canonical variables, 157–159
 - Garanin-Ishchenko-Panina model, 180–181
 - linear couplings, 159–169
 - fluctuating source properties, 167–169
 - Hamiltonian notations, 160–163
 - spin-dynamics case, 163–167
 - spin-environment Hamiltonian, 159–160
 - linear-plus-quadratic couplings, 169–174

- Brown-Kubo-Hashitsume model:**
 (continued)
 fluctuating source statistical properties, 173–174
 Hamiltonian notations, 170–171
 spin-dynamics case, 171–173
 spin-environment Hamiltonian, 169–170
 Markovian regime, 174–179
 damping coefficients, 176
 Fokker-Planck equations, 177–179
 Langevin equations, 174–176
 phenomenological equations, 153–156
 spin environment interactions, 156–157
 summary, 181–182
- Brown's high-energy approximation:**
 Fokker-Planck matrix calculations, asymptotic estimate vs. numerical calculations, 294–297
 nonaxial symmetry, 256–262
 analytic solutions, 263–265
 damping parameter and validity of, 267–268
- Brown's model, single-domain**
 ferromagnetic particle relaxation, 224–234
 high-energy-barrier approximation, 236–244
 low-energy-barrier approximation, 236
- B/T* superposition of M_B , nanometric magnetic particle partition functions, magnetization, 51**
- Canonical variables,**
 Brown-Kubo-Hashitsume model, free dynamics, 157–159
- Casimir-Polder formula, quantum electrodynamics (QED), resonance energy transfer, 362**
- Causality, quantum electrodynamics (QED), resonance energy transfer, molecular dynamics in dielectric media, 390–400**
- Chain rules:**
 Brown-Kubo-Hashitsume model, free dynamics, canonical variables, 158–159
 one-dimensional Ising spin-lattice model, Ising chain formation, 353–354
- Collective emission, quantum electrodynamics (QED), resonance energy transfer, 359–361**
- Condon factor, quantum electrodynamics (QED), resonance energy transfer: dielectric media, vibrational structure, 376–379**
 excited molecule decay, 387–389
- Continued fractions techniques, Fokker-Planck equation, dynamical properties, magnetoanisotropic nanoparticles, 99**
- Continuity equation, Fokker-Planck equation derivation, 111–113**
- Convergence, dynamical properties, magnetoanisotropic nanoparticles, order of, 129–130**
- Correlation time (T_c), Fokker-Planck matrix calculations:**
 numerical computation of, 297–306
 results tables, 323–329
- Coulomb interaction:**
 quantum electrodynamics (QED), resonance energy transfer, 358–361
 radiative (polariton) bath, 364
- Coupled initial conditions, statistical properties, fluctuating magnetic field:**
 linear couplings, 168–169
 linear-plus-quadratic couplings, 174
- CPU time requirements, Fokker-Planck matrix calculations:**
 correlation times, numerical calculation of, 306
 future research applications, 306–307
 lowest eigenvalues, 284–285

- Cubic anisotropy, Fokker-Planck matrix calculations and, 306–307
- Curie point:
 - defined, 330
 - in-phase linear
 - dynamical susceptibility, magnetic nanoparticle dynamics, 92–94
 - one-dimensional Ising spin-lattice model, 338–340
 - Ising ring formation, 347–353
 - statistical thermodynamics, 342–345
- Curie susceptibility law,
 - magnetoanisotropic nanoparticles:
 - linear susceptibility:
 - axial anisotropy, random distribution, 63, 67
 - isotropic regime, 61
 - nonlinear susceptibility, 76–77
 - temperature dependence, 83–85
- Current density, single-domain
 - ferromagnetic particle relaxation:
 - Brown's model, 228–229
 - high-energy-barrier approximation,
 - axial symmetry, 238–244
- Damping coefficient:
 - Langevin equation, 174–176
 - magnetoanisotropic nanoparticles,
 - transverse response, 150–153
 - Markovian regime, 176
- Dawson integral, Kummer functions and, 185–186
- Debye relaxation:
 - in-phase linear
 - dynamical susceptibility, magnetic nanoparticle dynamics, 91–92
 - low-frequency Shliomis-Stepanov model, magnetic nanoparticle dynamics, 87–88
 - magnetoanisotropic nanoparticles,
 - longitudinal response, 147–150
 - $\pi/2$ law, low-frequency Shliomis-Stepanov equation, 93–94
 - single-domain ferromagnetic particles, 221
 - Néel relaxation model, 224
- Decay factor, quantum electrodynamics (QED), resonance energy transfer, 360–361
 - dielectric media, vibrational structure, 378–379
 - spontaneous emission, excited molecule decay, 385–389
- Decoupling conditions, statistical properties, fluctuating magnetic field:
 - linear couplings, 167–169
 - linear-plus-quadratic couplings, 173–174
- Degenerate properties, defined, 330
- Degree of polarization,
 - magnetoanisotropic nanoparticles,
 - partition function, field series expansion, 24–25
- Deterministic dynamics,
 - magnetoanisotropic nanoparticles,
 - classical spins, Landau–Lifshitz (–Gilbert) equation, 101–103
- Deterministic Landau–Lifshitz equation,
 - dynamical properties, magnetoanisotropic nanoparticles, 100
 - axially symmetric Hamiltonians, 103–104
- Dexter exchange interaction, quantum electrodynamics (QED),
 - resonance energy transfer, 359–361
- Dielectric media:
 - quantum electrodynamics (QED),
 - resonance energy transfer:
 - background, 359–361
 - fluorescence depolarization range dependence, 379–383
 - molecular dynamics, 389–400
 - energy shifts, 392–396
 - nonrate regime, 397–400
 - rate description, 396–397
 - time evolution of quantum systems, 390–392
 - spontaneous emission, 384–389
 - transfer species vibrational structure, 376–379

- Dielectric media: (*continued*)
 resonance dipole-dipole interaction (RDDI):
 retarded RDDI tensor analysis, 403–406
 tensor analysis, 372–376
- Dimensionless anisotropy, magnetoanisotropic nanoparticles, thermal equilibrium quantities, 42–43
- Dipole moments:
 defined, 330
 magnetoanisotropic nanoparticles, linear susceptibility, random distribution, 66–67
- Dirac delta functions:
 magnetoanisotropic nanoparticles:
 Gaussian stochastic process, 108–110
 partition functions in free energy:
 Ising regime, 19
 plane-rotator regime, 19–20
 Shliomis-Stepanov model, longitudinal-field energy barriers, 89–91
 Markovian regime, phenomenological equations, 175–176
- Direction cosines, nonaxial symmetry, orthogonal transformations, 309–312
- Direct solution techniques, Fokker-Planck equation, dynamical properties, magnetoanisotropic nanoparticles, 99
- Discrete orientation:
 Brown's high-energy approximation, nonaxial symmetry, 263–265
 defined, 330
 magnetoanisotropic nanoparticles, linear susceptibility, heuristic treatment, 88
 single-domain ferromagnetic particle relaxation, 223–224
 high-energy-barrier approximation, axial symmetry, 238–244
 Néel relaxation model, 223
- Dispersion relationships, resonance dipole-dipole interaction (RDDI):
 dielectric media, 372–376
 photonic bandgap crystals, 369–372
- Displacement field operator, quantum electrodynamics (QED), resonance energy transfer, 401–403
- Distribution of magnetization orientation, single-domain ferromagnetic particles, domain structure, 219–220
- Domains, single-domain ferromagnetic particles:
 geometric structure, 216–220
 superparamagnetic effect, 212–213
- "Dressed" photons, quantum electrodynamics (QED), resonance energy transfer, transfer rates, 366–367
- Dynamical properties, magnetoanisotropic nanoparticles, 99–153
 heuristic approach, 85–99
 frozen magnetic fluid experiment, 96–99
 in-phase linear dynamical susceptibility, 91–92
 linear dynamical response, 86–89
 low-frequency Shliomis-Stepanov model, 88–96
 stochastic theory, 6, 99–153, 182–183
 Brown-Kubo-Hashitsume model, 107–128
 deterministic dynamics, classical spins, 101–107
 dynamics, 99–153
 individual spin trajectories, 133–138
 numerical method, 128–133
 spin ensemble dynamics, 138–153
- Easy-plane anisotropy:
 dynamical properties,
 magnetoanisotropic nanoparticles, numerical calculations, 128
 magnetoanisotropic nanoparticles:
 deterministic dynamics, 105–106

- linear susceptibility formulas, 70–71
- series expansion of magnetization, 56–57
- Effective eigenvalue, single-domain ferromagnetic particle relaxation, Brown's model, 233–234
- Effective-mass approximations, resonance dipole-dipole interaction (RDDI), photonic bandgap crystals, 371–372
- Eigenvalue calculations:
 - Brown's high-energy approximation, nonaxial symmetry, 263–265
 - Fokker-Planck equation:
 - dynamical properties, magnetoanisotropic nanoparticles, 99
 - longitudinal relaxation times, 123–125
 - matrix computation:
 - asymptotic estimates, comparison, 285–297
 - correlation time calculations, 297–306
 - lowest eigenvalue, 279–306
 - Sturm-Liouville problem, 119–120
 - integral relaxation time, 194
 - magnetoanisotropic nanoparticles, transverse relaxation time, 125–128
 - nonaxial symmetry, vs. intermediate-to-high damping (IHD) formula, 268–275
 - single-domain ferromagnetic particles:
 - Brown's model, 234
 - low-energy-barrier approximation, 236
 - nonaxially symmetric particles, 248
 - results tables, 316–322
- EISPACK library, Fokker-Planck matrix calculations, 279
- Energy transfer (ET). *See also* resonance energy transfer
- Entropy, magnetoanisotropic nanoparticles, thermal equilibrium quantities, 41–43
- Equations of motion, nonlinear couplings, 200–207
- Equilibrium probability distribution, magnetoanisotropic nanoparticles, partition function and free energy, 14–15
- Equilibrium properties, magnetoanisotropic nanoparticles, 6–8
 - Hamiltonians, 8–13
 - linear susceptibility, 61–73
 - longitudinal-field case, energy barriers, 11–13
 - magnetization, 49–61
 - nonlinear susceptibility, 73–85
 - thermal (caloric) quantities, 41–49
- Equilibrium transverse susceptibility, magnetoanisotropic nanoparticles, spin ensemble dynamics, 143–144
- Error functions, Kummer functions and, 184–186
- Euler-type algorithms, dynamical properties, magnetoanisotropic nanoparticles, numerical calculations, 130–132
- Excitation frequencies, quantum electrodynamics (QED), resonance energy transfer, molecular dynamics in dielectric media, 396–399
- Excited state molecules, quantum electrodynamics (QED), resonance energy transfer, spontaneous emission, excited molecule decay, 385–389
- Far-zone energy transfer, spontaneous emission, quantum electrodynamics (QED), 383–389
- excited molecule decay, 385–389
- research background, 383–385
- Fermi golden rule, quantum electrodynamics (QED), resonance energy transfer:
 - spontaneous emission, 384–389
 - transfer rates, 364–367

- Ferrofluids, nanometric particle magnetization, 49–50
- Ferromagnetic behavior:
- one-dimensional Ising spin-lattice model, 355–356
 - single-domain ferromagnetic particles, relaxation mechanisms, 221–222
- Field-cooled magnetization,
- Fokker-Planck matrix calculations, correlation times, numerical calculation of, 306
- Field series expansion:
- magnetoanisotropic nanoparticles:
 - coefficients, alternative expressions, 24–25
 - free energy expansion, 37–38
 - Ising regime, 27–28
 - isotropic regimes, 27
 - longitudinal-field regime, 28–29
 - magnetization, low-field expansion, 59–61
 - partition function, 23–29
 - plane-rotator regime, 28
 - resonance dipole-dipole interaction (RDDI), dielectric media, 373–376
- Finite-size spin systems, one-dimensional
- Ising spin-lattice model:
 - background, 337–340
 - chain structure, 353–354
 - ring structure, 345–353
 - statistical thermodynamics of, 340–345
 - temperature effects, 354–356
- Fluctuating magnetic field:
- dynamical response:
 - linear couplings, 164–167
 - linear-plus-quadratic couplings, 173–174
 - statistical properties, 167–169
- Fluctuation-dissipation reactions:
- magnetoanisotropic nanoparticles.
 - Brown-Kubo-Hashitsume model, 107
 - Markovian regime, phenomenological equations, 175–176
 - statistical properties, fluctuating magnetic field, decoupling conditions, 168
- Fluorescence depolarization, quantum electrodynamics (QED), resonance energy transfer, range dependence, 379–383
- Fokker-Planck equation:
- Brown-Kubo-Hashitsume model, phenomenological properties, 154–156
- Garanin-Ishchenko-Panina model, 181
- integral relaxation time formula:
- low-frequency dynamical susceptibility, 194
 - perturbational solution in low sinusoidal field, 195–197
- magnetoanisotropic nanoparticles,
- dynamic properties:
 - derivation of, 110–113
 - Itô calculus, 109–110, 115–117
 - longitudinal relaxation times, 123–125
 - longitudinal response, 148–150
 - spherical coordinates, 117–118
 - stationary solution, 113–115
 - stochastic approach using, 99–100
 - Sturm-Liouville expression, axial symmetry, 118–120
- Markovian regime, 177–179
- nonaxial symmetry:
- asymptotic expansion, field angles and, 275–277
 - Brown's high-energy approximation, 258–262
 - IHD vs. LD formulas, 268–275
 - Kramers low-damping and transition-state results, 265–267
 - numerical computation of matrix and lowest eigenvalue, 279–306
 - asymptotic estimates, comparison, 285–297
 - correlation time calculations, 297–306
- single-domain ferromagnetic particles:
- axial symmetry solutions, 234–236

- Brown's model, 227–234
- future research issues, 307–308
- high-energy-barrier approximation,
 - axial symmetry, 242–244
- matrix formulation, 249–252
- nonaxially symmetric particles, 244–248
- response function and initial conditions, 252–256
- Förster theory of short-range energy transfer, quantum electrodynamics (QED), resonance energy transfer, 358–361
- dielectric media, vibrational structure, 378–379
- excited molecule decay, 385–389
- Fourier integrals:
 - quantum electrodynamics (QED), resonance energy transfer, 402–403
 - single-domain ferromagnetic particle relaxation, Fokker-Planck equation, matrix formulation, 249–252
- Free-diffusion, dynamical properties, magnetoanisotropic nanoparticles, averaging equations, magnetic moments, 122
- Free dynamics, Brown-Kubo-Hashitsume model, canonical variables, 157–159
- Free energy distribution, magnetoanisotropic nanoparticles, 13–22
 - definitions, 13–15
 - ising regime, 19
 - isotropic case, 17
 - limiting regimes, 16–22
 - longitudinal field case, 20–22
 - plane-rotor regime, 19–20
 - series expansions, 35–40
 - anisotropy axes, random distribution averages, 35–36
 - anisotropy parameter, expansion in powers of, 38–39
 - asymptotic expansion for strong anisotropy, 40
 - field expansions, 37–38
 - logarithmic expansion, 35
 - simple axially symmetric anisotropy potential, 15–16
 - zero-field case, 17–19
- Frozen magnetic fluids, magnetic nanoparticle dynamics, heuristic approach, 96–99
- Gamma (factorial) functions:
 - Kummer functions and, 184–186
- magnetoanisotropic nanoparticles:
 - anisotropy axis, 36
 - asymptotic expansion, partition function, 33–35
- magnetoanisotropic nanoparticles, partition function, field series expansion, 26–27
- Garanin-Ishchenko-Panina model:
 - integral relaxation time, 197–198
- Markovian regime, 180–181
- Gaussian properties:
 - quantum electrodynamics (QED), resonance energy transfer, fluorescence depolarization, 382–383
 - statistical properties, fluctuating magnetic field:
 - linear decoupling conditions, 167–168
 - linear-plus-quadratic decoupling conditions, 173–174
- Gaussian stochastic process:
 - dynamical properties, magnetoanisotropic nanoparticles, implementation, 132–133
- Fokker-Planck equation derivation, 110–113
- magnetoanisotropic nanoparticles, 108–110
- Gibbs canonical ensemble, single-domain ferromagnetic particles, domain structure, 219–220
- Gibbs free energy:
 - defined, 330

- Gibbs free energy: (*continued*)
 single-domain ferromagnetic particle relaxation:
 axial symmetry solutions, 234
 high-energy-barrier approximation, 236–244
 nonaxially symmetric particles, 244–256
- Gilbert equation.
See also Landau–Lifshitz–Gilbert equation
 Brown-Kubo-Hashitsume model,
 Markovian regime, 179–180
 dynamical properties,
 magnetoanisotropic nanoparticles,
 deterministic dynamics,
 Landau–Lifshitz–Gilbert
 equation, 102–103
 linear couplings, spin-dynamics
 case, 164–167
 magnetoanisotropic nanoparticles,
 Brown-Kubo-Hashitsume model,
 107
 Markovian regime, 174–179
 nonaxial symmetry, asymptotic
 expansion, 278–279
 single-domain ferromagnetic particle
 relaxation:
 Brown's model, 226–234
 nonaxially symmetric particles,
 246–256
- Gittleman-Abeles-Bozowski model,
 magnetoanisotropic nanoparticles:
 linear susceptibility, heuristic
 treatment, 88
 longitudinal response, 146–150
 transverse response, 151–153
- Green function:
 quantum electrodynamics (QED),
 resonance energy transfer,
 spontaneous emission, 384–385
 resonance dipole-dipole interaction
 (RDDI), dielectric media,
 372–376
- Gyromagnetic function:
 defined, 330
- Fokker-Planck matrix calculations,
 asymptotic estimate vs. numerical
 calculations, 286–297
- single-domain ferromagnetic particle
 relaxation:
 axial symmetry solutions, 235–244
 Brown's model, 225, 230–234
 nonaxially symmetric particles,
 245–256
- Gyrosopic function, defined, 330
- Hamiltonian equations:
 Brown-Kubo-Hashitsume model:
 linear coupling, 160–163
 linear-plus-quadratic coupling,
 170–171
 magnetoanisotropic nanoparticles,
 8–13
 dynamic properties, axially
 symmetric Hamiltonians,
 deterministic Landau–Lifshitz
 equation, 103–104
 quantum electrodynamics (QED),
 resonance energy transfer,
 362–363
 molecular dynamics in dielectric
 media, 392–396
 radiative (polariton) bath, 363–364
 resonance dipole-dipole interaction
 (RDDI), vacuum calculations,
 367–369
- Heaviside function, quantum
 electrodynamics (QED),
 resonance energy transfer,
 molecular dynamics in dielectric
 media, 390–392
- Hegerfeldt theorem, quantum
 electrodynamics (QED),
 resonance energy transfer,
 molecular dynamics in dielectric
 media, 396
- Heisenberg-type Hamiltonian,
 Brown-Kubo-Hashitsume model,
 phenomenological equations,
 156–157

- Heitler-Ma method, quantum electrodynamics (QED), resonance energy transfer, molecular dynamics in dielectric media, 390–392
- Hermitian matrix, Fokker Planck equation, Sturm-Liouville problem, 120
- Hessian matrix, nonaxial symmetry, Kramers low-damping and transition-state results, 266–267
- Heun scheme, dynamical properties, magnetoanisotropic nanoparticles: numerical calculations, 130–132 spin ensemble response, 138–153 stochastic trajectories, individual spins, 132–138
- Heuristics, magnetoanisotropic nanoparticle dynamics, 85–86 frozen magnetic fluid experiments, 96–99 linear dynamical response, 86–88 low-frequency Shliomis-Stepanov model, 88–96
- High-energy-barrier approximations: Fokker-Planck matrix calculations, asymptotic estimate vs. numerical calculations, 286–297 low-frequency Shliomis-Stepanov equation, 96 magnetoanisotropic nanoparticles, linear susceptibility, heuristic treatment, 88 nonaxial symmetry, Brown's high-energy approximation, 256–262 single-domain ferromagnetic particles, axial symmetry solutions, 236
- Hilbert space, single-domain ferromagnetic particle relaxation, Fokker-Planck equation, matrix formulation, 249–252
- Hopfield model, resonance dipole-dipole interaction (RDDI), dielectric media, 373–376
- Hysteresis, defined, 330
- Imaginary infinitesimals, resonance dipole-dipole interaction (RDDI), vacuum calculations, 369
- Initial conditions, single-domain ferromagnetic particle relaxation, Fokker-Planck equation, 252–256
- In-phase linear dynamical susceptibility, magnetoanisotropic nanoparticles, 91–92
- Integral relaxation time: Fokker-Planck equation, low sinusoidal field, 195–197 Garanin-Ishchenko-Panina formula, 197–198 isotropic regime, 198 zero-field case, 198 low-frequency dynamical susceptibility, 194 magnetoanisotropic nanoparticles: Brown-Kubo Hashitsume model, 124–125 longitudinal response, 146–150
- Intermediate-to-high damping (IHD) formula, nonaxial symmetry: Brown's high-energy approximation, 258–262 damping parameter and validity of, 267–268 future research issues, 308 Kramers low-damping and transition-state results, 265–267 vs. low-density formulas, 268–275
- Inter-potential-well rotations, magnetoanisotropic nanoparticles: longitudinal response, 148–150 spin ensemble dynamics, 141–142
- Intra-potential-well relaxation modes, magnetoanisotropic nanoparticles, dynamic properties, longitudinal response, 148–150
- Ising model: finite-size spin systems, one-dimensional model: background, 337–340 chain structure, 353–354 ring structure, 345–353

- Ising model: (*continued*)
 statistical thermodynamics of, 340–345
 temperature effects, 354–356
 magnetoanisotropic nanoparticles:
 asymptotic expansion, partition function, 35
 dynamics, heuristic analysis, 97–98
 linear susceptibility, 61
 linear susceptibility, heuristic treatment, 88
 logitudinal-field energy barriers, 21–22
 magnetization, 50
 expansion for strong anisotropy, 57–59
 nonlinear susceptibility, 74, 82–85
 partition functions in free energy, 19
 series expansion, partition functions, 27–28
- Isothermal magnetic susceptibility,
 one-dimensional Ising spin-lattice model, statistical thermodynamics, 342–345
- Isotropic regimes:
 integral relaxation time, 198
 magnetoanisotropic nanoparticles:
 deterministic dynamics, 105
 free energy partition functions, 17
 linear susceptibility, 61
 magnetization, 50
 nonlinear susceptibility, 74, 81–85
 series expansion, partition functions, 27
 thermal equilibrium quantities, 43–44
 single-domain ferromagnetic particles, domain structure, 218–220
- Isserlis theorem, single-domain ferromagnetic particle relaxation, Brown's model, 227–228
- Itô calculus, dynamical properties, magnetoanisotropic nanoparticles:
 Fokker-Planck equations, 115–117
 Langevin equations, 109–110
 numerical calculations, 130–132
- Kernel derivation, equations of motion, nonlinear couplings, 205–207
- Klein-Kramers equation:
 Brown's high-energy approximation, nonaxial symmetry, 259–262
 Fokker-Planck matrix calculations, asymptotic estimate vs. numerical calculations, 294–297
 nonaxial symmetry:
 asymptotic expansion, field angles and, 276–277
 asymptotic expansion vs. experimental observations, 277–279
 vs. intermediate-to-high damping (IHD) formula, 273–275
- Kramers-Krönig relations, $\pi/2$ law, low-frequency Shliomis-Stepanov equation, 93–94
- Kramers problem, nonaxial symmetry, asymptotic expansion, 277–279
- Kramers theory:
 Fokker-Planck matrix calculations, asymptotic estimate vs. numerical calculations, 294–297
 nonaxial symmetry:
 Brown's high-energy approximation, 259–262
 damping parameter and validity of, 267–268
 vs. intermediate-to-high damping (IHD) formula, 269–275
 low-damping and transition-state results, 265–267
 single-domain ferromagnetic particle relaxation, 244–256
- Kröneckner delta, magnetoanisotropic nanoparticles, Gaussian stochastic process, 108–110
- Kummer functions:
 definition, 184–186
 large negative argument, 188–189
 magnetoanisotropic nanoparticles:
 anisotropy axis, 36
 asymptotic expansion, partition function, 33–35

- linear susceptibility, Hamiltonian formulas, 69
- magnetization formulas, 53
- partition function, series expansion, 24–27
- power series expansions, 187
- Lamb shift, quantum electrodynamics (QED), resonance energy transfer, molecular dynamics in dielectric media, 392–396
- Landau–Lifshitz equation:
 - Brown-Kubo-Hashitsume model, Markovian regime, 179–180
 - Garanin-Ishchenko-Panina model, 181
 - linear couplings, spin-dynamics case, 165–167
 - linear-plus-quadratic couplings, spin-dynamics case, 172–173
 - Markovian regime, 174–179
 - single-domain ferromagnetic particle relaxation, Brown's model, 225–227
- Landau–Lifshitz (–Gilbert) equation. *See also* Deterministic Landau–Lifshitz equation; Gilbert equation; Stochastic Landau–Lifshitz equation
 - dynamical properties, magnetoanisotropic nanoparticles:
 - averaging equations, 121–122
 - deterministic dynamics, 101–103
 - dimensionless quantities, 128–129
 - Fokker-Planck equation derivation, 111–113
 - integration of, 132–133
 - Itô calculus, Fokker-Planck equation, 116–117
 - numerical calculations, 131–132
 - vs. heuristic models, 146–153
 - spin ensemble response, 138–153
 - stationary solution, Fokker-Planck equation derivation, 114–115
 - stochastic approach, 99–100
 - stochastic trajectories, individual spins, 133–138
- Langevin-dynamics approach, Fokker-Planck equation, dynamical properties, magnetoanisotropic nanoparticles, 99–100
- Langevin equation:
 - asymptotic expansion, 308
 - Brown-Kubo-Hashitsume model:
 - linear couplings, 163, 165–167
 - Markovian regime, 179–180
 - phenomenological properties, 154–157
 - dynamical properties, magnetoanisotropic nanoparticles:
 - dimensionless quantities, 129
 - numerical schemes, 107–110, 131–132
 - stochastic forms, 99–100
- Fokker-Planck equation:
 - derivation, 110–113
 - Markovian regime, 177–179
 - stationary solution, 113–115
- Garanin-Ishchenko-Panina model, 180–181
- magnetoanisotropic nanoparticles:
 - linear susceptibility, Hamiltonian formulas, 69
 - magnetization, 49–50
 - isotropic regime, 50
 - series expansions along axial parameters, 53–57
 - partition functions in free energy, 17
 - reduced linear susceptibility, 63
 - series expansion, partition function, 30–32
 - thermal equilibrium quantities, isotropic regimes, 43–44
- Markovian regime, 174–176
- nonaxial symmetry, Kramers low-damping and transition-state results, 266–267
- single-domain ferromagnetic particles:
 - domain structure, 217–220
 - Fokker-Planck equation response function, 256

- Langevin equation: (*continued*)
 statistical properties, fluctuating
 magnetic field, coupled initial
 conditions, 169
- LAPACK library:
 DGESV equation solver, 297–306
 Fokker-Planck matrix calculations, 279
 correlation times, numerical
 calculation of, 297–306
 DGEEV subroutine, 283
- Laplace transform:
 single-domain ferromagnetic particle
 relaxation, Fokker-Planck
 equation response function,
 252–256
 single-domain ferromagnetic particles,
 nonaxially symmetric particles,
 247–256
- Legendre functions:
 magnetoanisotropic nanoparticles:
 linear susceptibility, Hamiltonian
 formulas, 68–69
 nonlinear susceptibility, 75
 quantum electrodynamics (QED),
 resonance energy transfer,
 fluorescence depolarization,
 381–383
 single-domain ferromagnetic particle
 relaxation, 313–315
 Fokker-Planck equation, matrix
 formulation, 249–252
 high-energy-barrier approximation,
 axial symmetry, 242–244
 low-energy-barrier approximation,
 236
- Leibnitz formula, low-frequency
 Shliomis-Stepanov equation:
 blocking temperatures, 95–96
 $\pi/2$ law, 93–94
- Lennard-Jones potential, quantum
 electrodynamics (QED),
 resonance energy transfer, 362
- Levi-Civita symbol:
 Brown-Kubo-Hashitsume model, free
 dynamics, canonical variables,
 158–159
 magnetoanisotropic nanoparticles,
 Fokker-Planck derivation,
 111–113
- Limiting regimes, magnetoanisotropic
 nanoparticles, partition functions
 in free energy, 16–22
- Ising regime, 19
 isotropic case, 17
 longitudinal-field case, 21–22
 plane-rotator regime, 19–20
 zero-field case, 17–19
- Linear couplings, stochastic dynamical
 equations, 159–169
 fluctuating source properties, 167–169
 Hamiltonian notations, 160–163
 spin-dynamics case, 163–167
 spin-environment Hamiltonian,
 159–160
- Linear-plus-quadratic couplings,
 stochastic dynamical equations,
 169–174
 fluctuating source statistical properties,
 173–174
 Hamiltonian notations, 170–171
 spin-dynamics case, 171–173
 spin-environment Hamiltonian,
 169–170
- Linear susceptibility, magnetoanisotropic
 nanoparticles:
 approximate formulas, 70–72
 axially symmetric Hamiltonian
 formula, 67–69
 formulas for, 62–64
 heuristic treatment, 86–88
 in-phase susceptibility, 91–92
 Ising regime, 61
 isotropic regime, 61
 longitudinal bias field formula, 71–72
 longitudinal field, 62
 out-of-phase dynamical susceptibility
 and energy-barrier distribution,
 89–91
 plane-rotator regime, 62
 probing-field magnetization derivative,
 64–66

- randomly distributed anisotropy axes, 66–67
- series expansion of magnetization, 55–57
- spin ensemble dynamics, 142–144
- strong-anisotropy range, 70–71
- temperature dependence, 72–73
- tensor structure, 66
- weak-anisotropy range, 70
- Local displacement field operator,
 - quantum electrodynamics (QED), resonance energy transfer, 401–403
- Logarithmic expansion:
 - magnetoanisotropic nanoparticles, thermal equilibrium quantities, 41–43
- series expansions of free energy, 35
- London potential, quantum electrodynamics (QED), resonance energy transfer, 362
- Longitudinal bias, magnetoanisotropic nanoparticles:
 - linear susceptibility, heuristic treatment, 87–88
 - spin ensemble dynamics, 144–145
 - vs. heuristic models, 146–150
- Longitudinal energy barriers, magnetoanisotropic nanoparticles, 11–13
- bias fields, 67, 71–72
- linear susceptibility, 62, 71–72
- magnetization, 49–53
 - series expansion along anisotropy axes, 55–57
- nonlinear susceptibility, 75
- partition function and free energy, 20–22
- series expansion, partition functions, 28–29
- Shliomis-Stepanov model:
 - frozen magnetic fluid example, 98–99
 - out-of-phase susceptibility, 89–91
- spin ensemble dynamical response, 139–141
- thermal equilibrium quantities, 46–49
- Longitudinal relaxation times:
 - Fokker-Planck matrix calculations, asymptotic estimate vs. numerical calculations, 287–289
- magnetoanisotropic nanoparticles, Brown-Kubo-Hashitsume model, 123–125
- quantum electrodynamics (QED), resonance energy transfer:
 - excited molecule decay, spontaneous emission, 389
 - spontaneous emission, 384–385
- single-domain ferromagnetic particle relaxation:
 - Brown's model, 234
 - high-energy-barrier approximation, axial symmetry, 240–244
- Lorenz factor, resonance dipole-dipole interaction (RDDI), dielectric media, 375–376
- Lossy dielectrics, quantum electrodynamics (QED), resonance energy transfer, spontaneous emission, 384
- Low-damping formulas, nonaxial symmetry:
 - future research issues, 308
 - transition-state results, 265–267
 - validity of, 267–268
 - vs. intermediate-to-high damping (IHD) formula, 268–275
- Low-energy-barrier approximations, single-domain ferromagnetic particles, axial symmetry solutions, 236
- Lowest eigenvalue (λ_1) calculations:
 - Fokker-Planck matrix, 279–285
 - numerical computation vs. asymptotic estimates, 285–297
 - results tables, 316–322
- Low-frequency equations:
 - magnetoanisotropic nanoparticles:
 - linear susceptibility, heuristic treatment, 87–88
 - Shliomis-Stepanov model, 88–96
 - transverse relaxation times, 199–200

- MacLaurin series, one-dimensional Ising spin-lattice model, statistical thermodynamics, 343–345
- Magnetic aftereffect behavior, single-domain ferromagnetic particles, relaxation mechanisms, 223
- Magnetic dipole moment, defined, 330
- Magnetic energy:
 - magnetoanisotropic nanoparticles, 10–13
 - Hamiltonian expression, 9–13
- Magnetic field absorption coefficient, one-dimensional Ising spin-lattice model:
 - Ising chain formation, 354
 - statistical thermodynamics, 345
- Magnetic flux:
 - defined, 331
 - nanometric particle magnetization, 49–50
- Magnetic hysteresis, single-domain ferromagnetic particles, superparamagnetic effect, 214–216
- Magnetic moments:
 - Brown-Kubo-Hashitsume model, phenomenological equations, 154–156
 - dynamical properties,
 - magnetoanisotropic nanoparticles:
 - averaging equations, 120–122
 - individual spin stochastic trajectories, 132–138
 - probing-field direction projection, 137–138
 - one-dimensional Ising spin-lattice model, statistical thermodynamics, 340–345
- Magnetic relaxation, magnetoanisotropic nanoparticles, linear susceptibility, heuristic treatment, 86–88
- Magnetization:
 - magnetoanisotropic nanoparticles, 49–61
 - formula, 53
 - Ising regime, 50
 - isotropic regime, 50
 - longitudinal-field regime, 51–53
 - plane-rotator regime, 51
 - probing-field derivative, linear susceptibility, 64–66
 - series expansions, 53–61
 - anisotropy parameter, 53–57
 - asymptotic expansion for strong anisotropy, 57–59
 - low-field expansion, 59–61
 - single-domain ferromagnetic particle relaxation, Fokker-Planck equation response function, 254–256
- Magnetization orientation, single-domain ferromagnetic particles:
 - Brown's relaxation model, 229–234
 - domain structure, 219–220
- MagnetoAnisotropic nanoparticles:
 - dynamical properties:
 - heuristic approach, 85–99
 - stochastic approach, 99–153
 - equilibrium properties, 6–8
 - Hamiltonians, 8–13
 - linear susceptibility, 61–73
 - longitudinal-field case, energy barriers, 11–13
 - magnetization, 49–61
 - nonlinear susceptibility, 73–85
 - thermal (caloric) quantities, 41–49
 - glassy properties, 5
 - partition function and free energy, 13–22
 - definitions, 13–15
 - ising regime, 19
 - isotropic case, 17
 - limiting regimes, 16–22
 - longitudinal field case, 20–22
 - plane-rotor regime, 19–20
 - simple axially symmetric anisotropy potential, 15–16
 - zero-field case, 17–19
 - physical properties, 4–6
- Markovian regime:
 - Brown-Kubo-Hashitsume model, 179–180

- Garanin-Ishchenko-Panina model,
180–181
- phenomenological equations, 174–179
damping coefficients, 176
Fokker-Planck equations, 177–179
Langevin equations, 174–176
- single-domain ferromagnetic particle
relaxation, discrete-orientation
model, 223–224
- Matrix elements, nonaxial symmetry,
orthogonal transformations,
309–312
- Maximum anisotropy field, nanometric
magnetic particle Hamiltonian,
11–13
- Maxwell-Boltzmann distribution:
defined, 331
nonaxial symmetry, Kramers
low-damping and transition-state
results, 265–267
single-domain ferromagnetic particles,
nonaxially symmetric particles,
247–256
- Maxwell equations, quantum
electrodynamics (QED),
resonance energy transfer,
spontaneous emission, 384–385
- Meier-Saupe potential,
magnetoanisotropic nanoparticles,
dynamic properties, longitudinal
response, 148–150
- Molecular interactions, quantum
electrodynamics (QED),
resonance energy transfer:
dielectric media, dynamics in,
389–400
energy shifts, 392–396
nonrate regime, 397–400
rate description, 396–397
time evolution of quantum systems,
390–392
Hamiltonian equations, 362–363
- Monte Carlo simulations:
asymptotic expansion, 308
dynamical properties, magneto-
anisotropic nanoparticles, 99
single-domain ferromagnetic particle
relaxation, Fokker-Planck
equation response function, 256
- Multiplicative fluctuation:
dynamical properties, magneto-
anisotropic nanoparticles,
stochastic approach, 99
Fokker-Planck equation derivation,
111–113
linear couplings, spin-dynamics case,
166–167
- Néel relaxation time:
dynamical properties,
magnetoanisotropic nanoparticles:
numerical calculations, 128
spin ensemble, longitudinal
response, 139
Fokker-Planck matrix calculations:
asymptotic estimate vs. numerical
computation, 285–297
correlation time calculations,
297–306
magnetoanisotropic nanoparticles:
Fokker-Planck derivation, 113
longitudinal relaxation times,
124–125
nonaxial symmetry, Kramers
low-damping and transition-state
results, 266–267
single-domain ferromagnetic particles,
221, 223–225
discrete-orientation assumption, 223
nonaxially symmetric particles,
245–256
rigid-coupling assumption, 223
- Nernst's theorem, magnetoanisotropic
nanoparticles, thermal equilibrium
quantities, isotropic regimes, 44
- Noise-induced drift coefficient, dynamical
properties, magnetoanisotropic
nanoparticles:
averaging equations, magnetic
moments, 121–122

- Noise-induced drift coefficient, dynamical properties, magnetoanisotropic nanoparticles: (*continued*)
 Fokker-Planck equations, 115–117
 Heun numerical calculations, 130–132
- Nonaxial symmetry:
 approximation formulas:
 asymptotic formulas as function of field angle, 275–277
 asymptotic formulas vs. experimental observations, 277–279
 Brown's high-energy approximation, 256–262
 analytic solutions, 263–265
 IHD vs. LD formulas, Fokker-Planck equation, 268–275
 Kramers low-damping and transition-state results, 265–267
 validity as function of damping parameter, 267–268
 defined, 331
 single-domain ferromagnetic particles, domain structure, 220
- Nonequilibrium phenomena, nanometric particles, 4
- Nonlinear coupling:
 Brown-Kubo-Hashitsume model, spin-environment Hamiltonians, 160
 reduced motion equations, 200–207
- Nonlinear susceptibility, magnetoanisotropic nanoparticles, 73–85
 anisotropy axes, random distribution, 75–76
 approximate formulas, 80–81
 formulas, 75–79
 Ising regime, 74
 isotropic regime, 74
 longitudinal-field energy barriers, 75
 plane-rotator regime, 74
 reduced nonlinear susceptibility, 76–77
 signs of magnetization departure, 77–79
 temperature dependence, 81–85
 zero-field averages, 79–80
- Nonrate regimes, quantum electrodynamics (QED), resonance energy transfer, molecular dynamics in dielectric media, 397–400
- Nonvanishing eigenvalues:
 Fokker-Planck matrix calculations, asymptotic estimate vs. numerical calculations, 296–297
 nonaxial symmetry, vs. intermediate-to-high damping (IHD) formula, 269–275
- Numerical calculations:
 dynamical properties, magnetoanisotropic nanoparticles:
 dimensionless quantities, 128
 dimensionless stochastic Landau–Lifshitz (–Gilbert) equation, 128–129
 Fokker-Planck equation, 99
 Landau–Lifshitz equation, 100
 stochastic equations, 128–132
 Fokker-Planck matrix calculations:
 asymptotic estimate vs., 285–297
 correlation time computation, 297–306
- One-dimensional Ising spin-lattice model:
 background, 337–340
 chain structure, 353–354
 ring structure, 345–353
 statistical thermodynamics of, 340–345
 temperature effects, 354–356
- Order-disorder transition temperature, one-dimensional Ising spin-lattice model, Ising ring formation, 347–353
- Orthogonal transformation:
 Fokker-Planck matrix calculations, correlation times, numerical calculation of, 305–306
 nonaxial symmetry, 308–312
 orthogonal coordinate transformation, nonaxial symmetry, 310–312
- Oscillator-bath representation, Brown-Kubo-Hashitsume model,

- phenomenological equations, 156–157
- Out-of-phase susceptibility,
 - magnetoanisotropic nanoparticles, Shliomis-Stepanov model, longitudinal-field energy barriers, 89–91
- Overbarrier rotation, magnetoanisotropic nanoparticles, 4
 - spin ensemble, longitudinal dynamical response, 139–141
 - stochastic trajectories, individual spins, 133–135
- Overdamping phenomenon,
 - magnetoanisotropic nanoparticles, transverse relaxation time, 126–128
- Paleomagnetism, defined, 331
- Paramagnetism, defined, 331
- Partition function:
 - magnetoanisotropic nanoparticles, 13–22
 - anisotropy parameter, 29–32
 - asymptotic expansion of, for strong anisotropy, 32–35
 - definitions, 13–15
 - field expansion of, 23–29
 - ising regime, 19
 - isotropic case, 17
 - limiting regimes, 16–22
 - longitudinal field case, 20–22
 - plane-rotor regime, 19–20
 - series expansions, 22–35
 - simple axially symmetric anisotropy potential, 15–16
 - thermal equilibrium quantities, 41–43
 - zero-field case, 17–19
 - one-dimensional Ising spin-lattice model:
 - Ising chain formation, 353–354
 - Ising ring formation, 345–353
- Phase transitions:
 - magnetoanisotropic nanoparticles, longitudinal response, 149–150
 - one-dimensional Ising spin-lattice model, 338–340
- Phenomenological equations,
 - Brown-Kubo-Hashitsume model, 153–156
 - stochastic Gilbert equation, 153
 - stochastic Landau–Lifshitz equation, 153–156
- Photonic bandgap crystals:
 - quantum electrodynamics (QED), resonance energy transfer, 360–361
 - spontaneous emission, 384–389
 - resonance dipole-dipole interaction (RDDI), tensor analysis, 369–372
- Photon states:
 - quantum electrodynamics (QED), resonance energy transfer:
 - radiative (polariton) bath, 364
 - transfer rates, 364–367
 - resonance dipole-dipole interaction (RDDI), vacuum calculations, 368–369
- $\Pi/2$ law, low-frequency Shliomis-Stepanov equation, magnetoanisotropic nanoparticles, 92–94
- Plane-rotator regime:
 - magnetoanisotropic nanoparticles:
 - linear susceptibility, 62
 - nonlinear susceptibility, 74
 - nanometric magnetic particle partition functions:
 - free energy expansion, 19–20
 - magnetization, 51
 - series expansion, 28
- Pochhammer symbols:
 - Kummer functions, 184–186
 - power series expansions, 187
- Poisson bracket,
 - Brown-Kubo-Hashitsume model:
 - free dynamics, canonical variables, 158–159
 - linear couplings, 160–163
 - linear-plus-quadratic couplings, 171–173

- Potential minima
 dynamics, magnetoanisotropic nanoparticles, 106–107
- Power series expansions:
 approximate formulas, 189–190
 Kummer functions, 187
- Precession, defined, 331
- Probability distribution,
 magnetoanisotropic nanoparticles,
 partition functions in free energy,
 zero-field regime, 17–19
- Probing-field geometry:
 dynamical
 properties, magnetoanisotropic nanoparticles, magnetic moment projection, 137–138
 magnetoanisotropic nanoparticles, spin ensemble dynamics, 141–142
 magnetoanisotropic nanoparticles,
 linear susceptibility, 64–66
- Product rule, Poisson bracket,
 Brown-Kubo-Hashitsume model,
 linear couplings, 162–163
- Quantum electrodynamics (QED),
 resonance energy transfer:
 dielectric medium, 376–383
 fluorescence depolarization range dependence, 379–383
 molecular dynamics, 389–400
 nonrate regime, 397–400
 rate description, 396–397
 resonance dipole-dipole interaction (RDDI) tensor, 372–376
 retarded resonance dipole-dipole interaction (RDDI) tensor, 403–406
 time evolution, 390–392
 vibrational structure, 376–379
 far-zone spontaneous emission, 383–389
 excited molecular decay in absorbing medium, 385–389
 Hamiltonian formulation, 361–363
 local displacement field operation, 401–403
 radiative (polariton) bath, 363–364
 research background, 358–361
 resonance dipole-dipole interaction (RDDI) tensor, 367–376
 dielectric media, 372–376
 photonic bandgap structure, 369–372
 retarded calculations, dielectric medium, 403–406
 vacuum case, 367–369
 transfer rates, 364–367
- Quartic equations, Fokker-Planck matrix calculations, asymptotic estimate vs. numerical calculations, 286, 292–297
- Radiationless energy transfer, quantum electrodynamics (QED), 358–359
 dielectric media, vibrational structure, 378–379
 fluorescence depolarization, 380–383
- Radiative (polariton) bath. *See* Bath polaritons
- Raikhher-Shliomis formula, transverse dynamical susceptibility, 199
- Range dependence, quantum electrodynamics (QED),
 resonance energy transfer,
 fluorescence depolarization, 379–383
- Real-cavity model, quantum electrodynamics (QED),
 resonance energy transfer,
 spontaneous emission, 384
- Recurrence relations:
 derivations of, 186–187
 single-domain ferromagnetic particle relaxation:
 Fokker-Planck equation, matrix formulation, 251–252
 Legendre functions, and spherical harmonics, 313–315
- Reduced linear susceptibility, magnetoanisotropic nanoparticles, 63–64

- Reduced nonlinear susceptibility,
 - magnetoanisotropic nanoparticles, 76–77
- Relaxation times. *See also* Debye relaxation
 - formula derivation, 194–200
 - integral relaxation time, 194–198
 - transverse relaxation time, 199–200
 - magnetoanisotropic nanoparticles,
 - Brown-Kubo-Hashitsume model, 107, 123–128
 - longitudinal relaxation, 123–125
 - transverse relaxation time, 125–128
 - single-domain ferromagnetic particles:
 - axial symmetry solutions, 234–236, 234–244
 - Brown's high-energy-barrier approximation, 236–244
 - Brown's low-energy-barrier approximation, 236
 - Brown's model, 224–234
 - discrete-orientation model, 222–223
 - domain structure, 216–220
 - mechanisms, 220–222
 - Néel model, 223–224
 - superparamagnetic effect, 212–216
- Remanant magnetization:
 - defined, 331
 - single-domain ferromagnetic particles,
 - superparamagnetic effect, 214–216
- Representing matrix, nonaxial symmetry,
 - orthogonal transformations, 309–312
- Resonance dipole-dipole interaction
 - (RDDI), quantum electrodynamics (QED), resonance energy transfer:
 - radiative (polariton) bath, 364
 - retarded tensor calculations, 403–406
 - tensor analysis, 367–376
 - dielectric media, 372–376
 - photonic bandgap structure, 369–372
 - retarded calculations, dielectric medium, 403–406
 - vacuum case, 367–369
 - transfer rates, 366–367
- Resonance energy transfer, quantum electrodynamics (QED):
 - dielectric medium, 376–383
 - fluorescence depolarization range dependence, 379–383
 - molecular dynamics, 389–400
 - nonrate regime, 397–400
 - rate description, 396–397
 - resonance dipole-dipole interaction (RDDI) tensor, 372–376
 - retarded resonance dipole-dipole interaction (RDDI) tensor, 403–406
 - time evolution, 390–392
 - vibrational structure, 376–379
- far-zone spontaneous emission, 383–389
 - excited molecular decay in absorbing medium, 385–389
- Hamiltonian formulation, 361–363
- local displacement field operation, 401–403
- radiative (polariton) bath, 363–364
- research background, 358–361
- resonance dipole-dipole interaction (RDDI) tensor, 367–376
 - dielectric media, 372–376
 - photonic bandgap structure, 369–372
 - retarded calculations, dielectric medium, 403–406
 - vacuum case, 367–369
 - transfer rates, 364–367
- Response function, single-domain
 - ferromagnetic particle relaxation, Fokker-Planck equation, 252–256
- Retarded dipole-dipole coupling, quantum electrodynamics (QED),
 - resonance energy transfer, molecular dynamics in dielectric media, 392–396
- Rigid-coupling assumption, single-domain ferromagnetic particles,
 - Néel relaxation model, 223

- Saddle points, nonaxial symmetry:
 Brown's high-energy approximation,
 257–262
 vs. intermediate-to-high damping
 (IHD) formula, 273–275
- Schottky effect, magnetoanisotropic
 nanoparticles, zero-field regimes,
 46
- Second-rank tensor,
 Brown-Kubo-Hashitsume model,
 phenomenological equations,
 155–156
- Self-adjoint operator, Fokker-Planck
 equation, Sturm-Liouville
 problem, 119–120
- Series expansions, magnetoanisotropic
 nanoparticles:
 anisotropy axes, random distribution
 averages, 35–36
 anisotropy parameter, expansion in
 powers of, 38–39
 field expansions, 37–38
 free energy, 35–40
 Kummer functions, 187–189
 asymptotic formula, large positive
 argument, 187–188
 power series, 187
 logarithmic expansion, 35
 low-field expansion of, 59–61
 magnetization, 53–61
 anisotropy power, expansion of
 powers, 53–57
 asymptotic expansion for strong
 anisotropy, 40, 57–59
 partition function, 22–35
 anisotropy parameter, 29–32
 asymptotic expansion for strong
 anisotropy, 32–35
 field expansion, 23–29
- Shliomis-Stepanov equation,
 magnetoanisotropic nanoparticles:
 frozen magnetic fluids, 97–98
 linear susceptibility, heuristic
 treatment, 88, 183
 longitudinal response, 146–150
 low-frequency model, 88–96
 blocking energies, energy-barrier
 distribution, 94–96
 in-phase linear dynamical
 susceptibility, 91–92
 out-of-phase linear susceptibility,
 energy-barrier distribution, 89–91
 $\pi/2$ law, 92–94
- Single-domain ferromagnetic particles,
 relaxation times:
 axial symmetry solutions, 234–236
 Brown's high-energy-barrier
 approximation, 236–244
 Brown's low-energy-barrier
 approximation, 236
 Brown's model, 224–234
 discrete-orientation model, 222–223
 domain structure, 216–220
 mechanisms, 220–222
 Néel model, 223–224
 nonaxially symmetric particles,
 244–256
 magnetic response functions,
 244–247
 matrix formulation, Fokker-Planck
 equation, 249–252
 response function and initial
 conditions, 252–256
 superparamagnetic effect, 212–216
- Spatial modulation, resonance
 dipole-dipole interaction (RDDI),
 photonic bandgap crystals,
 369–372
- Specific heat at constant field, magneto-
 anisotropic nanoparticles, thermal
 equilibrium quantities, 42–43
 longitudinal-field regimes, 47–49
 zero-field regimes, 44–46
- Spherical coordinates:
 Fokker-Planck equation, dynamical
 properties, magnetoanisotropic
 nanoparticles, 117–118
 nonaxial symmetry, orthogonal
 transformation, 312
 single-domain ferromagnetic particles:
 Brown's relaxation model, 230–234
 domain structure, 219–220

- Fokker-Planck equation:
 - matrix formulation, 249–252
 - response function, 252–256
- Legendre functions, 313–315
- Spin ensembles:
 - dynamical response, 138–153
 - bias field absence, 139–144
 - linear couplings, 163–167
 - linear-plus-quadratic couplings, 171–173
 - linear susceptibility vs. heuristic models, 146–153
 - in longitudinal bias field, 144–146
 - longitudinal response, 139–141
 - random distribution of anisotropy axes, 142–144
 - transverse response, 141–142
- one-dimensional Ising spin-lattice model:
 - background, 337–340
 - chain structure, 353–354
 - ring structure, 345–353
 - statistical thermodynamics of, 340–345
 - temperature effects, 354–356
- Spin-environment Hamiltonians:
 - Brown-Kubo-Hashitsume model:
 - linear couplings, 159–160
 - linear-plus-quadratic couplings, 169–170
 - phenomenological equations, 156–157
- Markovian regime, phenomenological equations, 175–176
- “Spin solids,” one-dimensional Ising spin-lattice model, thermodynamics, 340–345
- Spontaneous emission, quantum electrodynamics (QED):
 - far-zone energy transfer, 383–389
 - excited molecule decay, 385–389
 - research background, 383–385
 - resonance energy transfer, 359–361
- Spontaneous magnetization, single-domain ferromagnetic particles, domain structure, 218–220
- Static case, single-domain ferromagnetic particle relaxation, Brown’s model, 232–234
- Stationary properties:
 - Brown’s high-energy approximation, nonaxial symmetry, 257–262
- magnetoanisotropic nanoparticles, Fokker-Planck equation, 113–115
- Statistical-mechanical averaging, magnetoanisotropic nanoparticles: magnetization formulas, 53 partition function and free energy, 14–15 thermal equilibrium quantities, 41–43
- Statistical properties:
 - linear couplings, fluctuating terms, 167–169
 - one-dimensional Ising spin-lattice model, thermodynamics, 340–345
- Stern-Gerlach experiments, nanometric particle magnetization, 49–50
- Stochastic differential equation, magnetoanisotropic nanoparticles, Brown-Kubo-Hashitsume model, 107
- Stochastic Gilbert equation:
 - Brown-Kubo-Hashitsume model, phenomenological properties, 153–156
 - magnetoanisotropic nanoparticles, 107–128
 - Langevin dynamical equations, 109–110
- Stochastic Landau–Lifshitz equation:
 - Brown-Kubo-Hashitsume model, phenomenological properties, 153–156
 - dynamical properties, magnetoanisotropic nanoparticles, dimensionless quantities, 128–129
 - magnetoanisotropic nanoparticle dynamics, 109–110
 - averaging equations, 121–122
 - Fokker-Planck derivation, 111–113
 - numerical schemes, 131–132

- Stochastic Landau–Lifshitz equation:
 (*continued*)
 stationary solution, Fokker-Planck equation, 114–115
- Stochastic resonance, magnetoanisotropic nanoparticles, spin ensemble, longitudinal dynamical response, 141
- Stochastic theory:
 defined, 331
 dynamical equations:
 Brown-Kubo-Hashitsume model, 179–180
 free dynamics and canonical variables, 157–159
 Garanin-Ishchenko-Panina model, 180–181
 linear couplings, 159–169
 fluctuating source properties, 167–169
 Hamiltonian notations, 160–163
 spin-dynamics case, 163–167
 spin-environment Hamiltonian, 159–160
 linear-plus-quadratic couplings, 169–174
 fluctuating source statistical properties, 173–174
 Hamiltonian notations, 170–171
 spin-dynamics case, 171–173
 spin-environment Hamiltonian, 169–170
 Markovian regime, 174–179
 damping coefficients, 176
 Fokker-Planck equations, 177–179
 Langevin equations, 174–176
 phenomenological equations, 153–156
 spin environment interactions, 156–157
 summary, 181–182
 magnetoanisotropic nanoparticles, 6
 Brown-Kubo-Hashitsume model, 107–128
 deterministic dynamics, classical spins, 101–107
 dynamics, 99–153
 individual spin trajectories, 133–138
 numerical method, 128–133
 spin ensemble dynamics, 138–153
 Stochastic trajectories, dynamical properties, magnetoanisotropic nanoparticles, 132–133
 implementation, 132–133
 individual spins, 133–138
 Stratonovich stochastic calculus:
 dynamical properties, magnetoanisotropic nanoparticles, 99
 Fokker-Planck equations, 115–117
 Landau–Lifshitz (–Gilbert) equation, 100
 Langevin equation, 109–110
 numerical calculations, 130–132
 stochastic trajectories, individual spins, 133–138
 Fokker-Planck equation, Markovian regime, 177–179
 Strong anisotropy, magnetoanisotropic nanoparticles:
 asymptotic expansion, partition function, 32–35
 magnetization expansion, 57–59
 series expansion, 40
 linear susceptibility formulas, 70–71
 nonlinear susceptibility formulas, 81
 Sturm-Liouville equation:
 axially symmetric Fokker-Planck equation, 118–120
 integral relaxation time formula, 194
 magnetoanisotropic nanoparticles,
 dynamic properties, longitudinal relaxation times, 123–125, 147–150
 single-domain ferromagnetic particle relaxation, high-energy-barrier approximation, axial symmetry, 241–244
 Subtraction techniques, dynamical properties, magnetoanisotropic

- nanoparticles, stochastic trajectories, 132–133
- Summation indices, magnetoanisotropic nanoparticles, asymptotic expansion, partition function, 34–35
- Superparamagnetism:
 - magnetoanisotropic nanoparticles, equilibrium properties, 6–8
 - nonaxial symmetry, asymptotic expansion, 277–279
 - physical properties of, 4–6
 - single-domain ferromagnetic particle relaxation times, 212–216, 223
- Symmetric anisotropy potential, magnetoanisotropic nanoparticles, partition function, 15–16
- Taylor series:
 - nonaxial symmetry:
 - Brown's high-energy approximation, 260–262
 - vs. intermediate-to-high damping (IHD) formula, 268–275
 - power series expansions, 187
 - single-domain ferromagnetic particle relaxation, high-energy-barrier approximation, axial symmetry, 237–244
- Temperature dependence:
 - magnetoanisotropic nanoparticles:
 - linear susceptibility, 72–73
 - nonlinear susceptibility, 77–79, 81–85
 - reduced linear susceptibility, 63–64
 - stochastic trajectories, individual spins, 135–137
 - one-dimensional Ising spin-lattice model:
 - Ising chain formation, 353–354
 - Ising ring formation, 346–353
 - phase transitions, 339–340
- Tensor structure:
 - magnetoanisotropic nanoparticles, linear susceptibility, 66
 - resonance dipole-dipole interaction (RDDI):
 - dielectric media, 372–376
 - photonic bandgap structure, 369–372
 - retarded calculations, dielectric medium, 403–406
 - retarded RDDI, dielectric medium, 403–406
 - vacuum calculations, 367–369
- Thermal equilibrium:
 - defined, 331
 - magnetoanisotropic nanoparticles, 6–8, 41–49, 182
 - definitions, 41–43
 - isotropic case, 43–44
 - longitudinal field case, 46–49
 - spin ensemble dynamics, 142
 - zero-field case, 44–46
- Thermodynamic quantities. *See* free energy; partition function
- Thermodynamics:
 - magnetoanisotropic nanoparticles, thermal equilibrium quantities, 41–43
 - one-dimensional Ising spin-lattice model, 340–345
- Thermoremanent magnetization, Fokker-Planck matrix calculations, correlation times, numerical calculation of, 306
- Time-energy uncertainty principle, resonance dipole-dipole interaction (RDDI), photonic bandgap crystals, 370–372
- Time evolution, quantum electrodynamics (QED), resonance energy transfer, molecular dynamics in dielectric media, 390–392
- Transfer rates, quantum electrodynamics (QED), resonance energy transfer, 364–367
- Transition-state theory:
 - nonaxial symmetry, Kramers low-damping and transition-state results, 266–267

- Transition-state theory: (*continued*)
 single-domain ferromagnetic particle
 relaxation, high-energy-barrier
 approximation, axial symmetry,
 237–244
- Transverse relaxation time:
 effective formulas, 199–200
 magnetoanisotropic nanoparticles,
 Brown-Kubo Hashitsume model,
 125–128
 quantum electrodynamics (QED),
 resonance energy transfer:
 excited molecule decay, spontaneous
 emission, 389
 spontaneous emission, 384–385
 single-domain ferromagnetic particle
 relaxation, Brown's model, 234
- Transverse response, magnetoanisotropic
 nanoparticles:
 spin ensemble dynamics, 141–142,
 146
 vs. heuristic models, 150–153
- Uniaxial anisotropy, defined, 331
- Unit vector triad, nonaxial symmetry,
 orthogonal transformations,
 308–312
- Vacuum calculations:
 quantum electrodynamics (QED),
 resonance energy transfer,
 molecular dynamics in dielectric
 media, 395–396
 nonrate regimes, 398–400
 resonance dipole-dipole interaction
 (RDDI), 367–369
- Vacuum Rabi splitting, resonance
 dipole-dipole interaction (RDDI),
 photonic bandgap crystals, 372
- Vector-addition rules, single-domain
 ferromagnetic particle relaxation,
 Legendre functions, and spherical
 harmonics, 315
- Vibrational structure, quantum
 electrodynamics (QED),
 resonance energy transfer,
 dielectric media, 376–379
- Virtual-cavity case, quantum
 electrodynamics (QED),
 resonance energy transfer,
 spontaneous emission, 384
- Waldram variables, one-dimensional Ising
 spin-lattice model, 354–356
- Weak anisotropy, magnetoanisotropic
 nanoparticles:
 asymptotic expansion of magnetization,
 59
 linear susceptibility formulas, 70
 nonlinear susceptibility formulas, 80
 series expansion of magnetization,
 53–57
- Weak-damping case, dynamical
 properties, magnetoanisotropic
 nanoparticles, axially symmetric
 Hamiltonians, deterministic
 Landau–Lifshitz equation, 104
- Websites, www.netlib.no/netlib/lapack,
 279
- Weiss domains, single-domain
 ferromagnetic particles, 216
- Weiss mean-field theory, one-dimensional
 Ising spin-lattice model, research
 background, 338–340
- Wigner symbols, single-domain
 ferromagnetic particle relaxation,
 Legendre functions, and spherical
 harmonics, 315
- Zeeman energy:
 dynamical properties,
 magnetoanisotropic nanoparticles,
 axially symmetric Hamiltonians,
 deterministic Landau–Lifshitz
 equation, 104
 magnetoanisotropic nanoparticles:
 dynamic properties, longitudinal
 relaxation times, 123–125
 thermal equilibrium quantities, 43
 longitudinal-field regimes, 49

- Zero-field-cooled magnetization:
 Fokker-Planck matrix calculations,
 correlation times, numerical
 calculation of, 306
 integral relaxation time, 198
 magnetoanisotropic nanoparticles:
 deterministic dynamics,
 105–106
 free energy expansion, 17–19
 nonlinear susceptibility,
 79–80
 series expansion, 24–25
 thermal equilibrium quantities,
 44–46
 nonaxial symmetry:
 asymptotic expansion, 277–279
 future research issues, 308
 one-dimensional Ising spin-lattice
 model, statistical
 thermodynamics, 342–345
- Zero-point energy, quantum
 electrodynamics (QED),
 resonance energy transfer,
 radiative (polariton) bath, 364

