

1

The world as it was and the world as it is

1.1 Early ecological history

Our planet is old, around 4.53 billion years on current estimates, but we humans are very young. Only about 100 000 years have passed since we emerged distinctively as *Homo sapiens* from our previous ancestors. They had had comparatively little effect on the planet, and so did we until the last 15 000 years or so. Before then, the planet changed slowly but continually, under natural geological forces: volcanic eruption, plate separation and continental drift; natural cycles in the Earth's orbit around the Sun; and small changes in the rate at which the Sun emits energy. Its surface changed just as much because the inevitabilities of evolution were producing a succession of organisms that altered the chemistry of the atmosphere and oceans. Around 2 billion years ago, an atmosphere that had previously been free of molecular oxygen was steadily oxygenated because one group of bacteria, the Cyanobacteria (Fig. 1.1), had evolved the ability to use water as the hydrogen donor needed to reduce carbon dioxide in photosynthesis, and released oxygen as a by-product.

This created problems for a biosphere maintained by anaerobic bacteria, because free oxygen was toxic, but one consequence appears to have been the evolution of the eukaryotic cell, in which, through processes of symbiosis, host cells, probably Archaeobacteria, engulfed other bacteria whose enzymes could function deep in the combined cell, away from the increasing oxygen concentrations in the environment. Oxygen then built up steadily

in the atmosphere until concentrations were high enough (Fig. 1.2) for diffusion to be able to support bigger, multicellular organisation, between 500 and 600 million years ago. Multicellularity allows specialist systems to develop and was rapidly adopted. Multicellular systems could cope with conditions on land, and a biodiversity previously confined to water was joined by one that could take advantage of very high oxygen concentrations in the air. Oxygen is not very soluble in water (see Sections 5.2 and 5.3). On land there was also a greater supply of light energy (water absorbs the Sun's radiation very quickly; see Sections 7.2 and 7.3). In turn, these enhanced conditions allowed the eventual evolution of mammals that could start to modify conditions to their own ends through a high brain capacity. We were born.

The Earth is well supplied with the twenty or so elements that natural selection has used to produce and maintain living systems, but it has limited supplies, in available form at the surface, of some of them. The stock must be recycled. Moreover, liquid water is essential for living cells to function, and, from the end of the Hadean Period 4 billion years ago, when Earth had cooled sufficiently for water to condense from the steamy atmosphere of volcanic gases, there was established a water cycle. The essence of this is that water, evaporated from the oceans and land surfaces through solar heating, moves upwards or polewards in the atmosphere, cools and condenses. It falls as rain or snow, and runs off the land and back to the ocean.




Figure 1.1 Cyanobacteria, which are now ubiquitous in soils, fresh and salt waters, had a pivotal role in the history of the biosphere. They evolved the ability to use water as a hydrogen donor in photosynthesis, thus releasing molecular oxygen as a by-product. Individual cells of cyanobacteria (inset) are generally very small (around 1–2 μm) but may aggregate in much bigger filaments and colonies, sometimes occurring so abundantly as to colour the water prominently, as in this temple tank in Nepal. Ancient fossils suggest that the range of forms of cyanobacteria have not changed greatly since they first evolved. (Reproduced with permission of K. J. Irvine. Inset reproduced with permission of Matthew J. Parker.)




Figure 1.2 Reconstructed changes in oxygen and carbon dioxide concentrations in the Earth's atmosphere over geological time. The envelopes indicate the variation calculated from different geological models, but the trends are clear. Major events in evolution are also shown. A bar is the unit of atmospheric pressure. Current total pressure is close to 1 bar (or 10^0 on the logarithmic scale used). (Based on Mojzsis 2001.)




Figure 1.3 Linkages among parts of the freshwater system, the catchment area of the land, the atmosphere and the water cycle.

In doing so, it is retained for a time in a continuous system of hilly streams and rivers, groundwaters, pools, wetlands and lakes, and then in floodplain rivers and estuaries that connect them with the coastal seas and oceans (Fig. 1.3). Water dissolves a huge range of substances and carries them with it during the liquid phases of this cycle. Some are absorbed by aquatic organisms; others, like nitrogen compounds, are converted quickly to gases by bacteria and returned to the atmosphere, and yet others contribute to the saltiness of the ocean or are precipitated out into sediments and newly forming rocks. Recycling of these latter elements is long term. Movements of the Earth's plates against one another raise new mountains over millions of years and weathering slowly re-releases substances usable by organisms. Recycling of many essential elements, however, has to be much more rapid and depends on biological processes. Water thus acts like

a bloodstream for the Earth (Fig. 1.4), its rivers and lakes the equivalents of arteries and veins, its evaporative surfaces a sun-driven heart that pumps the water around, and its basins, especially lakes and wetlands, its digestive and excretal systems, foci of biological activity that shuttle dissolved substances between organisms and the water.

There appears to be some overall linkage of these activities, though we have no idea how it is achieved. The oxygen and carbon dioxide concentrations in the atmosphere and the saltiness of the sea have been maintained for a long time within limits that allow the persistence of liquid water and of multicellular organisms, despite geological forces that could threaten this. Carbon dioxide remained between 190 and 260 ppm by volume for at least a million years until very recently – a period that included a number of advances of the polar and mountain glaciers – and oxygen concentrations have been



Figure 1.4 Seen from space, the freshwater systems of the Earth support the analogy that they are the bloodstream of the biosphere. The many mouths of the River Ganges shown here discharge into the Indian Ocean. (Reproduced with permission of USGS EROS Data Center Satellite Systems Branch.)

Table 1.1 Composition of the Earth's atmosphere, compared with those of its closest planets, Mars and Venus. Equilibrium Earth is calculated from chemical models that assume that all possible reactions are allowed to run to equilibrium. Present-day composition is as measured on Earth, and for Mars and Venus is deduced from spectroscopic measurements. (Based on Lovelock 1979. Reproduced with permission of Oxford University Press.)

	Venus	Equilibrium Earth	Mars	Earth as it is
Carbon dioxide (%)	98	98	95	0.03
Nitrogen (%)	1.9	1.9	2.7	79
Oxygen (%)	Trace	Trace	0.13	21
Argon (%)	0.1	0.1	2	1
Surface temperature (°C)	477	290	-53	13
Total pressure (bars)	90	60	0.0064	1

around 21% for at least as long. Carbon dioxide was prevented from rising much higher through the storage of carbon compounds in waterlogged soils, peats and lake sediments, whilst gases produced by living organisms, such as methane and dimethyl sulphide, react with oxygen and temper its concentration downwards. Much higher oxygen concentrations would promote uncontrollable forest and grassland fires and indeed this had happened earlier in geological history.

The chemistry of both seawater and the atmosphere is maintained in a non-equilibrium state by the activities of living organisms (Table 1.1). Without them, we would have a much hotter planet and possibly no liquid water, and then only with a crushing atmospheric pressure. But how this system is maintained in an apparently orderly way is a mystery. There appears to be cooperation, but cooperation and natural selection do not easily fit together.

1.2 The more recent past

We know a great deal about the ecology of our planet but the detail is greater for the past few million years, and particularly for the last few tens of thousands of years, than for any time previously. This latter time embraces the final melting back of the ice sheets that had advanced and retreated over the previous several million years. As ice advances, it bulldozes the land surfaces, widens pre-existing river valleys and changes the former courses of rivers. It scrapes out new basins for eventual lakes, and changes the outlines and depths of previous ones if they were large enough for their basins still to be recognisable. When ice melts, enormous amounts of fractured rock, gravel, sand and finer sediments wash out from under the glaciers, and may be deposited as moraines or in extensive washout plains. Organisms from previous periods of retreat, when the land was exposed and ecological communities developed, do survive, but they are a small proportion of the former communities.

We are not talking about a few metres of ice and snow, the consequences of even a severe blizzard. Over the land in the upper forty degrees of latitude in both the southern and the northern hemispheres, we must imagine ice several kilometres in thickness. We must hear the deafening noise of crashing bergs and roaring floods at the ice front; and towards the Equator, we must see lands that were frosted in winter, melting only in summer for their rivers to flow. Closer to the Equator, there was not such devastation, but it was cooler and wetter: one of the reasons for ice to advance is that the Earth's orbit is slightly farther from the Sun than at other times and its tilt on its axis slightly less extreme. Heating and evaporation were reduced; droughts were still frequent in arid areas, but lakes were bigger, rivers flowed more prolifically. And eventually, from a reservoir of species in the now warmer lands, came a steady migration of organisms polewards as the ice retreated and the now temperate and polar lands were re-exposed.

The final advance of the ice and its melting back did not completely re-set ecological history, even for the temperate and polar lands, but particularly not for the regions that were not covered by ice. There continued a long process of evolutionary change since the times that they too may have been devastated by even more severe glaciations some hundreds of millions of years before, and particularly following the impact of collision with a large meteor in what is now Mexico 66 million years ago. At that time, the continents were approaching their present positions but were still on the move. North and South America had not yet joined up, India was yet to smash into Asia and cause the upthrust of the Himalayas, and Australia was only just separating from Antarctica and moving equatorward, whilst Antarctica moved farther south. The land surfaces that were to become the present tropics and warm-temperate regions were mostly ancient, having been subjected to millions of years of weathering down to flat plains (Fig. 1.5), whose soils were deep and had been leached for long periods, in contrast to the glaciated regions where new soils were formed from rock freshly plucked and scoured by the ice (Fig. 1.6).

There were to be immense consequences for freshwaters, because the minerals that leach out from the soils determine the nature of the water that fills the basins. And cutting across this simple picture of ancient, un-iced surfaces and new just-glaciated ones, was the legacy of past plate movements. There were (and still are) huge mountain chains along the western edges of the Americas, continuing across the arc of the Aleutian Islands around the far side of the Pacific through Russia and Japan, and dwindling into South East Asia. Sixty-six million years ago, the Himalayas were yet to form, but the Carpathians, the Alps and the Appalachians, formed by earlier plate collisions, were still high features. In many places, even under the ice, the effects of plate movement continued to be reflected in volcanic activity. The flat continent of Africa acquired, only a million or two years ago, deep basins, the rift valleys,



Figure 1.5 Mount Conner, Northern Territory, Australia. Ancient land surfaces have been planed flat by erosion, often have deep but infertile soils, from which minerals have long been leached, leaving only bright iron oxides and quartz with little organic matter. Such landscapes are now mostly found in the warm temperate and tropical regions. (Reproduced with permission of Gabriele Delhay.)



Figure 1.6 Youthful landscapes are those where rock has been exposed by volcanic action or scoured and ground by ice. Soils are shallow and still forming but are rich in minerals, which leach to the streams and rivers. In glaciated landscapes like this near Cader Idris in Wales, with Cregennan Lake in the foreground, the northerliness and dampness of the climate also promotes the build-up of organic matter in the soils.

in which rest the East African Great Lakes. Volcanoes continued to erupt, leaving isolated high mountains, Kenya and Kilimanjaro, for example, with dozens of streams and small tarns. Lakes like Kivu (Fig. 1.7) were

formed as lava flows dammed the rivers. Nothing has rested for very long.

We enter the last 20000 years, even the last few million years, on a still changing stage but one that would be easily recognisable to a



Figure 1.7 Fishing on Lake Kivu, which is shared between the Democratic Republic of the Congo and Rwanda. The lake was formed by the damming of the River Rutshuru sometime before 15 000 years ago, when the Virunga volcanoes to the north erupted and filled the rift valley between Lake Edward and Lake Tanganyika with lava flows. This changed the drainage so that the lake overflows now to the south whereas previously the flow was northwards into Lake Edward. The lake is very deep and does not mix from top to bottom and so has accumulated very large stocks of carbon dioxide and methane in its deeper layers. (Reproduced with permission of Steve Evans.)

modern biologist. The groups of organisms that are now familiar to us were by then already dominating. We are not dealing with trilobites and dinosaurs, huge trees of the fern families, or even exotic microorganisms. Indeed there has been a great turnover in individual species and genotypes, even over a hundred thousand years, but all the groups with which we are now familiar were established before the disruption of the meteorite collision 66 million years ago. In freshwaters, the bacteria, algae and protozoans, though now reclassified into seven or so new kingdoms that would be unfamiliar to a biologist trained just twenty-five years ago, are old groups.

1.3 Characteristics of freshwater organisms

The freshwater invertebrates have been dominated by the annelid worms, molluscs, crustaceans and insects for many millions of years, and the flowering plants, birds and

mammals had established their dominance over the ferns and mosses, amphibians and reptiles long previously. But it is, and was, a highly changing world. Freshwaters contain only a very small proportion of the world's water. Most water is contained in a set of linked oceans that has been a permanent body of water (though greatly changing its shape) since water first condensed out of the atmosphere around 4 billion years ago. The next most abundant source is in the polar and mountain glaciers, then the soil and groundwaters. Only a tiny fraction, a percentage point or two, is stored in lakes or flowing through rivers at any one time, but with the even tinier proportion in the atmosphere, this water maintains the movements and exchanges of nutrients and minerals throughout the land. Its amounts in any given place are basically determined by climate, but within the climatic zones, weather has an immense effect. There are droughts and floods; lake levels can rise and fall; streams, even rivers, can move sideways,




Figure 1.8 Freshwater animals often produce structures that allow them to survive drought or other difficult conditions by reducing their metabolism in protective shells or spores. A–D are cysts of protozoans; E and F are resting eggs of tardigrades (water bears). G shows dispersive structures (gemmules) of a freshwater sponge; H is a resting egg of a freshwater shrimp (*Eulimnadia*); and J and K the resting eggs of rotifers. L and M are side and front views of a statoblast of a bryozoan. (Based on Pennak 1985. Reproduced with permission of Oxford University Press.)

even dry up, and there is little detailed predictability from year to year. The loss and re-formation of the entire system in the polar and temperate regions during the recent glaciations is a manifestation of this. Unless it lives in a particularly big lake, a freshwater organism can rely on little. Its habitat may be very different next year.

Freshwater communities are well adapted to change and disturbance. They might produce resting stages to survive a dry period; they might be very efficient at dispersal as spores or cocoons (Fig. 1.8) or by flood or on the wing. They generally have high tolerance of varying habitat conditions and they invest more energy in reproduction and replacement than in developing liaisons among different species, in breeding colours or elaborate behaviours. When these features are prominent, the organisms will usually be in one of the more stable basins: a large, deep tropical lake for example. In contrast, marine organisms can rely on a permanent water body. Tides vary but highly predictably; ocean currents involve such huge volumes of water that

they can be relied upon; and below a shallow surface layer is a steady habitat, tranquil even, that supports multicoloured animals that have developed many mutually supportive symbioses (Fig. 1.9). As in freshwaters, the open-water planktonic habitat in the oceans is hazardous, but the bottom is steadier and that is where most of the biodiversity lies. Marine organisms are generally fussy. It is much easier to maintain a freshwater aquarium than a marine one.

1.4 Freshwater biodiversity

Freshwaters are sometimes thought of as Cinderellas when it comes to richness of diversity. Apparently they have many fewer species, though not necessarily families, than the land, or especially the oceans. But the more valid comparison is in richness compared with extent of habitat. Freshwaters occupy a tiny proportion of the land space or total water space, but have broadly comparable diversities in terms of order of magnitude

Figure 1.9 Marine animals are frequently brightly coloured compared with freshwater ones. The ocean is a much more predictable habitat than freshwaters and investment in warning or breeding colours is worthwhile, whereas in freshwaters, more energy is invested in reproduction in a habitat where death rates are often high. (Reproduced with permission of Mark Peter.)



Table 1.2 Relative biodiversity of fishes in marine and freshwater habitats. NA, not applicable. (Based on Cohen 1970; Horn 1972; and Balian *et al.* 2008.)

Habitat	Area (million km ²)	Volume (million km ³)	Mean depth	Species number (percentage)	Species per million km ²	Species per million km ³
<i>Fish</i>						
Oceans	361	1371	3.8 km	18 000 (58)	49.9	13.1
Freshwaters	1.5	0.13	87 m	13 000 (41)	8667	100 000
<i>Total animals</i>						
Oceans	361	1371	3.8 km	349 000 (26.8)	967	255
Freshwater	1.5	0.13	87 m	126 000 (9.7)	84 000	969 200
Land	149	NA	NA	827 000 (63.5)	5550	NA

to land or marine biota (Table 1.2). Speciation has been much more vigorous in freshwaters, perhaps because of the continual moderate and occasionally high disturbance that they experience. The continually shifting habitat means continual evolutionary adjustment. Freshwater fish are especially diverse.

After the ice retreated, there was plentiful water flow in the rivers, and on flat, long-eroded landscapes, irregularities held water

and created extensive networks of shallow wetlands or deeper ponds and lakes. Rivers meandered over wide floodplains, carrying silt from the uplands, depositing it in the spring or wet season floods, and creating a mosaic of habitats. The land was covered in natural vegetation: forests of many kinds, grasslands in drier areas, deserts in the most arid. But most deserts had some vegetation and plant-lined ephemeral streams ran

through them. Estuaries, mangrove swamps and salt marshes made linkages with offshore sea grass meadows, coral and algal reefs and eventually the deeper ocean. It was the continuous bloodstream needed to maintain a functioning biosphere. Rich in species, rich in the processes it sustained, magnificent in its extent and scope, it stitched together the land systems into a continuous, unboundaried whole. Its continuity allowed migrations of animals to take advantage of seasonal availability of fodder; its undisturbed forests stored water and mitigated flooding, maintaining habitats wet even during low rainfall periods.

Large amounts of organic matter were washed from the land as leaf litter, twigs, downed timber and dissolved substances, and processed by stream, river and lake communities. Some of this material was respired. Other material was stored in sediments and peats, and the rest washed downriver to be processed in the ocean. The freshwaters also effectively controlled the Earth's nitrogen cycle, as well as being significant influences on the carbon cycle. Although there is plenty of molecular nitrogen available in the

atmosphere, it is only available to the biosphere when fixed into amino groups by bacteria that occur in waterlogged soils, root nodules and other places where oxygen is reduced to low levels. And when they have been fixed and taken up, recycled and decomposed, ammonium and nitrate ions are used by bacteria, again in waterlogged places, for the release of energy, and returned to nitrogen gas in the process. The floodplains, with their productive communities, were extensive features of every river and their high fertility, from plentiful water and accumulated silt, attracted large flocks of birds and herds of mammals (Fig. 1.10), which daily moved between the drier uplands and the plains, redistributing nutrients and dung. Even close to the edge of the retreating ice, the herds were very large.

The system was controlled by evolutionary processes capable of responding rapidly when necessary. It managed itself even as a naturally changing climate altered the compositions of the communities. There are several features necessary for independent existence, adjustment and repair of ecological systems: relatively scarce nutrients are



Figure 1.10 Bison in Yellowstone National Park, USA. Floodplains, fertilised by eroded silt from upstream and with rich, damp grasslands and wetlands, attract large herds of grazers, where such animals are still protected.




Figure 1.11 *Homo sapiens* progressively differentiated from common ancestors with the other great apes, through a complex of *Paranthropus*, *Australopithecus* and several other *Homo* species. We were in essentially our present form by only around 100 000 years ago.

conserved in biomass and the risk of them being lost to the ocean by leaching and washout is minimised; physical characteristics like river flows are determined by natural climate and fall into patterns, albeit variable ones, to which organisms adjust through natural selection; food webs are intact, with top predators always present; and the system is extensive and interconnected. Only natural barriers like mountain chains and tracts of ocean interrupted a potential free flow of organisms. It was an Eden that required no cash budget and functioned indefinitely.

1.5 A spanner in the works?

But even as this system was emerging from the effects of the ice, it was threatened. For 13 million years, hominoids had been differentiating from great apes and steadily

speciating. Several *Australopithecus* and *Homo* species developed, all of them omnivores or meat-eaters, and by 100 000 years ago the three or four coexisting *Homo* species collapsed into a single species, *Homo sapiens*: ourselves (Fig. 1.11). *Homo* had survived the glaciations, emerging from tropical refuges to recolonise the temperate regions in the warmer periods between the many small and large advances of the ice, and finally followed the ice margin into the Arctic. Before the start of the Holocene, 11 700 years ago, we had a foothold on every continent except Antarctica but we were just one among dozens of effective predators. We had, however, a number of advantages owing to our very large brain to body ratio, a cerebellum within the brain complex that was large enough to allow a degree of planning and foresight, and a sophisticated ability to make tools and manipulate our environment rather than merely react to changes in it.

As the Holocene unfolded, we were increasingly to change the world, completely ignorant at the time that we could alter the biosphere in ways that would bring severe problems only a few thousand years later. Indeed for most of our history we have perceived ourselves as a force for the good, rather than a spanner in the works.

Climate was still changing 15 000 years ago, and this confounds interpretations, but strong evidence suggests that our first major impact was to hunt out and make extinct many of the other larger mammals that had survived the ice alongside our immediate ancestors. We were, and are, very effective invasive species; we had our own predators but we learnt to use shelter, fire and weapons to dominate them. We were highly mobile and could soon construct boats and primitive bridges and slash convenient routes through dense vegetation. We learnt to cache food and to extend our range through the use of animal skins as clothes. Other mammals had the protection of size and strength, but they did not have guile, and only our low numbers at first delayed the extinctions that were to characterise the later part of the Pleistocene and early part of the Holocene.

The evidence lies in archaeological remains including cave paintings and petroglyphs (Fig. 1.12), in the huge quantities of mammal bones preserved in the wind-blown soils of Alaska and Siberia and in the spore and pollen changes seen in lake sediment cores from that time. Analysis of lake sediments is an important tool in understanding freshwater ecology (see Section 12.7). The cores show initially rich deposits of a very easily recognisable fungal spore, *Sporormiella*, which comes from a fungus specific to the dung of large wild herbivores (Fig. 1.13). Over a few hundred or thousand years, the spores in cores yet examined from North America, Madagascar and Australia became scarce whilst the amount of charcoal deposited in the sediment steadily increased. As the plant biomass of the grasslands was less grazed and more of it dried in the late season, grassland fires were fuelled. Perhaps also, we used fire to drive the last of a dwindling source of



Figure 1.12 Cave paintings, and petroglyphs cut into rock surfaces, of large mammals and hunting scenes abound from the late Pleistocene. This exquisite engraving of a sleeping antelope is one of around 15 000 from a small area at Tin Taghirt in the Tassili n'Ajjer mountains in southern Algeria. It was cut probably around 9–10 thousand years ago. (Reproduced with permission of Linus Wolf.)

prey into traps for slaughter. Traditional buffalo jumps were used in Canada until the sixteenth century CE.

About 9000 years ago, in the Middle East, Meso-America and China and not long afterwards on other continents, the hunter-gatherer humans realised that a more settled existence was possible, at first by attracting game towards their camps, or by locally planting seeds of favoured food plants that they had previously gathered. The details of how husbandry and agriculture began are inevitably few and vague, but there is no doubt that these were to be planet-changing moves. Domesticated stock is easy prey to wild predators, so removal of these became paramount. Cultivated fields contain attractive, nutritious food that must be protected from wild herbivores, which must be eliminated. Cultivation allows human populations to increase, for food supplies are greater and reliably harvested. It encourages societies to become more hierarchical, work to become more specialised and exploitation more efficient, for there are needs for labourers, supervisors and investors. Tillage disturbs soils and increases the loss of nutrients and other elements from them through exposure to rain (Fig. 1.14). The soils become less productive and cropping rates can be




Figure 1.13 Evidence of the effects of large herbivore populations comes from lake sediment cores, illustrated here for lakes in Indiana (USA; left), Madagascar (centre) and Australia (right) for periods corresponding to major loss of large herbivores. The numbers of spores of the dung fungus, *Sporormiella* (illustrated), are shown relative to the amounts of charcoal particles. Sediments have been dated using the carbon-14 method. Sometimes these dates are calibrated against tree-ring data, so that cal. ka BP means calibrated thousands of years before present. 'Present' is set at 1950 CE; yr BP means years before present, without cross calibration. The calibration generally makes only small differences. (Left, based on Burney *et al.* 2003. Reproduced with permission of AAAS. Centre, based on Gill *et al.* 2009. Reproduced with permission of National Academy of Sciences, USA. Right, based on Rule *et al.* 2012. Reproduced with permission of AAAS.)

maintained only by fertilisation; at first by the fresh silts that entered the floodplains where cultivation began in damp soils, but on the higher land only by adding animal manures, and later inorganic fertilisers. But agricultural systems do not have the many subtle ways that a biodiverse natural ecosystem calls on to retain and recycle nutrients. They are leaky and so for the freshwaters a process of nutrient pollution, or eutrophication inevitably began when agriculture developed.

Meanwhile, as populations increased through the advantages of permanent settlement, more land needed to be cleared. Agriculture, once begun, was unstoppable. It generated a need for more of itself and social structures that began to embrace larger and larger settlements. As the fields became more extensive, roads became necessary; as the proceeds built up, land became valuable, ownership became an issue; armies were born to defend the spoils. The hunting camp of the Mesolithic 9000 years ago became the



Figure 1.14 Natural vegetation conserves its nutrient supply, but agriculture replaces it with a leaky system where soil is exposed to rain, the annual crops do not have the retention mechanisms that perennial woodland has and where continual losses of nutrients must be replaced by fertiliser, which compounds the problem. Nutrient pollution (eutrophication) is inevitable whenever natural vegetation is disturbed.

city-state of the modern period. There were pressures to make agriculture produce more. Fertiliser was increasingly used but also chemicals that defended the crops and the domesticated stock from diseases that multiplied in plants and animals that were themselves no longer tempered by natural selection for survival, but under human selection for high productivity. Inevitably such substances, designed to kill, had effects on the remaining semi-natural lands as they spread as aerosols or even vapour and eventually reached the freshwaters. Some pesticides are now detectable even in seabirds in the remoter areas of Antarctica.

Land that was naturally wet and had borne the river floods was coveted for more agriculture, and drained. The propaganda grew that this was land that was inconveniently sometimes flooded and needed to be

defended by pumps and embankments, and displaced the understanding that it was the river bed, occupied for only part of the year but necessary for the river to be able to cope with varying flows. In some years the water supply failed for the storages had been destroyed. And so, farther upstream, reservoirs had to be created. Some were needed also to supply domestic water to where the cities, almost always clustered by the rivers, had burgeoned and polluted the water with sewage and other wastes. Reservoirs mean dams and dams block the passage of fish, some of which have life histories that require movements between the headwater rivers and the ocean. Reservoir management also disrupts natural flow patterns so that the life histories even of invertebrates are unable to cope. By the twenty-first century, with more and more human pressures, there is nothing ecologically pristine that is left, but about 25% of the land surfaces masquerade as ‘close to natural’ though they are far from it. The rest has been manifestly converted from biome to anthrome: human-dominated and managed systems. Geologists are formally considering naming the recent period as the Anthropocene, in which the surface of the planet is shaped to a considerable extent by human activity rather than solely by the natural forces of the past.

1.6 Politics and pollution

The citizens, as their cities expanded and gave the visual impression of self-containment, have become less and less aware of their dependence on natural processes. The composition of the atmosphere was of little concern as lifestyles became more isolated, and luxuries came to be regarded as necessities. A huge range of chemicals was manufactured to serve these purposes; there are probably at least as many ‘unnatural’ chemicals in the freshwater system as ‘natural’ ones but it is impossible to be sure. There may be as many as a million different substances, mostly organic and at very low concentrations, in unadulterated waters anyway.

Homo sapiens came to be seen as a heroic species conquering nature, improving its lot, the new king of the beasts. Political systems, with this idea at heart, have developed over the past few centuries and are inward-looking, with acquisition of wealth (so long as revolution by the workforce is avoided) their main aim. Their proponents have little knowledge of ecological science. Indeed they deride it because its findings are inconvenient to their self-serving aims. Energy has been generated on an unprecedented scale from the burning of fossil fuels, increasing the carbon dioxide concentration in the atmosphere from 190 ppm by volume or lower at the start of the Holocene to over 400 by the first few years of the twenty-first century (see Section 18.2). That such a gas retains heat radiation in the atmosphere and thence promotes higher temperatures, evaporation and heavier rainfall is trivialised. Propaganda that increasingly heavy flooding is ‘unprecedented’ might be true, but the reasons for it are not divine will, but simple physics. In tandem, the burning of coal and oil has acidified the atmosphere, reduced the pH of rain and caused serious problems for streams and lakes that have lost much of their diversity. And so a disrupted, poisoned landscape has a bloodstream that is itself sick, and indeed connects with an equally damaged ocean.

Until recently, the ocean, remote and largely unobserved, was regarded as unfettered wilderness. But it is not. Acidification from carbon dioxide absorption, overexploitation of invertebrates and fish, even of birds and mammals, pollution and loss of coastal ecosystems to development, have put paid to that. Without realising it, we face a complex of different threats (see Section 18.7). They consist of climate change, loss of much of our natural biodiversity, excessive consumption and indulgence in the developed world, and their consequences in the developing world. There is increasing poverty and unrest, many small wars, appalling air and water quality in many cities, poor health and overpopulation in the developing world as, under the economic influence of the West and of strangulating trade agreements and hard currency

loans, it is forced to repeat the West’s history of mistakes. And all of these problems, directly or indirectly are connected by the wastage of freshwater of sufficient quality.

Our future is very clouded. We have done less to mitigate the problems than to deride them with convenient propaganda. We spend large sums of money in meetings and conferences to make empty resolutions and minimise the problems. Political diplomacy froths over scientific reality. The only consolation is that science understands what must be done. Failure to act lies in the politics of a ruthless, self-indulgent invasive species that eternally schemes and deceives, covers up and lies. Eventually there must be a denouement, but it could take any of many forms. We can only hope that there will be enough left as a basis for repair when that time comes.

1.7 On the nature of textbooks

Textbooks on freshwater ecology are usually propaganda-ridden too. Most freshwater ecologists used to head for the hills in their research. They sought the least disturbed, ideally the pristine, environments where the results of evolutionary play are still intact and where the design of the environmental theatre has not been wrecked by ignorant demolition by ambitious people. As a result, we can draw pictures of intensely fascinating ecologies: the floodplain forests of the Amazon, with their seed-eating fish; the connections between wolves and bears in the functioning of river systems of the Pacific north-temperate zone; the division of the available food in the African Great Lakes among fish so specialised that some scrape the scales or bite the eyes of others (see Fig. 3.13). We can take you to lakes and wetlands in seemingly endless landscapes of forest and savannah, steppe and even desert. We can be more escapist than the glossiest of travel brochures (Fig. 1.15). But we mislead if you come away with the idea that a basically high-quality freshwater system is just sullied on a local scale. The truth is that a generally



Figure 1.15 The wilderness of Africa. Buffalo, egrets and a crocodile make a photogenic tableau at the edge of the Kazinga Channel in Uganda. But this is no wilderness. Not 500 m away is a fishing village.

deteriorating system still has a few relics of what it formerly was.

The bears and wolves have long gone in most places, as have many of the fish that were once common. Wild Atlantic salmon in Europe, for example, are now rare. Previous ecological textbooks, including earlier editions of this one, pretended that the world was intact, and tagged on human influences as secondary. This is no longer the case. The world is heavily damaged and more than half of its natural ecosystems are seriously degraded. An honest textbook must reflect this.

So a final difficulty is one of motivation, for both of us, reader and author. Ecologists do not particularly want to write as historians of what once was, but no longer is. They do not want to appear as doomsayers, perpetually depressed and backward-looking. Nor do they want to instil the idea that things are going so steeply downhill that nothing can be done. To undermine optimism and morale is a great disservice and self-fulfilling. Better to see things like this. We are studying perhaps the most interesting scene of the evolutionary play so far. People are immensely successful organisms, arch competitors, who have

already out-competed many other species and brought many others already to extinction. That milk is spilt.

Yet when one looks at the population fluctuations of an animal or plant, there are times when numbers (and thus impact) rise temporarily above the notional line of carrying capacity. For a very small part, perhaps only a few centuries, of our 100 000-year history we have been above this line. It has had immense effects, but we live on a planet that has its own mechanisms of stabilising its environment and may make its own adjustments, though these may be something of a shock to our eventual numbers. We may be able to avoid this, however, for we know a lot about how these systems work. We already have the capacity to start readjusting and re-designing so that we maintain ecological systems, which cannot be the same as those in the past, but which could still be interesting, pleasant and, above all, functioning to maintain an equable environment. It is here that freshwater, indeed all of ecological science, starts to involve the human social sciences, for much ecology is now connected with human societies. In the final analysis, our present difficulties, frustrations and

depressions can be a significant but reversible blip of human history not the start of an unending nightmare.

And against all these difficulties, there is one monumental bastion. Freshwater ecology is immensely interesting. I do not want to sound parochial about this. Anything is very interesting if you go deeply enough into it. But freshwaters are so much a part of our daily lives (go and turn the kitchen tap on) that they have a special place. One of my colleagues illustrated this by asking students to imagine somewhere where they would feel most at home. Almost invariably the scene involved water.

1.8 Further reading

G.E. Hutchinson's classic '*A Treatise on Limnology*' is published as four volumes, with enormous reference value though some sections are now out-of-date. There are many other texts on freshwater ecology, each with its own approach, and looking at several is always a good idea. Kalff (2003; a new edition is expected in 2018) and Lampert & Sommer (2007) have north-temperate emphases. Brock & Boulton (1999) covers Australasia. Davies & Day (1998) is a book on African freshwaters, and Talling &

Lemoalle (1999) cover tropical limnology. Bronmark & Hansson (2005) emphasises more the biology of organisms, whilst Burgis & Morris (2007) is a very readable account of lakes, Hynes (1966, 1979) of rivers and Worthington & Worthington (1933) a delightful and eminently readable account of east African waterways and ecology as they once were. Smol (2008) approaches the subject through what can be gleaned by a study of palaeolimnology. Mason (2002) is an introduction to the problems that all waters are now suffering from. The global situation is amply documented in the Millennium Ecosystem Assessment reports (www.millenniumassessment.org/en/index.aspx, accessed September 2016). Hutchinson (1965) introduces the analogy of the ecological theatre and the evolutionary play. There are some references to specialist topics in this overview. Regulation of the composition of the biosphere is interestingly covered in Lovelock (1979), Watson & Lenton (2011) and Tyrell (2013). It is a controversial area, but an exceptionally important one. There are several readable books on the mechanics of human evolution, but the meaning of it is much more profoundly covered in Rowlands' (2008) book, *The Philosopher and the Wolf*. The rise and implications of agriculture are well covered in Diamond (2007).