

# Subject Index

Page numbers with 'f' indicate figures. Page numbers with 't' indicate tables.

- Abiotic factors
  - in communities, 572
  - in CR habitats, 217
  - in harsh environments as, 616
  - in predator–prey cycles, 325
  - stressful, effects of, 419
- Above-ground NPP (ANPP), 658, 658f
- Abscission, 301, 361
- Abundance
  - alternative stable states of, 476
  - correlations with, 453–4
  - data, on species richness, 606
  - determination of, 454–5, 455f
  - elasticity analysis for, 129–31, 129f, 130f
  - extinction risk and, 502
  - index of, 108, 108f
  - key factor analysis of, 456–8, 456t, 457f, 459t
  - of keystone species, 553
  - lambda ( $\lambda$ ) contribution analysis of, 458, 459t, 460f
  - in metacommunity models, 601, 601f
  - metapopulations and, 194f, 195
  - multiple equilibria for, 474–5, 474f, 475f
  - outbreaks in, 474f, 475f
  - patterns, predation and, 285
  - rank–abundance diagrams, 530, 575–7, 576f
    - for metacommunity, 501f, 601
  - regulation of, 454–5, 455f, 477–8
  - stability of, 454–6, 454f
  - theories of, 454–5
  - time series of, 460–5, 462f–5f
- Abyssal zone, 359, 618, 635
- AC. *See* Adaptive capacity
- Acclimation, 45, 46, 69
- Acclimatisation, 45, 45f
- ACF. *See* Autocorrelation function analysis
- Acidification, of oceans, 707, 707f, 720–1, 722f, 723f
- Acid rain, 60–1, 60f, 681, 720, 721f, 722
- Acquired immunity, 385–6, 385f
- Activation energy, 98
- Active dispersal, 177–9, 177f, 178f
- Adaptation, 1–2
  - in foraging, food chain length and, 565, 565f
  - local, 3–5, 5f
- Adapting mosaic, 727, 727f
- Adaptive capacity (AC), 714
- Adenosine triphosphate (ATP), 100, 447
- Aerobic decomposition, 359
- Aerobic respiration, 90–1, 91f
- Age
  - of maturity, 220
  - reproductive allocation and, 218–20, 219f
  - reproductive value and, 209–10
- Age-specific fecundity schedules, 114
- Aggregated dispersal, 180–3, 181f–4f
- Aggregated distributions

- aggregation of risk and, 349, 349f
- interference and, 347–9, 348f
- interspecific competition and, 266–7, 267f
- of parasites in hosts, 381–2, 382f
- of parasitoids, 340f, 349–50, 349f
- in predation population dynamics, 339–41, 340f
- of predators, density of prey and, 349–50
- Aggregate ecosystem properties, 552
- Aggregation of risk
  - aggregated distributions and, 349, 349f
  - direct density dependence and, 350–1
  - for parasitoids, 355f
- Agriculture. *See also* Crops
  - conservation and, 610–11, 611f
  - mutualism in, 423–4
  - nitrogen in, 725–6
  - phosphorus in, 725–6
  - pollution from, stream invertebrates and, 29, 29f
- AITs. *See* Automatic isolating traits
- Algae
  - allelopathy of, 248
  - coexistence of, 258f
  - fungi and, mutualism of, 444–5, 445f, 446t
  - as modular organisms, 102
  - photosynthesis and, 69
  - salinity and, 57
  - specialisation of, 3
  - symbiosis of, with corals, 438f, 439–40
  - symbiosis of, with invertebrates, 438f, 439–40
- Alkaloids, 298
- Allee effect, 337, 338, 338f, 474, 476, 493
- Allelopathy, 240f, 248–9
- Allochthonous production, 652–4
- Allometric relationships
  - in life histories, 235–6, 235f, 236f
  - for mammals, 238–9, 239f
  - of metabolism, 97–8
  - in plants, 100, 100f
- Allopatric speciation, 8–14, 9f, 10f, 12f–13f
- Alpha ( $\alpha$ ) richness, 575, 592, 601
- Alternative stable states, of abundance, 476
- Ammonia, 60f
  - gut mutualism and, 432
  - nitrogen from, 95
  - in terrestrial biomes, 682
- Amoebic dysentery, 379
- Amphibians
  - biodiversity conservation of, 15
  - dispersal of, by sex, 185–6, 186f
  - extinction risk of, 503, 509, 510f
  - in extreme temperatures, 53f
  - humidity and, 54
  - as invasive species, 24f
  - migration of, 172
  - population size of, 709, 710f
  - species richness of, 615f, 633f
- Amphicarpy, 184
- AMs. *See* Arbuscular mycorrhizas
- Anaerobic decomposition, 359
- Anaerobic respiration, 90, 91f
- Animals
  - carbon : nitrogen ratio in, 93
  - defences of, 314–19, 314f, 315f, 317f–19f
  - as detritivores, 357–77
  - dormancy in, 111–12
  - migration of, 170
  - as modular organisms, 104
  - natural selection in, 2
  - nutritional contents of, 93–5
  - oxygen for, 90–1, 91f
  - resources of, 92–3
  - semelparous life cycle of, 110
  - sodium chloride for, 32
  - species distribution of, temperature and, 49–54, 50f–3f
  - in succession, 587–8
  - in tropical rainforest, 27
  - water for, 74
- Annual–biennials, 111
- Annual plants, 110–11
  - dormancy of, 114–15
- Anoxic habitats, 359
- ANPP. *See* Above-ground NPP
- Antagonistic coevolution, in plants, 394–5, 395f
- Antagonistic resources, 96–7
- Anthromes, 718–19, 719f
  - expansion of, 724
- Anthropocene, 708–9
- Ants
  - mutualism of
    - with aphids, 424, 424f
    - with butterflies, 425–6, 426f
    - with fungi, 426, 427–8, 427f
    - with plants, 422–5, 422f, 423f, 425f
  - species richness of, 609, 609f
- Aphids, 95
  - ants and, mutualism of, 424, 424f
  - defensive symbioses of, 438
  - dispersal of, polymorphism and, 185, 186f
  - IPM for, 485

- Aphids (*cont'd*)  
 plant defences against, 296, 302  
 as secondary pests, 480  
 viruses on plants from, 310, 310f
- Aposematism, 316
- Apparency theory, 298–9, 298f
- Apparent competition, 270–4, 271f–3f
- Apparent Trophic Cascade Hypothesis (ATCH), 550–1, 550f, 551f
- Aquatic biomes/communities/environments, 27. *See also*  
 Corals and coral reefs; Estuaries; Lakes; Ocean; Rivers;  
 Streams  
 carbon dioxide in, 78–9, 79f, 80f  
 clonal growth dispersal in, 179  
 detritivores in, 364, 364f  
 high temperatures in, 48  
 hypoxia in, 728, 728f  
 nutrient budgets in, 680f, 689–707  
 nutrient pollution in, 681–2, 682f  
 primary productivity in, 664–7, 664f–8f  
 regime shifts in, 569f  
 species richness by depth in, 635–6, 636f
- Arbuscular mycorrhizas (AMs), 441–2, 441f
- Areal argument, for self-thinning, 165–6
- Arthropods  
 augmentation of, 484  
 in extreme temperatures, 53f  
 humidity and, 54  
 as invasive species, 24f
- Assimilation efficiency, 672–3
- Assisted evolution, 723
- Asymmetric competition  
 and population size, 159, 159f  
 and size, 155–8, 156f–8f  
 and territoriality, 159–61, 160f, 161f
- Asymptotic stability, 552
- ATCH. *See* Apparent Trophic Cascade Hypothesis
- Atmosphere  
 continental shelves' nutrient budget and, 695  
 nutrient loss in, 684–5  
 phosphorus from, 683f
- ATP. *See* Adenosine triphosphate
- Augmentation, for biological pest control, 484
- Autochthonous production, 652–4  
 in estuaries, 665  
 in lakes, 665
- Autocorrelation function analysis (ACF), 461, 462f, 464f, 465
- Automatic isolating traits (AITs), 14
- Autotrophic respiration (RA), 649
- Autotrophs, 65  
 nutritional contents of, 95
- Bacteria, 136f. *See also* Nitrogen-fixing bacteria  
 allelopathy of, 248–9  
 bacteriophages and, coevolution of, 395  
 as biological pesticides, 484  
 carbon : nitrogen ratio in, 93  
 cockroaches and, 371–2  
 as decomposers, 357–77  
 effects on plants of, from herbivores, 310  
 at high temperatures, 48  
 for insect pest control, 438, 438f  
 insects and, symbiosis of, 436–7  
 interspecific competition of, 282, 282f  
 as microparasites, 379  
 as modular organisms, 102  
 oxygen and, 90  
 plastic pollution and, 731–2  
 in ruminant gut mutualisms, 434f  
 virulence of, 395, 395f
- Bacteriophages, 395
- Barnacles, 123, 134, 134f  
 birds and, food web trophic cascades of, 545, 545f  
 community-wide character displacement of, 532f  
 mussels and, 536  
 specialisation of, 3
- Basic reproductive number, 403  
 immunisation and, 406, 406f
- Batesian mimicry, 314f, 316
- Bats, 510–11  
 birds and, parallel evolution of, 18  
 pollination by, 429  
 seeds and, 588
- Bees  
 gut metagenome of, pesticides and, 436  
 plants and, mutualism of, 451, 452f  
 subpopulations of, 197, 198f
- Beetles, 133  
 character displacement of, 280, 280f  
 defensive symbioses of, 437–8  
 dormancy and, 113f  
 dung, 373–4, 373f, 374f  
 in four-trophic-level system, 545, 546f  
 fungi and, mutualism of, 425–6  
 importation of, for biological control, 483  
 at low temperatures, 45  
 plant defences against, 296–7  
 predation population dynamics of, 326, 326f  
 protozoan parasites and, 411f

- species richness of, 621, 621f
- temperature and, 40f, 41f
- Beta ( $\beta$ ) diversity, 592, 601, 612
- Beta ( $\beta$ ) richness, 575
- Bicarbonate, 79
- Biennial plants, dormancy of, 114–15
- Biliproteins, 69
- Biodiversity
  - conservation of, 15, 15f, 603–46
    - area selection for, 639–41, 640f, 641f
    - PAs for, 604–5, 604f
    - sustainability and, 642
  - food web stability and, 401, 559
  - humans and, 663, 663f
  - loss of, humans and, 709–11, 738
  - of plants, 267–8, 268f
  - species richness and, 603–5
  - of tropics, 53
  - Unified Neutral Theory of, 530
- Bioengineering, of crops
  - from CCMs, 83, 83f
  - for photoprotection, 70, 70f
- Biogeochemical cycles
  - global, 698–708
  - of matter flux, 679–80
- Biological control
  - desirable properties of agents, 356
  - generalist predators in, 337
  - and integrated pest management, 485
  - of myxamatosis, 393
  - of pests, 483–5, 483t
  - switching of agents, in horticulture, 289
  - of thistles, elasticity analysis of, 130
- Biological pesticides, 484
- Biomagnification, 480–1
- Biomass
  - of carnivores, 670, 670f
  - density and, 167–8
  - energy flux and, 648–9
  - NPP and, 655–6, 655f
  - of prey, 160f, 670, 670f
- Biomes. *See also* Aquatic biomes/communities/environments; Terrestrial biomes/communities/environments
  - climate change and, 718–19, 719f
- Bioremediation, 60
- Biotrophic parasites, 379
- Birds
  - barnacles and, food web trophic cascades of, 545, 545f
  - bats and, parallel evolution of, 18
  - biodiversity conservation of, 15
  - brood parasitism by, 381
  - climate variability hypothesis for, 54
  - clutch size of, 226
  - community size spectra of, 577, 577f
  - conservation of, 515, 516
  - Darwin's finches, 401, 401f, 535, 535f
    - allopatric speciation of, 9–11, 10f
  - as detritivores, 376–7
  - dispersal of, 170, 178, 178f
    - patchiness of, 180
    - by sex, 185–6, 186f
  - as endotherms, 42–3
  - extinction of, from parasites, 534–5, 535f
  - extinction risk of, 503
    - from invasive species, 509, 509f
  - extinction vortex of, 509–10, 511f
  - in extreme temperatures, 53f
  - foraging by, climate change and, 717–18, 717f
  - fundamental niche of, 38–9, 39f
    - as herbivores, 38–9, 39f
  - inbreeding depression of, 505, 505f
  - industrial melanism and, 7–8, 7f
  - interspecific competition and coexistence in, 244–5, 245f
  - as invasive species, 23, 24f
  - on islands, checkerboard distributions of, 532, 532f
  - life histories of, 232
  - LTRE for, 128, 128f
  - metabolic scaling of, 99
  - metapopulation capacity for, 204–5, 204f
  - migration of, 170–1, 173
    - climate change and, 716
  - nematodes and, 411–14, 412f, 413f
  - niches of, 34f
  - optimal foraging theory for, 289–92, 290f
  - parasite host tolerance and resistance of, 388, 389f
  - population cycles of, 466–7, 466f
  - population size of, 709, 710f
  - reproductive allocation in, 219f, 220
  - seeds and, 93
  - species distribution modelling for, 195, 196f
  - species richness of, 615f
  - territoriality of, 162
    - as true predators, 285
    - as unitary organisms, 102
- Births and birth rate
  - cohort generation time and, 123
  - counting, 108
  - density-dependent, abundance and, 455, 455f
  - of humans, 145, 146f
  - metapopulations and, 198

- Births and birth rate (*cont'd*)  
 of modular organisms, 106f  
 monitoring of, 114–21, 114f, 115t, 116t, 118f–20f  
 net recruitment curves for, 142–3, 142f, 143f  
 reproductive value and, 209
- Body size. *See* Size
- Boltzmann's constant, 98
- Boron, 87
- Bottom-up food webs, 548–9, 549f
- Boundaries  
 in communities, 581  
 planetary, 738–40, 741f, 741t
- Boundary lines, population, 164–5, 164f, 166f
- Bovine tuberculosis (BTB), 387, 387f
- Brood parasites, 381
- BTB. *See* Bovine tuberculosis
- Butterflies, 136f  
 agriculture and, 610, 611f  
 ants and, mutualism of, 425–6, 426f  
 chemical defences of, 314, 314f  
 crypsis of, 316  
 dispersal of, invasion, 190–1, 190f  
 extinction risk of, 508–9, 508f  
 inbreeding depression of, 504–5, 504f  
 metapopulations of, 198, 198f, 201f, 202–3, 203f  
 migration of, 170  
 reactive–diffusion modelling for, 193–4, 193f  
 resources of, 97
- C<sub>3</sub> pathway, 79–82, 81f, 85f
- C<sub>4</sub> pathway, 79–82, 81f, 85f
- Cacti  
 in deserts, 27  
 in extreme temperatures, 51
- Cadmium, 60
- Calcium, 87
- Calcium carbonate, 63
- CAM pathway, 82–3
- Canonical correspondence analysis (CCA), 38f
- Capacity-limited thermal tolerance theory, 48
- Carbohydrates  
 cold tolerance and, 46  
 defoliation and, 309  
 low temperatures and, 45  
 SCFAs and, 432
- Carbon  
 cycle, 705–7, 706f  
 decomposers and, 369–70, 370f  
 forests and, 685–6, 686f  
 in photosynthesis, 66  
 in plants, 95
- Carbon concentrating mechanisms (CCMs), 79–80, 83, 83f
- Carbon dioxide, 61–3, 61f, 77–87, 78f–86f  
 in aquatic biomes, 78–9, 79f  
 in the atmosphere, 706–7, 707f  
 changing concentrations of, 83–7, 84f, 85f  
 climate change from, 570–1, 571f  
 energy flux and, 678–9  
 enhancement of, harmful effects of, 86–7, 86f  
 GPP and, 652  
 as inhibitory resource, 97  
 in oceans, photosynthesis and, 63  
 photoautotrophs and, 65  
 photosynthesis and, 69  
 photosynthetic capacity and, 71  
 plant defences and, 306–7, 307f  
 radiative forcing and, 711–12, 712f  
 terrestrial biomes and, 682
- Carbonic acid, 683
- Carbon : nitrogen ratio, 93, 95  
 decomposers and, 370
- Carbon : nutrient balance hypothesis, 304
- Carboxylation, 73
- Carnivores  
 biomass of, 670, 670f  
 faeces of, detritivores and, 373  
 in four-trophic-level system, 545, 546f  
 invertebrates as, 364  
 nutritional contents of, 95
- Carotenoids, 69
- Carrying capacity, 141–2  
 consumer–resource models for interspecific competition  
 and, 255  
 for global human population, 144–8, 144f–6f  
 in population models, 149–50  
 RCPs and, 711–12, 712f
- Catastrophes, 59
- Catchment management, in wetlands, 729, 729f
- CCA. *See* Canonical correspondence analysis
- CCMs. *See* Carbon concentrating mechanisms
- CE. *See* Consumption efficiency
- Cellulose, 93, 95  
 decomposers and, 371  
 detritivores and, 371, 371f  
 gut mutualism and, 432
- CFCs. *See* Chlorofluorocarbons
- Chagas disease, 438
- Chain of extinctions, 510–11, 511f
- Chamaephytes, 29

- Chaos, in population models, 152–3
- Chaparral, 25f, 26
- Character displacement, 279–80, 279f, 280f, 531–2, 532f
- Checkerboard distributions, 532–3, 532f
- Chemical defences  
 of animals, 314–15, 314f, 315f  
 of plants, 295
- Chemoautotrophs, 65
- Chilling injury, 44
- Chlorofluorocarbons (CFCs), 62, 730
- Chlorophyll, 66–7, 69
- Chrysophytes, 69
- Climate change  
 biomes and, 718–19, 719f  
 energy flux and, 676–7, 677f  
 evolution and, 19–22, 19f–22f  
 extinction risk and, 508–9, 508f  
 fish and, 718, 719f  
 food webs and, 722–3  
 foraging by birds and, 717–18, 717f  
 forests and, 21f, 713–15, 714f, 715f  
 GPP and, 652  
 groundwater and, 701  
 gut metagenome and, 434  
 hydrological cycle and, 701, 701f  
 invasive plants and, 736–8, 736f  
 LUC and, 724–9, 724f  
 migration of birds and, 716, 716f  
 ocean acidification and, 707, 707f  
 predictions for, 42  
 reactive–diffusion modelling and, 193–4, 193f  
 as regime shift, 570–1, 571f  
 in tropics, 53
- Climate matching, 35–6
- Climate variability hypothesis, 52–4
- Climax, succession to, 588–9
- Clonal growth dispersal, in aquatic biomes, 179
- Clutch size, 222f–5f, 223–6
- Coarse-grained environment, 180, 181f
- Coarse particulate organic matter (CPOM), 365f
- Cockroaches  
 bacteria and, 371–2  
 as invasive species, 23  
 protozoans and, 371–2
- Coevolution  
 of bacteria and bacteriophage, 395  
 of parasites and hosts, 392–5, 394f, 395f  
 of predator–prey, 292, 295
- Coexistence  
 of algae, 258f  
 exploiter-mediated, 535–7, 540  
 interspecific competition and  
 in birds, 244–5, 245f  
 in diatoms, 244, 245f  
 in fish, 243–4, 243f  
 niche overlap and, 261–2, 261f  
 niche similarity and, 262–3, 263f  
 with niche differentiation, 262  
 without niche differentiation, 247–8, 248f  
 of plankton, 247–8, 248f  
 supply points of resources and, 255
- Cohort generation time, 123
- Cohort life tables, 114–17, 114f, 115f, 116f, 123, 123f
- Coinfection, 392, 392f, 393f
- Cold tolerance, 45–7, 46f, 47f
- Cole’s paradox, 221
- Collector–filterers, 364
- Collectors, 364
- Colonisation  
 competition and, trade-offs between, 586–7, 599, 600f  
 competition and, in succession, 582  
 of fungi, in dead wood, 366–7, 367f  
 gaps in  
 in mangroves, 596, 596f, 596t  
 of mussels, 595f, 596  
 interspecific competition and, 263–9, 264f–9f  
 on islands, 622, 624, 625f  
 in patches, 595f  
 of pests, 484
- Commensalism  
 competition and, trade-offs between, 418  
 of corals, 416  
 defined, 416  
 evolution of, 419–20  
 of fish, 416–17  
 in habitats, 417  
 with herbivores and grasses, 417–18  
 intercropping and, 418–19
- Communities. *See also* Aquatic biomes/communities/  
 environments; Metacommunity; Succession; Terrestrial  
 biomes/communities/environments  
 boundaries in, 581  
 competition in, 523–34, 523f, 531f–3f  
 demonstrable effects of, 524, 525f  
 literature reviews on, 524  
 niche complementarity, 525–8, 526f–8f  
 niche differentiation in, 525–31, 529f, 531f  
 null model analysis for, 528–33, 529f, 534  
 composition of, 572–602  
 disturbance in, 593–4, 594f

- Communities. *See also* Aquatic biomes/communities/  
environments; Metacommunity; Succession; Terrestrial  
biomes/communities/environments (*cont'd*)  
diversity in, 574–5, 575f  
diversity of, of fish, 597  
environment and, 24–9, 25f, 28f, 29f  
food web trophic cascade in, 549  
gaps in, 592–7, 595f, 595t, 596f, 596t  
habitat hierarchy of, 572–3, 573f  
levels of, 573–4  
life form spectra of, 27–9, 28f  
ordination of, 578–81, 580f  
rules for, 572–3  
size spectra for, 577–8, 577f, 578f  
in space, 578–81  
spatiotemporal context of, 592–602  
species richness in, 574–5, 574f  
in time, 581–91
- Community modules  
defined, 522  
facilitation in, 540–1, 543f  
omnivores in, 540  
plurality in, 540–3  
predation in, 534–40, 534f–9f  
stochastic models for, 541–3  
stress-gradient hypothesis for, 540–1, 542f
- Community-wide character displacement, 531–2, 532f
- Compartmentalisation, of food webs, 561,  
562f, 563f
- Competition. *See also* Interspecific competition; Intraspecific  
competition  
apparent, 270–4, 271f–3f  
asymmetric  
for population size, 159, 159f  
by size, 155–8, 156f–8f  
for territoriality, 159–61, 160f, 161f  
coefficient, 249  
colonisation and  
succession and, 582  
trade-offs between, 586–7, 599, 600f  
commensalism and, trade-offs with, 418  
in communities, 523–34, 523f, 531f–3f  
demonstrable effects of, 524, 525f  
literature reviews on, 524  
niche complementarity, 525–8, 526f–8f  
niche differentiation in, 525–31, 529f, 531f  
null model analysis for, 528–33, 529f, 534  
contest, 140–1  
between decomposers and plants, 370–1  
defined, 132  
environmental conditions for, 525  
exact compensation and, 140–1  
exploitation and, 132–3, 132f, 133f  
facilitation to, shift from, 419, 420f  
in food webs, 557  
guilds and, 526–7, 529–30, 529f  
interference and, 133, 134f  
intransitive, 533–4  
of modular organisms, 139, 139f  
neutral, 262  
non-kin, 183, 183f  
of plants  
herbivores and, 308, 309f  
parasites and, in salt marshes, 402, 402f  
in population models, 149, 150–2, 151f, 152f  
pre-emptive, 263–9  
for resources, of parasite host, 388–92, 389f–92f  
of roots, 158, 158f
- Competitive exclusion principle  
Lotka–Volterra model and, 240, 252, 255–6  
productivity and, 609  
ZNGI and, 254, 254f
- Competitive strategy, 232
- Complementarity, of niches, in community  
competition, 525–8
- Complementarity effects  
food security and, 559, 559f  
of niche differentiation in food webs, 557  
species richness and, 662–3, 662f
- Complementarity selection, 640–1, 641f
- Complementary resources, 96
- Complex dynamics, in consumer–resource models for  
interspecific competition, 255–7, 256f
- Conservation, 501–21. *See also* Extinction  
agriculture and, 610–11, 611f  
of biodiversity, 15, 15f, 603–46  
area selection for, 639–41, 640f, 641f  
PAs for, 604–5, 604f  
sustainability and, 642  
decision analysis for, 518–21, 519f, 520f  
defined, 501  
of dung beetles, 374, 374f  
ecotypes for, 4, 4f  
life history fast–slow continuum and, 230–1, 231f  
of metapopulations, 515–18, 516f–18f  
of migratory species, 172–5, 173f, 174f  
optimal sowing rates for, 138, 138f  
successional mosaics for, 598, 598f  
value of, 502  
of water, photosynthesis and, 74–5
- Constant final yield, 135, 137f, 388–90, 390f
- Constitutive defences, 294–5

- Consumer–resource models
  - of interspecific competition, 253–61, 254f, 256f, 258f–60f
  - of predation population dynamics, 320–5, 322f, 323f
- Consumption efficiency (CE), 672, 672f
- Contact rate, 396–7
- Contemporary reproductive output, 209
- Contest competition, 140–1
- Continental drift, 16–17, 16f
- Continental shelves, nutrient budgets on, 693–5, 695t
- Continuous breeding, logistic equation for, 154–5, 155f
- Continuous redistribution, of parasitoids, 351
- Copper, 87
- Coprophagy, 433
- Corals and coral reefs
  - algae and, symbiosis of, 438f, 439–40
  - allelopathy of, 249f
  - assisted evolution for, 723
  - bleaching of, 440
  - calcium carbonate for, 63
  - commensalism of, 416
  - extinction risk of, 503
  - fish diversity in, 597
  - fungi and, mutualism of, 426–7
  - as modular organisms, 102
  - NPP of, 649t
  - PAs for, 604f
  - plastic pollution on, 731–2, 732f
  - regime shifts in, 569, 570f
  - tsunamis and, 58
- Cost of reproduction (CR)
  - clutch size and, 224–5
  - habitats and
    - age of maturity and, 220
    - comparative classification of, 216–17, 218f
    - reproductive investment and, 218
  - life history trade-offs with, 210f, 213
- Coupled oscillations
  - of parasitoids, 286f
  - in predation population dynamics, 286f, 324
- Cover crops, 55, 55f
- COVID-19, 378, 384
- C POM. *See* Coarse particulate organic matter
- CR. *See* Cost of reproduction
- Critically endangered, 503, 711
- Critical population size, 403–4
- Crops
  - bioengineering of
    - from CCMs, 83
    - photoprotection for, 70, 70f
  - cold tolerance of, 47, 47f
  - earthworms and, 362–3, 363f
  - in extreme temperatures, 53
  - GM of, 481, 482f, 484, 484f
  - intercropping, 277, 277f
    - commensalism and, 418–19
  - macroparasites in, 381, 408, 409f
  - NPP of, 649t
  - optimal sowing rates for, 138, 138f
  - parasites of, 378
  - soil pH and, 55, 55f
  - temperature and, 55, 55f
- Crowding
  - density and, 136–9
  - discrete breeding seasons and, 150
  - population growth and, 471–2
  - predation population dynamics and, 334
- Cryoprotective dehydration, 45
- Crypsis, 316
- Cryptophytes, 29, 69, 71f
- Cultural services, 58
- Cyanide, 295
- Cyanogenic glycosides, 298
- Cyclone Hugo, 58–9
- Dark diversity, 234–5, 234f
- Darwin, Charles, 1–2, 2f, 93, 429, 536
- Darwin's finches, 401, 401f, 535, 535f
  - allopatric speciation of, 9–11, 10f
- DD. *See* Dominance decay
- DDT. *See* Dichlorodiphenyl-trichloroethane
- Dead bodies
  - decomposition of, 358, 375, 375f, 376t
  - detritivores and, 375–6
  - microplastics in, 732
- Dead-end hosts, 383
- Dead zones, 681–2, 682f
- Dear enemy hypothesis, 161, 161f
- Deaths
  - counting, 108
  - density-dependent, abundance and, 455, 455f
  - of humans, 145, 146f
  - metapopulations and, 198
  - of modular organisms, 106f
  - monitoring of, 114–21, 114f, 115t, 116t, 118f–20f
  - net recruitment curves for, 142–3, 142f, 143f
  - of plants, defoliation and, 307
  - of unitary organisms, 102
- Decision support systems (DSSs), 485, 486f
- Decomposition and decomposers, 357–77
  - aerobic, 359
  - anaerobic, 359
  - of dead bodies, 358

- Decomposition and decomposers (*cont'd*)
- insects as, 375, 375f, 376t
  - of dead wood, 366–7, 367f
  - defined, 357–8
  - domestic processes and, 359
  - ecological stoichiometry of, 369–71, 370f
  - energy flux and, 674–6
  - of faeces, 358
  - food webs of, 357, 362f
  - industrial processes and, 359
  - isopods for, 372–3, 372f
  - of mammal carcasses, 367
  - of plant litter, home field advantage for, 367–8, 368f, 369f
  - plants and, 370–1
  - resources of, 357
  - in salt marshes, 365–6, 366f
  - size of, 361, 362f
  - in streams, 365
  - succession of, 360–1, 360f
- Defences
- of animals, 314–19, 314f, 315f, 317f–19f
  - of parasite hosts, 384–8, 385f–7f
  - of plants
    - against herbivores, 294–307, 296f–307f
    - apparency theory and, 298–9, 298f
    - carbon dioxide and, 306–7, 307f
    - fertilisers and, 305, 305f
    - of roots, 300, 301f
    - of shoots, 300, 301f
    - stress and, 305–6, 306f
    - timing of, 299–303, 299f–303f
- Defensive symbioses, 437–8
- Defoliation, by herbivores, 307–8, 309, 309f
- Deforestation
- extinction risk of, 507, 507f
  - nutrient flux and, 688–9, 689f
  - zero policy for, 720
- Degradation, of habitat, 507–8
- Delayed density dependence
- for parasitoids, 323f
  - of predation population dynamics, 323, 323f
  - satiation and, 329–30, 329f
- Demographic transition, 145
- Demographic uncertainty, of extinction, 503
- Density
- biomass and, 167–8
  - crowding and, 136–9
  - and emigration, 183
  - of prey, predator aggregate distributions and, 349–50
  - self-thinning and, 161–8, 163f, 164f, 166f–8f
- Density-dependence, weather and, 465
- Density-dependent birth, 455, 455f
- Density-dependent death, 455, 455f
- Density-dependent dispersal, 181–3, 181f–4f, 390f
- Density-dependent emigration, 184f, 186
- Density-dependent growth, 135, 136f
- Density-dependent transmission, 396–7, 397f
- Density management diagrams (DMDs), 168–9, 169f
- Depensation, in harvest management, 493–5, 494f
- Deserts, 25f, 26–7, 111, 649t
- Desiccation, 47
- Destruction, of habitat, 507
- Deterministic chaos, 152
- Detritivores, 357–77, 362f–5f
- in aquatic biomes, 364, 364f
  - ATCH for, 550, 550f
  - carnivore faeces and, 373
  - cellulose and, 371, 371f
  - dead bodies and, 375–6
  - ecological stoichiometry of, 369–71, 370f
  - food chains of, 550–1
  - herbivore faeces and, 373–4, 373f, 374f
  - invertebrates as, 375–6
  - invertebrates faeces and, 372–3, 372f
  - mutualism of, 371–2
  - NPP and, 675, 675f
  - parasites as, 379
  - resources and, 371–7, 371f–5f, 376t
  - in streams, 364–5, 365f
  - in terrestrial biomes, 363–4
  - vertebrates as, 376–7
- Development
- life histories and, 207–8
  - temperature and, 39–41, 41f
- DI. *See* Dissolved inorganic
- Diapause, 111–12
- Diatoms, 244, 245f, 265–6, 266f
- Dichlorodiphenyl-trichloroethane (DDT), 484, 731
- Diet width, 289–92
- Differential resource utilisation, 244–5
- Dilution effect, 400–1, 400f
- DIN. *See* Dissolved inorganic nitrogen
- Dinosaurs, 118
- DIP. *See* Dissolved inorganic phosphorus
- Direct density dependence, of predation population dynamics, 323–4
- Direct life cycles, of parasites, 379
- Direct transmission, 396, 403–4
- Disasters, 58–9
- Disease. *See also* Infections
- extinction risk of, 509, 510f
  - immunisation for, 406–7, 406f

- from invasive species, 24f
- parasite hosts diversity and, 399
- superspreaders of, 398, 398f
- Dispersal. *See also* Dispersion; Invasion
  - active, 177–9, 177f, 178f
  - active–passive continuum of, 175
  - defined, 170
  - demographic significance of, 186–95
  - density-dependent, 181–3, 181f–4f, 390f
  - eccentric, 190–1, 190f
  - ESS for, 183
  - by inbreeding and outbreeding, 182, 182f
  - in metacommunity, 601, 601f
  - metapopulations and, 192–205, 194f, 196f, 198f–201f, 203f, 204f
  - modes of, 175–6, 175f
  - non-kin competition and, 183, 183f
  - passive, 175–9, 178f
  - patchiness of, metapopulations and, 192–5
  - philopatry and, 181
  - polymorphism, 184–5, 184f, 186f
  - population cycles and, 466
  - of seeds, mutualism of, 428–9
  - by sex, 185–6, 186f
  - of single populations, 186–95, 187f–91f
  - turnover, 180
  - by wind, 176–7
- Dispersion
  - aggregated, 180–3, 181f–4f
  - patterns of, 179–83, 180f–4f
  - random, 180, 180f
  - regular, 180, 180f
- Dissolved inorganic (DI), 727
- Dissolved inorganic nitrogen (DIN), 727–8, 727f
- Dissolved inorganic phosphorus (DIP), 727–8, 727f
- Dissolved organic (DO), 727
- Dissolved organic carbon (DOC), 722
- Dissolved organic matter (DOM), 365f
- Dissolved organic nitrogen (DON), 727, 727f
- Dissolved organic phosphorus (DOP), 727, 727f
- Disturbance
  - in communities, 593–4, 594f
  - food chain length and, 565–6, 567f
  - of food webs, 552, 557
  - of habitat, 508
    - by grazing, 537
  - species richness and, 608
- Diversity. *See also* Beta diversity; Biodiversity; Gamma diversity; Zeta diversity
  - in communities, 574–5, 575f
  - of fish, 597
    - of habitat, island biogeography and, 621
    - of parasite hosts, disease and, 399
- DMDs. *See* Density management diagrams
- DO. *See* Dissolved organic
- DOC. *See* Dissolved organic carbon
- DOM. *See* Dissolved organic matter
- Domestic processes, decomposition and, 359
- Dominance decay (DD), 576, 576f
- DON. *See* Dissolved organic nitrogen
- Donor control, 357, 548
- DOP. *See* Dissolved organic phosphorus
- Dormancy
  - of animals, 111–12
  - of plants, 112–14, 113f
  - of seeds, 112–14, 113f
- Dryfall, 683–4, 684f
- DSSs. *See* Decision support systems
- Dung beetles, 373–4, 373f, 374f
- Dutch elm disease, 411, 411f
- Dynamic pool models, for harvest management, 495–6, 496f
- Dynamic thinning lines, 162–3, 162f
  
- Earthworms, 54, 362–3, 363f, 364
- Ebola, 404, 405f, 406
- Eccentric dispersers, 190–1, 190f
- ECMs. *See* Ectomycorrhizas
- Ecological Society of America, 477
- Ecological stoichiometry, 95, 369–71, 370f, 675
- Economically optimum yield (EOY), 492, 492f
- Economic injury level (EIL), 478–9, 478f, 485
- Economic threshold (ET), 478–9, 485
- Ecosystem engineers, 417
- Ecosystem Exploitation Hypothesis (EEH), 550
- Ecosystem respiration (RE), 665, 665f, 676, 676f
- Ecosystem services, 58, 502, 648
- Ecosystem size, food chain length and, 564, 568, 568f
- Ecotypes, 2–5
  - for conservation, 4, 4f
  - polymorphism of, 5–8, 6f
- Ectomycorrhizas (ECMs), 442
- Ectotherms, 41–4, 43f, 52, 53f
- EEH. *See* Ecosystem Exploitation Hypothesis
- EESC. *See* Equivalent effective stratospheric chlorine
- Effective population size, 503–4
- EIL. *See* Economic injury level
- Elasticity analysis, 129–31, 129f, 130f, 458
- Electivities, 529–30
- Elephantiasis, 378
- Elevation, as species richness gradient, 633–6, 634f–6f
- El Niño–Southern Oscillation (ENSO), 49, 50f, 676–7

- Eltonian pyramids, 669
- Emigration, 183
- deaths and, 108
  - density-dependent, 184f, 186
  - in metapopulations, 192, 198
- Endangered species, 503, 711
- Endemism
- climate change and, 20
  - conservation and, 15, 15f
  - in fish, 13
  - on islands, 18, 627
  - of parasites, 403–11
- Endosymbionts, 437–40, 437f, 438f
- Endotherms, 41–4, 43f, 52, 53f
- Enemy-free space, of prey, 607
- Enemy interaction theory, 591
- Energy
- endotherms and, 43–4
  - flux, 647–77
    - biomass and, 648–9
    - carbon dioxide and, 678–9
    - climate change and, 676–7, 677f
    - decomposers and, 674–6
    - ENSO and, 676–7
    - grazers and, 674–6
    - nutrient flux and, 678–9, 679f
    - primary productivity and, 649–56, 649t, 650f–5f
    - productivity and, 648
    - through food webs, 671–2
    - trophic compartments for, 668–71, 671f
  - from foraging, 289–92
  - species richness and, 613–16, 614f–16f
- Enforced dormancy, 113, 113f
- ENSO. *See* El Niño–Southern Oscillation
- Environment. *See also* Aquatic biomes/communities/  
environments; Terrestrial biomes/communities
- community and, 24–9, 25f, 28f, 29f
  - conditions of
    - for competition, 525
    - niches, 33–9, 34f–9f
    - species and, 32–64
  - evolution and, 1–31
  - fluctuations of
    - harvest management and, 495
    - interspecific competition and, 265–6, 266f
- Environmental space, 169
- Environmental uncertainty, of extinction, 503
- Enzymes, in insects, 95
- EOY. *See* Economically optimum yield
- Ephemerals, 111
- Epidemic curve, 404, 405f, 406
- Equilibrium theory, of island biogeography, 619–31, 620f–31f
- Equitability, 574–5
- Equivalent effective stratospheric chlorine (EESC), 730, 730f
- Ericoid mycorrhizas, 442–3
- ESS. *See* Evolutionarily stable strategy
- Essential resources, 96
- Estuaries, 654
- anoxic habitats in, 359
  - autochthonous production in, 665
  - minerals in, 27
  - nutrient flux in, 692–93, 692f, 693f
- ET. *See* Economic threshold
- Eutrophication, 728, 728f
- Evenness, 574–5
- Evolution. *See also* Coevolution
- assisted, 723
  - climate change and, 19–22, 19f–22f
  - of commensalism, 419–20
  - environment and, 1–31
  - interspecific competition and, 247, 278–83, 278f–83f
  - on islands, 18–19, 18f
  - life histories and, 206–39
  - parallel, 17–18
  - of virulence, 392–4
- Evolutionarily stable strategy (ESS), 183
- Evolutionary time, 618
- Exactly compensating density dependence, 135, 140–1
- Experimental evolution, 395
- Exploitation, 132–3, 132f, 133f. *See also* Overexploitation
- EEH for, 550
  - of groundwater, 701
  - by humans, destabilising Allee effects and, 338
  - interference and, 248–9, 249f
- Exploiter-mediated coexistence, 535–7, 540
- Extinction
- causes of, 505–11, 506f–11f
  - chain of, 510–11, 511f
  - on islands, 620–1
  - LTRE for, 128, 128f
  - metapopulations and, 197–203
  - modern and historical rates of, 502
  - MVPs and, 511–12
  - from overexploitation, 505–7, 506f
  - from parasites, 534–5, 535f
  - PPMs for, 513
  - PVAs for, 512–14, 512f
  - risk of, 502–5, 503f–5f

- of climate change, 508–9, 508f
  - of disease, 509, 510f
  - habitat and, humans and, 507–8, 507f
  - invasive species as, 509, 509f
  - vortex, 509–10, 511f
- Extreme events, 58–9
- Extreme temperature, 39
- FACE. *See* Free air carbon dioxide enrichment facilities
- Facilitation, 416–51. *See also* Commensalism; Mutualism
  - in community modules, 540–1, 543
  - to competition, 419, 420f
  - species richness and, 608
  - in succession, 587, 590–1
- Facultative diapause, 112
- Facultative mutualism, 371
- Faeces
  - of carnivores, detritivores and, 373
  - decomposition of, 358
  - of herbivores, detritivores and, 373
  - of invertebrates, detritivores and, 372–3, 372f
  - microplastics in, 732
- FAO. *See* Food and Agricultural Organization
- Fast–slow continuum
  - for life histories, 227–31, 229f–31f
  - life histories and, for mammals, 238–9
- Fecundity
  - intraspecific competition and, 135, 136f, 140
  - of mammals, predator intimidation and, 318–19, 319f
  - of parasite hosts, 399–401
  - of plants, herbivores and, 311–12, 311f
  - schedules, 117
    - age-specific, 114
    - for variables, 122–3, 125f
  - in Y model for life history trade-offs, 211f
- Feeding commensalisms, 417–18
- Fertilisers
  - grasshoppers and, 317
  - IFS and, 485–6
  - plant defences and, 305, 305f
- Field capacity, 76, 77f
- Fine particulate organic matter (FPOM), 365f
- Fire
  - in forests, 715, 715f
  - high temperatures of, 48
- First come best served, in interspecific competition, 265
- Fish
  - acid rain and, 60, 722
  - age of maturity of, 220
  - allopatric speciation of, 12–13, 13f, 14
  - climate change and, 42, 718, 719f
  - commensalism of, 416–17
  - community diversity of, 597
  - detritivores and, 376
  - dispersal of, by sex, 185–6, 186f
  - endemism in, 13
  - in extreme temperatures, 52f
  - in food chains, 488–9
  - food switching by, 289f
  - foraging in presence of predators, 293–4, 293f
  - fundamental niche of, 244
  - harvest management of, 490–2, 491f, 492f, 495–9, 496f–500f
  - interspecific competition in
    - coexistence and, 243–4, 243f, 265
  - as invasive species, 23, 24, 24f
  - in lakes, acid rain and, 61
  - microplastics and, 732
  - migration of, 172
  - mutualism of, 421, 421f
  - in NAO, 52f
  - niches of, 36
    - complementarity of, 525–7, 526f
    - ordination for, 38, 38f
  - ocean warming and, 718, 719f
  - optimal foraging theory for, 290f, 291
  - overexploitation of, 733–6, 734f, 735f
  - polymorphism in, 14
  - realised niche of, 244
  - reproductive allocation in, predators and, 220, 221f
  - specialisation of, 3
  - species richness of, 559, 559f, 609, 609f
  - stream nutrient spiraling and, 690
  - temperature and, 40f, 42f
- Fitness
  - contours, life history trade-offs and, 216, 217f
  - of host, symbiosis and, 421–2
  - number of offspring life history trade-offs and, 213–14, 214f
  - of parasite hosts, 388, 389f
- Fixed escapement harvesting, 490–2, 491f, 492f
- Fixed harvesting rate, 488f, 489
- Fixed proportion harvesting, 490–2, 491f, 492f
- Floaters, in territoriality, 159, 160–1
- Flood risk, 701
- Food
  - consumers and, match diversity of, 30
  - donor controlled, 357, 548
  - of moths, 54
  - parasites and, 414–15, 415f
  - population cycles and, 466
  - predation and, 317, 317f

- Food (*cont'd*)  
 of predators, preferences of, 286–8, 287f  
 quality of, predation population dynamics and, 330, 330f, 331t  
 switching of  
 of predators, 288–9, 289f  
 versus optimal foraging theory, 292  
 water from, 74
- Food and Agricultural Organization (FAO), 171, 171f
- Food chains, 522, 544–51  
 of detritivores, 550–1  
 fish in, 488–9  
 length of, 564–6, 564f–7f  
 pesticide biomagnification in, 480–1
- Food webs, 464f, 522–3  
 acid rain and, 61  
 aggregate properties of, 557–9  
 bottom-up, 548–9, 549f  
 climate change and, 722–3  
 compartmentalisation of, 561, 562f, 563f  
 competition in, 557  
 of decomposers, 357, 362f  
 energy flux through, 671–2  
 keystone species in, 553–4, 554f, 555f  
 parasites in, 567, 568f  
 population stability in, 555–6  
 productivity of, 551, 552, 557  
 regime shifts in, 568–71, 568f–571f  
 robustness of, 552, 559–60, 560f  
 parasites in, 567, 568f  
 stability of, 552–7  
 biodiversity and, 401  
 complementarity effects and, 559, 559f  
 from fish, 559, 559f  
 top-down, 548–9, 549f  
 trophic cascades in, 544–51, 545f–51f  
 in mesopredator release, 547, 547f  
 trophic loops in, 561–4, 563f
- Foraging and foragers  
 adaptation in, food chain length and, 565, 565f  
 by birds, climate change and, 717–18, 717f  
 diet width for, 289–92  
 energy from, 289–92  
 of grasslands, 310  
 as handlers, 290–1  
 optimal foraging theory of  
 for plants, 343–5, 344f  
 for predators, 289–92, 290f  
 versus food switching, 292  
 in patchiness of environment, 339–49, 340f–2f, 344f–6f, 347t, 348f, 349f  
 in presence of predators, 292–4, 293f  
 roots as, 76–7, 78f, 87–8  
 as searchers, 290  
 widths and composition of diets of, 285–94
- Forests. *See also* Deforestation; Mangroves; Temperate forests;  
 Tropical rainforest  
 acid rain and, 60  
 carbon and, 685–6, 686f  
 carbon dioxide in, 78, 78f  
 climate change and, 21f, 713–15, 714f, 715f  
 at extreme temperatures, 52f  
 fires in, 715, 715f  
 management of, global warming and, 686–8, 687t  
 nitrogen in, 726, 726f  
 NPP in, 686, 686f  
 phosphorus in, 726, 726f  
 stability of, 21f  
 in wetlands, seed banks for, 112
- Founders  
 evolution of tolerance and, 59  
 gaps and, 596–7
- Four-trophic-level systems, 545–6, 546f
- FPOM. *See* Fine particulate organic matter
- Free air carbon dioxide enrichment facilities (FACE), 84–5, 84f, 85f
- Frequency-dependent predation, 538–9, 538f, 539f
- Frequency-dependent transmission, 397, 397f, 404
- Frogs  
 migration of, 172  
 population size of, 108
- Fundamental net per capital rate of increase, 122
- Fundamental net reproductive rate (R), 122, 125
- Fundamental niche, 36, 38–9, 39f, 244
- Fungi, 140f  
 algae and, mutualism of, 444–5, 445f, 446t  
 allelopathy of, 248–9  
 ants and, mutualism of, 426, 427–8, 427f  
 beetles and, mutualism of, 425–6  
 carbon : nitrogen ratio in, 93  
 colonisation of, in dead wood, 366–7, 367f  
 corals and, mutualism of, 426–7  
 as decomposers, 357–77  
 at high temperatures, 48  
 local adaptation of, 4  
 as macroparasites, 380–1  
 as microparasites, 379  
 plants and, 36  
 herbivores and, 310

- mutualism of, 407
- roots and, 88–90
  - mutualism of, 440–3, 441f, 441t
  - virulence of, 395f
- Gamma ( $\gamma$ ) diversity, 575, 592, 612, 613f
- Gaps
  - in colonisation
    - in mangroves, 596, 596f, 596t
    - of mussels, 595f, 596
  - in communities, 592–7, 595f, 595t, 596f, 596t
- GCMs. *See* Global climate models
- Generation lengths, 121–7
- Genet, 104, 106f
- Genetic modification (GM), of crops, 481, 482f, 484, 484f
- Genetic polymorphism. *See* Polymorphism
- Genetic variation, as extinction risk, 503–4
- Genotypes, 212–13, 296
- Geographic space, 169
- Geometric series (GS), 576, 576f
- Geophytes, 29
- GFCL. *See* Gross percentage forest cover loss
- GHG. *See* Greenhouse gas
- Gini coefficient, 157
- Giving-up times, 341
- Global climate models (GCMs), 711
- Global human population, carrying capacity for, 144–8, 144f–6f
- Global orchestration (GO), 727, 727f
- Global stability, 552
- Global warming, 22, 63–4, 63f, 711–20
  - coral bleaching and, 440
  - forest management and, 686–8, 687t
  - iron in oceans and, 698
  - permafrost methane anaerobic respiration and, 91–2, 92f
- Glucosinolate, 296, 297f, 298, 302, 304f
- GM. *See* Genetic modification
- GO. *See* Global orchestration
- GPP. *See* Gross primary productivity
- Grasses, 135
  - grasshoppers and, 132
  - herbivores and, commensalism with, 417–18
  - mast years of, 328, 328f
  - modular populations of, 104
  - niche divergence in, 281, 281f
  - salinity and, 57
  - species distribution modelling for, 195
  - tolerance of, 308
  - trade-offs in, 257, 259f
  - in tundra, 25
- Grasshoppers, 22f, 111, 132, 317
- Grasslands, 25f, 26
  - colonisation gaps in, 596
  - foraging of, 310
  - invasive species and, 23
  - NPP of, 649t
  - species richness in, 558f
  - succession in, 585, 585f
- Grazers and grazing
  - by detritivores, 361
  - energy flux and, 674–6
  - by herbivores, 285
  - as keystone species, 553
  - plant species richness and, 536–8, 536f–8f
  - as predators, 284
  - primary productivity and, 670–1
  - resources of, 92
- Grazer–scrapers, 364
- Greenhouse effect, 62, 63–4
- Greenhouse gas (GHG), 62f. *See also* Carbon dioxide; Methane
  - climate change from, 570–1, 571f
  - global warming and, 64
- Grime's CSR triangle, 232, 233f, 234–5, 234f
- Gross percentage forest cover loss (GFCL), 507f
- Gross primary productivity (GPP), 649
  - climate change and, 652
  - ocean depth and, 667, 667f
  - RE and, 665, 665f, 676, 676f
  - in seasonal environments, 652
- Groundwater, 685, 701
- Growth
  - essential resources for, 96
  - of genet, 106f
  - interspecific competition and, 240
  - intrinsic, 255
  - life history trade-offs and
    - survivorship and, 212, 212f
    - Y model for, 211f
  - of modular organisms, 104
  - of parasite hosts, 399–401, 401f
  - plant defences and, 296–7, 297f
  - of plants, defoliation and, 307–8
  - of populations, 455–6, 458–9, 459f, 459t, 461
  - rate hypothesis, 304
  - resources and, niche differentiation and, 255
  - RGR, 80f
  - temperature and, 39–41, 41f
- Growth-differentiation balance hypothesis, 304–5
- GS. *See* Geometric series
- Guilds, 526–7, 529–30, 529f

- Gulls, 11
- Gut
- metagenome in
    - of bees, 436
    - of vertebrates, 433–4, 435f
  - mutualism in
    - in herbivores, 432–3, 433f
    - in insects, 434–6, 435f
    - in ruminants, 432–3, 433f, 434t
    - in vertebrates, 433–4, 433f, 434t
- Habitats
- anoxic, 359
  - area, species richness and, 619, 619f
  - checkerboards, 533
  - commensalism in, 417
  - CR and
    - age of maturity and, 220
    - comparative classification of, 216–17, 218f
    - reproductive investment and, 218
  - degradation of, 507–8
  - destruction of, 507
  - disturbance of, 508
    - by grazing, 537
  - diversity of, island biogeography and, 621, 621f
  - extinction risk and, 503, 505
  - hierarchy of, of communities, 572–3, 573f
  - humans and, extinction risk and, 505f, 507–8, 507f
  - life histories and, 214–21, 217f–19f, 221f
  - restoration of, for squirrels, 188–9, 189f
  - stress on, 232
- Habitat/species management areas, 605
- Handlers, 290–1
- Handling time, 326
- HANPP. *See* Human appropriation of net primary production
- Harrison, Smith and Slobodkin (HSS), 549–50
- Harsh environments, species richness and, 417, 616
- Harvesting effort, 489–90, 490f
- Harvest management
- depensation in, 493–5, 494f
  - dynamic pool models for, 495–6, 496f
  - environmental fluctuations and, 495
  - EOY and, 492, 492f
  - of fish, 490–2, 491f, 492f, 495–9, 496f–500f
  - fixed harvesting rate and, 488f, 489
  - mortality compensation and, 491–2, 493
  - MSY and, 488–90, 488f–90f
  - multiple equilibria for, 493–5, 494f
  - overexploitation and, 487–8
  - profit sharing in, 493, 493f
  - sustainability in, 491, 497–8, 498f
  - underexploitation and, 487–8
- Hazards, 58–9
- Heavy metals, 54, 60, 418, 616
- Helminths, 380, 392, 392f, 407
- Hemicryptophytes, 29
- Hemiparasites, 381
- Herbivores, 93
- ATCH for, 550, 550f
  - bacteria on plants from, 310
  - birds as, 38–9, 39f
  - CE of, 672, 672f
  - defoliation by, 307–8, 309, 309f
  - faeces of, detritivores and, 373–4, 373f, 374f
  - food preferences of, 288
  - in four-trophic-level system, 545
  - grasses and, commensalism with, 417–18
  - grazing by, 285
  - gut mutualism in, 432–3, 433f
  - meta-analyses of, 312–14, 313f
  - nutritional contents of, 95
  - plants and
    - coevolutionary arms race of, 295
    - competition and, 308, 309f
    - defences against, 294–307, 296f–307f
    - fecundity and, 311–12, 311f
    - fungi on, 310
    - tolerance of, 307–14, 308f–13f
  - predation by, 285
  - secondary productivity by, 669
  - survivorship of plants and, 308–10
  - timing of, 312, 312f
  - viruses on plants from, 310
- Herd immunity, 406–7, 406f
- Herpes simplex virus, 394
- Heterocarpy, 184
- Heterogeneity
- inbreeding and, 510
  - interspecific competition and, 263–9, 264f–9f
  - invasion and, 269, 269f
  - for parasitoids, 350–1
  - predation population dynamics of, 349–55, 352f–5f
  - spatial, species richness and, 616
- Heterotrophic respiration (RH), 649, 686
- Heterotrophs, 65, 95
- Hidden trade-offs, 211
- Hill numbers, 575
- Historical checkerboards, 533
- Holdfasts, 57
- Holoparasites, 381

- Home field advantage, for plant litter decomposition, 367–8, 368f, 369f
- Home-loving. *See* Philopatry
- Hosts. *See also* Parasite hosts  
 fitness of, symbiosis and, 421–2  
 microparasites and, life cycle of, 407
- Hot spots, of species richness, 603, 640, 735
- HSS. *See* Harrison, Smith and Slobodkin
- Human appropriation of net primary production (HANPP), 650–1, 651f
- Humans. *See also specific topics*  
 in Anthropocene, 708–9  
 biodiversity and, 663, 663f  
 biodiversity loss and, 709–11, 738  
 births of, 145, 146f  
 community size spectra and, 578, 578f  
 deaths of, 145, 146f  
 exploitation by, destabilising Allee effects and, 338  
 global carrying capacity for, 144–8, 144f–6f  
 habitat and, extinction risk and, 505f  
 as hyper-keystones, 554–5, 555f  
 infections of, cycles of, 405, 406f  
 intraspecific competition of, 144–8, 144f–6f  
 invasive species and, 23–4, 23f, 24t  
 LUC and, 724–9, 724f  
 nitrogen cycle and, 704  
 phosphorus cycle and, 702
- Humidity, 54, 71
- Hybridisation, 3, 6
- Hydrological cycle, 699–701, 699f, 701f
- Hyperaccumulators, 60
- Hyper-keystones, 554–5, 555f
- Hypoxia, 728, 728f
- IBP. *See* International Biological Programme
- Ice age, 19f, 20–2
- ICS. *See* International Commission on Stratigraphy
- Ideal free distribution, 347, 348f
- IFS. *See* Integrated farming systems
- Immigration  
 to islands, 619–21  
 in metapopulations, 192, 198
- Immobilisation, 358
- Immune response  
 to infections, 405  
 of parasite host, 390
- Immunisation, herd immunity and, 406–7, 406f
- Importation, for biological control, 483–4
- Inbreeding, 182, 182f  
 depression, as extinction risk, 504–5, 504f  
 heterogeneity and, 510
- Index of abundance, 108, 108f
- Indirect life cycles, of parasites, 379
- Indirect transmission, 396
- Individual-based data, on species richness, 606
- Induced dormancy, 113
- Inducible defences, 294–5  
 for parasitoids, 303, 303f
- Industrial melanism, 7–8, 7f
- Industrial processes, decomposition and, 359
- Infections, 378  
 of humans, cycles of, 405, 406f  
 immune response to, 405  
 population dynamics of, 402–10, 405f, 406f, 409f, 410f  
 protozoan, prevalence of, 405  
 transmission threshold of, 403  
 zoonotic, 384
- Inhibitory resources, 97
- Innate dormancy, 113
- Inoculation, for biological control, 483–4
- Insecticides  
 DDTs, 484, 731  
 in IPM, 485  
 plant defences and, 296
- Insects, 140f  
 allopatric speciation of, 12  
 bacteria and, symbiosis of, 436–7  
 biological pesticides for, 484  
 cold tolerance of, 46  
 as decomposers of dead bodies, 375, 375f, 376t  
 enzymes in, 95  
 food switching by, 289, 289f  
 frequency-dependent predation of, 539f  
 for fungi colonisation, in dead wood, 366–7, 367f  
 gut mutualism in, 434–6, 435f  
 importation of, for biological control, 483  
 interspecific competition of, 525f  
 at low temperatures, 45  
 migration of, 171, 171f  
 oxygen and, 90  
 pest control of, bacteria for, 438, 438f  
 plant defences against, 300–3  
 as pollinators  
 as keystone species, 553  
 mutualisms of, 429–32, 430f  
 as unitary organisms, 102
- Integral projection model (IPM), 125–7, 126f
- Integrated farming systems (IFS), 485–6, 487f
- Integrated pest management (IPM), 485–7, 486f, 487f
- Intensity, of parasites, 382

- Intercropping, 277, 277f, 418–19
- Interference, 133, 134f
  - aggregated distributions and, 347–9, 348f
  - coefficients, 348–9, 348f
  - exploitation and, 248–9, 249f
  - predation population dynamics and, 330–9, 332f, 333f, 335f, 336f, 337t, 338f
  - pseudo-, 350
- Intergovernmental Panel on Climate Change (IPCC), 570–1, 647, 711–13
- Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem services (IPBES), 711
- Intermediate disturbance hypothesis, 593–4, 594f
- International Biological Programme (IBP), 647
- International Commission on Stratigraphy (ICS), 708, 709f
- International Union for Conservation of Nature and Natural Resources (IUCN), 502–3
  - on population size, 709–10
  - protected area classification by, 604–5
  - Red Line assessment of, 520–1
- Interspecific competition
  - aggregated distributions and, 266–7, 267f
  - apparent competition and, 270–4, 271f–3f
  - of bacteria, 282, 282f
  - coexistence and
    - in birds, 244–5, 245f
    - in diatoms, 244, 245f
    - in fish, 243–4, 243f
    - niche overlap and, 261–2, 261f
    - niche similarity and, 262–3, 263f
  - colonisation and, 263–9, 264f–9f
  - ecological effects of, 274–6, 275f–7f
  - environmental fluctuations and, 265–6, 266f
  - evolution and, 247, 278–83, 278f–83f
  - first come best served, of fish, 265
  - in fish, 243–4, 243f
  - fundamental niche and, 244
  - general features of, 247–9
  - growth and, 240
  - heterogeneity and, 263–9, 264f–9f
  - of insects, 525f
  - Lotka–Volterra model for, 249–52, 250f, 251f, 252t
  - of parasites, 389–92, 391f, 392f
  - with phytoplankton, 259f
    - phosphorus and, 241–2, 241f
  - for plants and nitrogen, 242–3, 242f
  - prevalence of, 534
  - of protozoans, 282–3, 283f
  - realised niche and, 244
  - of roots, 260, 260f
    - of shoots, 260, 260f
    - survivorship and, 240
    - between unrelated species, 245–6, 246f
- Intertidal zone, 57–8, 57f
- Intimidation, 318–19, 319f, 331, 338–9, 338f
- Intransitive competition, 533–4
- Intraspecific competition, 132–68
  - density-dependent growth and, 135, 136f
  - fecundity and, 135, 136f, 140
  - of humans, 144–8, 144f–6f
  - mathematical models for, 148–61, 149f–52f, 154f–61f
  - of parasites, 389–90
  - of plant modular organisms, 137f
  - in population models, 149–50, 150f
  - population size and, 141–8, 141f–6f
  - prey isoclines with, 332–4, 332f
  - quantification of, 139–41, 140f
  - self-thinning and, 161–8, 163f, 164f, 166f–8f
- Intrinsic growth rate, 255
- Intrinsic rate of natural increase, 54, 122
- Inundation, for biological pest control, 484
- Invasion, 180
  - dynamics of, 189–91, 190f, 191f
  - heterogeneity and, 269, 269f
  - in lakes, 181f, 190–1, 190f
  - life history fast–slow continuum and, 230–1, 231f
  - LUC and, 738
  - plant tolerance for, 308, 308f
  - species distribution modelling for, 195, 196f
- Invasive plant species (IPSSs), 736–8, 736f, 737f
- Invasive species
  - current and projected distribution of, 738, 739f
  - as extinction risk, 509, 509f
  - humans and, 23–4, 23f, 24t
- Invertebrates
  - algae and, photosynthesis symbiosis of, 438f, 439–40
  - as carnivores, 364
  - detritivores and, 375–6
    - faeces of, 372–3, 372f
  - interspecific competition of, 266–7, 267f
  - at low temperatures, 45
  - as modular organisms, 102
  - neonicotinoids in, 481
  - niche models for, 37–8, 37f, 38f
  - parasite host defences of, 385
  - pH and, 616, 617
  - in streams, agricultural pollution and, 29, 29f
- IPBES. *See* Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem services
- IPCC. *See* Intergovernmental Panel on Climate Change

- IPM. *See* Integral projection model; Integrated pest management
- IPSS. *See* Invasive plant species
- Iron, 87  
 as inhibitory resource, 97  
 in oceans, global warming and, 698
- Irreplaceability, 640–1, 641f
- Islands  
 biogeography of, equilibrium theory of, 619–31, 620f–31f  
 birds on, checkerboard distributions of, 532, 532f  
 colonisation on, 622, 624, 625f  
 evolution on, 18–19, 18f  
 extinction on, 620–1  
 immigration to, 619–21  
 metapopulations and, 197, 198f  
 as nature reserves, 629–31, 629f–31f  
 species richness and, 619–31, 619f–31f
- Isoclines  
 of interspecific competitors, 250–1, 250–1f, 474, 474f  
 of predator, with mutual interference, 333f  
 in predator–prey models, 321–2, 332–8, 474, 322f, 333–8f, 474f  
 of prey, with intraspecific competition, 332–4, 333f  
 vertical, 334, 350  
 ZNGI, 95–7, 96f, 253–7, 254f
- Isopods, 111  
 for decomposition, 372–3, 372f  
 humidity and, 54  
 plant defences and, 301–2, 301f
- Iteroparous life cycle, 108–10
- IUCN. *See* International Union for Conservation of Nature and Natural Resources
- Ivermectin, 414
- JA. *See* Jasmonic acid
- Janzen–Connell hypothesis, 538–9, 538f
- Jasmonic acid (JA), 299–300, 300f
- Key factor analysis, of abundance, 456–8, 456t, 457f, 459t
- Keystone species, 553–4, 554f, 555f
- Kleiber’s law, 99
- Knock-out mutants, 296
- K-selection, 226–7, 228f, 238
- K values, 139–40, 140f
- LA. *See* Leaf area
- Lack clutch size, 223–4, 224f, 225f
- LAI. *See* Leaf area index
- Lakes  
 acid rain and, 60  
 anoxic habitats in, 359  
 autochthonous production in, 665  
 carbon dioxide in, 79, 79f  
 eutrophication in, 728, 728f  
 fish in, acid rain and, 61  
 invasion in, 181f, 190–1, 190f  
 nutrient flux in, 690–2, 691f  
 photosynthesis in, 71f  
 solar radiation in, 71f
- Lambda ( $\lambda$ ) contribution analysis, of abundance, 458, 459t, 460f
- Land sharing, 610
- Land sparing, 610
- Land-use change (LUC), 724–9, 724f  
 invasion and, 738
- La Niña, 49, 50f
- Latitude  
 productivity and, 651  
 as species richness gradient, 631–3, 632f, 633f
- LDMC. *See* Leaf dry matter content
- Leaf area (LA), 232
- Leaf area index (LAI), 142
- Leaf dry matter content (LDMC), 232
- Lichen  
 from algae-fungi mutualism, 444–5, 445f, 446t  
 in tundra, 25
- Life cycles, 108–11  
 graph, 124, 124f  
 of microparasites and hosts, 407  
 of parasites, 379  
 of plants, 110  
 semelparity of, 108–11  
 of unitary organisms, 102, 109f
- Life histories  
 allometric relationships in, 235–6, 235f, 236f  
 for mammals, 238–9  
 components of, 207–10, 207f–9f  
 development and, 207–8  
 evolution and, 206–39  
 fast–slow continuum for, 227–31, 229f–31f  
 for mammals, 238–9, 239f  
 habitats and, 214–21, 217f–19f, 221f  
 K-selection and, 226–7, 228f, 238  
 natural selection and, 208  
 phylogenetic constraints on, 232–9, 236f–9f  
 on mammals, 238–9, 239f  
 reproductive allocation and, 208, 208f, 218–20, 219f  
 reproductive value and, 208–10, 209f  
 resources in, 210  
 r-selection and, 226–7, 228f, 238  
 seeds and, 206

- Life histories (*cont'd*)  
 size and, 207, 207f  
 trade-offs in, 210–14, 210f–12f  
   with CR, 210f, 213  
   fitness contours and, 216, 217f  
   genotypes and, 212–13  
   with offspring size and number, 213–16, 214f, 215f, 221–6, 222f–5f  
   option sets and, 214–15, 217f  
   for survivorship and growth, 212, 212f
- Life table response experiment (LTRE), 127–8, 128f
- Life tables  
 cohort, 114–17, 114f, 115f, 116f, 123, 123f  
 for modular organisms, 121, 121f  
 static, 118–20, 120f  
 variables of, 122–3
- Light compensation point, 69–71, 71f
- Living planet index, 604, 604f
- Lizards  
 as ectotherms, 44  
 food chain length of, 566  
 in four-trophic-level system, 545  
 frequency-dependent predation of, 539f  
 guild competition of, 529–30, 529f
- Local adaptation, 3–5, 5f
- Local stability, 552
- Logistic equation, for continuous breeding, 154–5, 155f
- Log series (LS), 576, 576f
- Long-lived agents, 396
- Lotka–Volterra model  
 competitive exclusion principle and, 240, 255–6  
 food chain length and, 564  
 for interspecific competition, 249–52, 250f, 251f, 252t  
 for predation population dynamics, 320–2, 322f, 350
- LS. *See* Log series
- LTRE. *See* Life table response experiment
- LUC. *See* Land-use change
- Lyme disease, dilution effect and, 400–1, 400f
- MacArthur fraction (MF), 576, 576f
- Macroecology, defined, 605
- Macronutrients, 87
- Macroparasites, 380–1  
 in crops, 408, 409f
- Magnesium, 87
- Mainland–island metapopulations, 198, 198f
- Malaria, 407
- Malathion, 480
- Mammals  
 allometric relationships in life histories for, 238–9, 239f  
 biodiversity conservation of, 15  
 carcasses of, decomposition of, 367  
 clutch size of, 225, 225f  
 coinfection of, 392, 393f  
 conservation of, 515, 517f  
 as detritivores, 376  
 dispersal of, by sex, 185–6, 186f  
 extinction risk of, 503  
 in extreme temperatures, 53f  
 fecundity of, predator intimidation and, 318–19, 319f  
 interspecific competition of, 278–9, 278f  
 as invasive species, 24f  
 life histories of, 210  
   fast–slow continuum for, 229–30, 230f, 238–9, 239f  
   phylogenetic constraints on, 238–9, 239f  
 marginal value theorem for, 346–7, 346f, 347t  
 marsupials, 17–18, 17f, 27–8, 429  
 in mesopredator release, 547, 547f  
 metabolic scaling of, 99  
 metapopulation capacity for, 204–5, 204f  
 migration of, 170, 173–4, 174f  
 phocine distemper in, 409, 410f  
 physical defences of, 295  
 population cycles for, 467–74, 467f–9f, 472f, 473f  
 population size of, 709, 710f  
 predator–prey cycles of, 325  
 reproductive allocation in, 219f, 220  
 species richness of, 615f, 633f  
 survivorship curves of, 119–20, 119f  
 as unitary organisms, 102
- Manganese, 87
- Mangroves  
 colonization gaps in, 596, 596f, 596t  
 tsunamis and, 58
- Maquis, 25f, 26
- Marginal value theorem, 341–7, 342f–5f, 347t
- Marine protected areas (MPAs), 637–8, 637f
- Marine sediments, zonation of, 359, 359f
- Marine zoning plans, 642–3, 643t, 644f
- Markov chain model, 586
- Marsupial mammals, 17–18, 17f, 27–8, 429
- Mass-effects metacommunity, 600–1
- Mast seeding, 212
- Mast years, 181, 327–8, 328f
- Matter flux, 678–707. *See also* Nutrient flux  
 biogeochemical cycles of, 679–80
- Maximum sustainable yield (MSY), 488–90, 488f–90f
- MEA. *See* Millennium Ecosystem Assessment
- Mean intensity, of parasites, 382
- Mean resistance, virulence and, 395f
- Mean trophic level (MTL), 498–9, 499f, 500f
- Measles, immunisation for, 407

- Megafauna, 361
- Mercury, 731, 733
- Mesofauna, 361
- Mesopredator release, 547, 547f
- Metabolic theory of ecology, 97–101
- Metabolism
  - allometric scaling of, 97–101, 98f, 99f
  - temperature and, 39–41, 40f
- Metacommunity, 530, 599–602, 599f–602f
  - abundance in, 601, 601f
  - dispersal in, 601, 601f
  - neutral theory for, 600
  - patches in, 599
  - species richness in, 601
- Metagenome, in gut
  - of bees, 436
  - of vertebrates, 433–4, 435f
- Metapopulation
  - abundance and, 194f, 195
  - of butterflies, 198, 198f, 201f, 202–3, 203f
  - capacity, 201
    - for birds, 204–5, 204f
    - for mammals, 204–5, 204f
  - conservation of, 515–18, 516f–18f
  - defined, 192
  - dispersal and, 192–205, 194f, 196f, 198f–201f, 203f, 204f
  - dynamics of, 195–205, 198f–201f, 203f, 204f
  - extinction and, 197–203
  - islands and, 197, 198f
  - mainland–island, 198, 198f
  - parasites in, 408
  - of parasitoids, 353f
  - persistence and, 194f, 195
  - recolonisation and, 197–203
  - uninhabited habitable patches and, 195–7
- Methane, 62, 685
- MF. *See* MacArthur fraction
- Microbivores, 361–5, 362f–5f
- Microparasites, 379, 403–4
  - life cycle of, 407
- Microplastics, 731–2, 732f, 733
- Migration. *See also* Emigration; Immigration
  - of birds, climate change and, 716, 716f
  - conservation for, 172–5, 173f, 174f
  - defined, 170
  - of insects, 171, 171f
  - as one-way trip, 172
  - patterns of, 170–5
  - of predators, 349
    - between resources, 172
- Mildew, 381
- Millennium Ecosystem Assessment (MEA), 647–8, 727, 738
- Mimicry, 314f, 316
- Mineralisation, 358
- Minerals, 87–90, 88f, 89f
  - decomposers and, 369–70, 370f
  - in estuaries, 27
  - as inhibitory resource, 97
  - NPP and, 659–61, 660f
- Minimum viable populations (MVPs), 511–12
- Mites
  - as macroparasites, 380–1
  - temperature and, 40, 41f
- Modular organisms, 102–4, 103f
  - birth of, 106f
  - competition of, 139, 139f
  - constant yield of, 135
  - counting individuals of, 106–8
  - death of, 106f
  - growth of, 104
  - integration of, 106, 107f
  - life table for, 121, 121f
  - neighbours of, 139, 139f
  - plants as, intraspecific competition of, 137f
- Modular population, 104
- Molluscs
  - calcium carbonate for, 63
  - humidity and, 54
    - as invasive species, 24, 24f
- Molybdenum, 87
- Monomodal community size spectra, 577f
- Monophagous, 93
  - foragers, 285
- Monotonical community size spectra, 577f, 578
- Monotonic damping, 151
- Mother knows best hypothesis, 288
- Moths
  - dispersal of, 178, 178f
  - food of, 54
  - industrial melanism and, 7–8, 7f
  - temperature and, 54
- Movement, 169–205. *See also* Dispersal; Migration
  - resources and, 169
  - of seeds, 169
- MPAs. *See* Marine protected areas
- MSY. *See* Maximum sustainable yield
- MTL. *See* Mean trophic level
- Müllerian mimicry, 316
- Multimodal community size spectra, 577–8, 577f
- Multiple equilibria
  - for abundance, 474–5, 474f, 475f
  - for harvest management, 493–5, 494f

- Mussels  
 barnacles and, 536  
 colonisation gaps of, 595f, 596  
 specialisation of, 3
- Mutual antagonism, Lotka–Volterra model and, 252
- Mutual interference, 331, 332f, 334
- Mutualism  
 of ants  
 aphids and, 424, 424f  
 butterflies and, 425–6, 426f  
 fungi and, 426, 427–8, 427f  
 plants and, 422–5, 422f, 423f, 425f  
 between decomposers and plants, 370–1  
 of corals and algae, 439–40  
 defined, 416  
 of detritivores, 371–2  
 of fungi  
 beetles and, 425–6  
 plants and, 407, 440–3, 441f, 441t  
 roots and, 440–3, 441f, 441t  
 in gut  
 in herbivores, 432–3, 433f  
 in insects, 434–6, 435f  
 in ruminants, 432–3, 433f, 434t  
 in vertebrates, 433–4, 433f, 434t  
 in human agriculture, 423–4  
 of keystone species, 553  
 models of, 451, 452f  
 in mycorrhizas, 440–3, 441f, 441t  
 of nitrogen-fixing bacteria, 30, 416, 446–51, 447f–9f  
 of plants  
 bees and, 451, 452f  
 fungi and, 407, 440–3, 441f, 441t  
 of pollination, 429–32, 430f  
 protection in, 420–3, 421f, 422f  
 as reciprocal, 417  
 of rhizobia and legumes, 446–50, 448f, 449f  
 of seed dispersal, 428–9
- MVPs. *See* Minimum viable populations
- Mycorrhizal networks, 443, 443f
- Mycorrhizas, 88–90  
 mutualism in, 440–3, 441f, 441t
- Myxomatosis, 393–4, 394f
- NAME. *See* Numerical Atmospheric-dispersion Modeling Environment
- NAO. *See* North Atlantic Oscillation
- Nasty neighbour hypothesis, 161, 161f
- Natal dispersal, 170
- National parks, 605
- Natural hosts, 383
- Natural monuments, 605
- Natural selection, 1–2  
 life histories and, 208  
 optimal foraging theory and, 291  
 phylogenetic constraints and, 232  
 of sessile organisms, 3  
 for virulence, 394
- Nature reserves, 604  
 islands as, 629–31, 629f–31f
- N*-dimensional hypervolume, 33, 97
- Near-threatened species, 711
- Necrotrophic parasites, 379
- Negative binomial encounters, 350
- Neighbours  
 of modular organisms, 139, 139f  
 nasty hypothesis for, 161, 161f
- Nematodes  
 birds and, 411–14, 412f, 413f  
 humidity and, 54  
 as macroparasites, 380
- Neonicotinoids, 481
- Net ecosystem productivity (NEP), 649
- Net photosynthesis, 69–73
- Net primary productivity (NPP), 615, 615f, 649, 649t, 650, 650f, 652f  
 biomass and, 655–6, 655f  
 detritivores and, 675, 675f  
 in forests, 686, 686f  
 minerals and, 659–61, 660f  
 PAR and, 656–7, 656f  
 in oceans, 666, 666f  
 in seasonal environments, 659, 659f  
 soil texture and, 658–9  
 SST and, 666, 666f  
 in succession, 652, 661  
 temperature and, 658, 658f  
 water and, 657, 657f, 658f, 660f
- Net recruitment curves, 142–3, 142f, 143f
- Net reproductive rate, 143, 149–50
- Neutral competition, 262
- Neutral theory  
 for metacommunities, 600  
 of niche differentiation, 530–1, 531f
- Niches, 34f–9f  
 breadth of, in species richness, 607  
 complementarity of, community module competition, 525–8, 526f–8f  
 of cover crops, 55, 55f  
 differentiation of  
 coexistence and, 244–5, 262

- coexistence without, 247–8, 248f  
 in community module competition, 525–31, 529f, 531f  
 neutral theory of, 530–1, 531f  
 resources and growth and, 255  
 in seasonal environments, 618  
 divergence of, in grasses, 281, 281f  
 fundamental, 36, 38–9, 39f, 244  
 models for, 35–6, 36f  
 ordination of, 33–5, 37–8, 37f, 38f  
 overlap of  
   interspecific competition coexistence and,  
     261–2, 261f  
   in species richness, 607  
 realised, 36, 244, 294  
 resources of, 97  
 similarity of, interspecific competition coexistence and,  
     262–3, 263f  
 species distribution modelling for, 195, 196f  
 successional, 587, 587f  
 temperature in, 39–56, 40f–3f, 45f–7f, 50f–3f, 55f  
 Nicholson, A. J., 454–5  
 Nicholson–Bailey model, of predation population dynamics,  
     324–5, 350  
 Nickel, 60  
 Nitrates, 88, 95, 695–6, 696f  
 Nitrogen, 87  
   in agriculture, 725–6  
   from ammonia, 95  
   carbon dioxide and, 79  
   carbon : nitrogen ratio, 93, 95  
   decomposers and, 370  
   on continental shelves of oceans, 693–5, 695t  
   cycle, 703f, 704  
   decomposers and, 369–70, 370f  
   in estuaries, 692–3, 692f, 693f  
   in forests, 726, 726f  
   from nitrates, 88, 95  
   NPP and, 659–61, 660f  
   in oceans, 665–6  
   photosynthetic capacity and, 71–3, 72f  
   plants and, interspecific competition for, 242–3, 242f  
   in rivers, 727–8  
 Nitrogen dioxide, 60f  
 Nitrogen-fixing bacteria  
   as keystone species, 553  
   mutualism of, 30, 416, 446–51, 447f–9f  
   succession and, 450–1  
 Nitrous oxide, 62  
 Non-kin competition, 183, 183f  
 Non-seasonal environments, specialisation in, 618  
 Normalisation constant, 97, 98  
 North Atlantic Oscillation (NAO), 49, 50f, 51f  
 NPP. *See* Net primary productivity  
 Null model analysis, for community competition, 528–33,  
     529f, 534  
 Numerical Atmospheric-dispersion Modeling Environment  
   (NAME), 176, 176f  
 Nurse plants, 418  
 Nutrient budgets, 680–1  
   in aquatic biomes, 689–707  
   on continental shelves of oceans, 693–5, 695t  
   in oceans, 696–7, 697f, 698f  
   in terrestrial biomes, 680f, 682–9, 683f–6f, 689f  
 Nutrient flux  
   deforestation and, 688–9, 689f  
   downstream effects of, 726–7  
   energy flux and, 678–9, 679f  
   in estuaries, 692–3, 692f, 693f  
   in lakes, 690–2, 691f  
 Nutrients  
   loss of, in atmosphere, 684–5  
   pollution of, in aquatic biomes, 681–2, 682f  
   spiraling, in streams, 689–90, 690f  
 Obligate mutualism, 371–2  
 Oceans  
   acidification of, 720–1, 722f, 723f  
     climate change and, 707, 707f  
   anoxic habitats in, 359  
   carbon dioxide in, photosynthesis and, 63  
   continental shelves of, nutrient budgets on, 693–5, 695t  
   depth of, GPP and, 667, 667f  
   eutrophication in, 728, 728f  
   high temperatures in, 48  
   iron in, global warming and, 698  
   NPP of, 649t, 650  
   nutrient budgets in, 696–7, 697f, 698f  
   PAR of, NPP and, 666, 666f  
   plankton in, 696–7  
   primary productivity in, SST and, 666  
   productivity of, 651  
   upwelling in, 665–6, 668f  
   warming of, fish and, 718, 719f  
 Offspring size and number, life history trade-offs with, 213–16,  
     214f, 215f, 221–6, 222f–5f  
 Oligophagous foragers, 285  
 OLS. *See* Ordinary least squares analysis  
 Omnivores  
   in community modules, 540  
   food preferences of, 288  
 One-generation cycles, in predation population  
   dynamics, 324–5

- Optimal foraging theory  
 for plants, 343–5, 344f  
 for predators, 289–92, 290f  
 versus food switching, 292
- Optimal sowing rates, for conservation, 138, 138f
- Option sets, 214–15, 217f
- Orchid mycorrhizas, 443
- Ordinary least squares analysis (OLS), 237, 238f
- Ordination  
 of communities, 578–81, 580f  
 of niches, 33–5, 37–8, 37f, 38f
- Osmoregulation, 297
- Outbreaks  
 in abundance, 474f, 475f  
 secondary pest, 479–80, 480f
- Outbreeding, 182, 182f
- Overcompensating density dependence, 134, 140
- Overexploitation  
 density-dependent emigration and, 186  
 extinction from, 505–7, 506f  
 of fish, 733–6, 734f, 735f  
 harvest management and, 487–8
- Overyielding, 557, 662
- Oxalic acid, 295
- Oxygen, 90–1, 91f  
 hypoxia and, 728, 728f
- Ozone depletion, 730, 730f, 733
- PACF. *See* Partial autocorrelation function
- PAR. *See* Photosynthetically active radiation
- Paradox of enrichment, 609
- Paradox of the plankton, 247–8, 248f, 265–6
- Parallel evolution, 17–18
- Parasite hosts  
 aggregated distributions of, 381–2, 382f  
 coevolution of, 392–5, 394f, 395f  
 competition for resources of, 388–92, 389f–92f  
 constant final yield with, 388–90, 390f  
 defences of, 384–8, 385f–7f  
 trade-offs with, 386f, 387f  
 diversity of, disease and, 399  
 fecundity of, 399–401  
 growth of, 399–401, 401f  
 immune response of, 390  
 population dynamics of, 410–15, 411f–13f, 415f  
 as predators, 384  
 resistance of, 388, 389f  
 as resources, 384  
 specificity of, 383–4  
 as superspreaders, 398, 398f  
 survivorship of, 399–401  
 tolerance of, 388, 389f  
 transmission in, 396–402, 397f–401f  
 virulence and, 388, 389f
- Parasites  
 allopatric speciation of, 12  
 of crops, 378  
 as detritivores, 379  
 direct life cycles of, 379  
 disease and, 378–415  
 diversity of, 378–81, 380f  
 essential resources of, 96  
 exploiter-mediated coexistence of, 540  
 extinction from, 534–5, 535f  
 food and, 414–15, 415f  
 in food webs, 567, 568f  
 indirect life cycles of, 379  
 interspecific competition of, 389–92, 391f, 392f  
 life cycles of, 379  
 in metapopulations, 408  
 plants and, competition of, in salt marshes, 402, 402f  
 population cycles and, 466  
 as predators, 284–5  
 protozoan as, beetles and, 411f  
 resources of, 92  
 on seeds, 286  
 transmission and virulence of, trade-offs with, 394
- Parasitoids  
 aggregated distributions of, 340f, 349–50, 349f  
 aggregation of risk for, 355f  
 as biological control agents, 356  
 consumer–resource systems for, 320  
 continuous redistribution of, 351  
 coupled oscillations of, 286f  
 defensive symbioses of, 438  
 delayed density dependence for, 323f  
 heterogeneity for, 350–1  
 importation of, for biological control, 483  
 inducible defences for, 303, 303f  
 marginal value theorem for, 343, 344f, 346  
 metapopulations of, 353f  
 Nicholson–Bailey model for, 324–5  
 as predators, 285  
 as searchers, 324  
 type 2 functional response for, 326, 326f
- Paris Climate Agreement, 733
- Partial autocorrelation function (PACF), 461–2, 462f
- Partial rate correlation function (PRCF), 462, 462f, 464f, 465
- Partial refuges, 350
- Particulate organic carbon (POC), 612, 612f
- PAs. *See* Protected areas

- Passive dispersal, 175–9, 178f
- Patches and patchiness  
 colonisation in, 595f  
 of dispersal, 180  
 metapopulations and, 192–5  
 foraging in, 339–49, 340f–2f, 344f–6f, 347t, 348f, 349f  
 in metacommunities, 599  
 metapopulations in, 195–7  
 species richness of, 575  
 stochastic colonisation of, 543
- PCMs. *See* Phylogenetic comparative statistical methods
- Perfectly substitutable resources, 96
- Perimeter : area ratio, 594, 594f, 596
- Permafrost, 91–2, 92f, 570–1, 571f
- Permanent wilting point, 76, 77f
- Persistence, metapopulations and, 194f, 195
- Persistent organic pollutant (POP), 731, 733
- Pest control, 478–87  
 biological agents for, 483–5, 483t  
 of insects, bacteria for, 438, 438f  
 IPM for, 485–7, 486f, 487f
- Pesticides. *See also* Insecticides  
 bee gut metagenome and, 436  
 biological, 484  
 habitat loss from, 508  
 resistance to, 481–2, 482f  
 unintended consequences of, 479–81, 480f
- Pests  
 colonisation of, 484  
 defined, 478  
 EIL of, 478–9, 478f  
 ET of, 478–9
- PET. *See* Potential evapotranspiration
- Pgi*. *See* Phosphoglucose isomerase gene
- PGLS. *See* Phylogenetic generalised least squares
- pH  
 acid rain and, 722  
 carbon dioxide and, 79  
 in harsh environments, 616, 617f  
 in niches, 33  
 of soil, 54–56, 55f  
 species richness and, 616, 617f
- Phalanx-type plants, 179, 179f
- Phanerophytes, 29
- Phenolics, 298
- Philopatry (home-loving), dispersal and, 181  
 sex and, 186
- Phocine distemper, 409, 410f
- Phosphoglucose isomerase (*Pgi*) gene, 203, 203f
- Phospholipid fatty acids (PLFA), 361
- Phosphorus, 87, 89–90  
 in agriculture, 725–6  
 from atmosphere, 683f  
 cycle, 702, 703f  
 decomposers and, 369–70, 370f  
 in forests, 726, 726f  
 NPP and, 659–61, 660f  
 in oceans, 665–6  
 phytoplankton and, interspecific competition for, 241–2, 241f  
 in primary succession, 680–1, 681f  
 in rivers, 727–8
- Photoautotrophs, 65
- Photoinhibition, 67
- Photosynthesis  
 in aquatic biomes, 27  
 carbon dioxide and, 69  
 in oceans, 63  
 carbon in, 66  
 defoliation and, 307  
 in lakes, 71f  
 net, 69–73  
 photoautotrophs and, 65  
 symbiosis of, of invertebrates and algae, 438f, 439–40  
 water conservation and, 74–5, 76f
- Photosynthetically active radiation (PAR), 67  
 light compensation point and, 69–71  
 NPP and, 656–7, 656f  
 of ocean, NPP and, 666, 666f  
 photosynthetic capacity and, 73  
 in streams, 664, 664f
- Photosynthetic capacity, 72–3, 72f
- Photosynthetic photon flux density (PPFD), 71f
- Phylogenetic comparative statistical methods (PCMs),  
 237–8, 238f
- Phylogenetic constraints  
 on life histories, 232–9, 236f–9f  
 on mammals, 238–9  
 natural selection and, 232
- Phylogenetic generalised least squares (PGLS), 237, 238f
- Physical defences  
 of animals, 315–16, 315f  
 of plants, 295
- Phytomining, 60
- Phytoplankton  
 food web trophic cascade of, 546, 546f  
 GPP of, 667, 667f  
 interspecific competition in, 259f  
 phosphorus and, 241–2, 241f  
 niches for, 34–5, 35f  
 species richness of, 609, 611
- Placental mammals, species distribution of, 17–18, 17f
- Plague, 384

- Planetary boundaries, 738–40, 741f, 741t
- Plankton. *See also* Phytoplankton  
 calcium carbonate for, 63  
 coexistence of, 247–8, 248f  
 in estuaries, 692–3, 692f, 693f  
 microplastics and, 732  
 in oceans, 696–7  
 paradox of, 247–8, 248f, 265–6  
 physical defences of, 295
- Plants. *See also* Photosynthesis; Resources  
 allometric relationships in, 100, 100f  
 antagonistic coevolution in, 394–5, 395f  
 ants and, mutualism of, 422–5, 422f, 423f, 425  
 apparent competition of, 271–2, 272f  
 in aquatic biomes, clonal growth dispersal in, 179  
 bacteria on, from herbivores, 310  
 bees and, mutualism of, 451, 452f  
 biodiversity of, 267–8, 268f  
 carbon in, 95  
 carbon : nitrogen ratio in, 93  
 cold tolerance of, 46, 46f  
 competition of, herbivores and, 308, 309f  
 constant final yield of, 137f  
 death of, defoliation and, 307  
 decomposers and, 370–1  
 defences of  
   against herbivores, 294–307, 296f–307f  
   apparency theory and, 298–9, 298f  
   carbon dioxide and, 306–7, 307f  
   fertilisers and, 305, 305f  
   of roots, 300, 301f  
   of shoots, 300, 301f  
   stress and, 305–6, 306f  
   timing of, 299–303, 299f–303f  
 dispersal of  
   invasion, 190–1, 190f  
   polymorphism and, 184, 184f  
 dormancy in, 112–14, 113f  
 fecundity of, herbivores and, 311–12, 311f  
 in four-trophic-level system, 545, 546f  
 fungi and, 36  
   from herbivores, 310  
   mutualism of, 407, 440–3, 441f, 441t  
 growth of, defoliation and, 307–8  
 herbivores and, coevolutionary arms race of, 295  
 at high temperature, 47–8  
 as hyperaccumulators, 60  
 inbreeding and outbreeding of, 182, 182f  
 as invasive species, 24f  
 life cycles of, 110  
 life histories of, 210, 210f  
   fast–slow continuum for, 228–9, 229f  
 litter decomposition, home field advantage for, 367–8, 368f, 369f  
 at low temperature, 44–5  
 macroparasites of, 381  
 as modular organisms, 102–4, 103f  
   intraspecific competition of, 137f  
 natural selection in, 2  
 nitrogen and, interspecific competition for, 242–3, 242f  
 nutritional contents of, 93–5, 94f  
 optimal foraging theory for, 343–5, 344f  
 oxygen for, 90–1, 91f  
 parasite and, competition of, in salt marshes, 402, 402f  
 parasite host defences of, 385  
 phalanx-type, 179, 179f  
 physical defences of, 295  
 population boundary lines for, 164–5, 164f, 166f  
 rank–abundance diagram for, 576f  
 in Raunkiaer’s life forms, 28–9, 28f  
 reproductive investment in, 218  
 reproductive value of, 209f  
 salinity and, 57–8, 57f  
 secondary metabolites of, 295–7  
 semelparity in, 222f  
 SESs of, 533  
 sodium chloride for, 32  
 species distribution of, temperature and, 49–54, 50f–3f  
 species richness of, 609, 609f, 610f  
   grazers and, 536–8, 536f–8f  
 subpopulations of, 200, 200f  
 survivorship of, herbivores and, 308–10  
 tolerance of  
   of herbivores, 307–14, 308f–13f  
   for invasion, 308, 308f  
 in tropical rainforest, 27  
 in tundra, 25  
 viruses on, from aphids, 310, 310f
- Plastics, 731–2, 732f
- PLFA. *See* Phospholipid fatty acids
- POC. *See* Particulate organic carbon
- Pollination and pollinators  
 chain of extinction of, 510–11, 511f  
 insects as, as keystone species, 553  
 mutualisms of, 429–32, 430f
- Pollution, 59–61, 60f, 730–3, 731f, 733f  
 from agriculture, stream invertebrates and, 29  
 habitat loss from, 507–8  
 industrial melanism and, 7–8, 7f  
 sodium chloride from, 32
- Polymorphism  
 dispersal, 184–5, 184f, 186f

- of ecotypes, 5–8, 6f
- in fish, 14
- Polyphagous foragers, 285
- POP. *See* Persistent organic pollutant
- Population. *See also* Abundance; Metapopulation
  - boundary lines, 164–5, 164f, 166f
  - cycles, 466–74, 466f–9f, 472f, 473f
  - defined, 107
  - experimental perturbation of, 459–60
  - growth rate of, 455–6, 458–9, 459f, 459t, 461
  - modular, 104
  - MVPs, 511–12
  - natural selection and, 2
  - of prey, predation and, 316–19, 317f–19f
  - resilience of, 29
  - stability of
    - in food webs, 555–6
    - species richness and, 560, 561f
  - variability of, species richness and, 556–7, 556f
- Population dynamics
  - of infection, 402–10, 405f, 406f, 409f, 410f
  - of parasite hosts, 410–15, 411f–13f, 415f
  - of predation
    - aggregated distributions in, 339–41, 340f
    - consumer–resource models of, 320–5, 322f, 323f
    - coupled oscillations in, 286f
    - crowding and, 334
    - delayed density dependence of, 323, 323f
    - direct density dependence of, 323–4
    - food quality and, 330, 330f, 331t
    - of heterogeneity, 349–55, 352f–5f
    - interference and, 330–9, 332f, 333f, 335f, 336f, 337t, 338f
    - intimidation in, 331, 338–9, 338f
    - Lotka–Volterra model for, 320–2, 322f, 350
    - Nicholson–Bailey model of, 324–5, 350
    - type 1 functional response in, 325–6, 326f
    - type 2 functional response in, 326–7, 326f, 337, 338f
    - type 3 functional response in, 327, 327f, 335–6, 336f
- Population projection matrices (PPMs), 123–7, 124f
  - for extinction, 513, 513f
- Population size, 107–8, 108f, 122
  - asymmetric competition and, 159, 159f
  - as extinction risk, 503–4
  - intraspecific competition and, 141–8, 141f–6f
  - IUCN on, 709–10
  - territoriality and, 159–61, 160f, 161f
- Population viability analyses (PVAs), 512–14, 512f
- Postbreeding dispersal, 170
- Potassium, 87, 90
- Potential evapotranspiration (PET), 613–14, 614f
- PPFD. *See* Photosynthetic photon flux density
- PPMs. *See* Population projection matrices
- Praziquantel, 408
- PRCF. *See* Partial rate correlation function
- Predation, 284–319
  - abundance patterns and, 285
  - in community modules, 534–40, 534f–9f
  - compensatory reactions to, 316–17
  - defined, 284
  - food and, 317, 317f
  - frequency-dependent, 538–9, 538f, 539f
  - by herbivores, 285
  - population dynamics of, 320–56
    - aggregated distributions in, 339–41, 340f
    - consumer–resource models of, 320–5, 322f, 323f
    - coupled oscillations in, 286f
    - crowding and, 334
    - delayed density dependence of, 323, 323f
    - direct density dependence of, 323–4
    - food quality and, 330, 330f, 331t
    - of heterogeneity, 349–55, 352f–5f
    - interference and, 330–9, 332f, 333f, 335f, 336f, 337t, 338f
    - intimidation in, 331, 338–9, 338f
    - Lotka–Volterra model for, 320–2, 322f, 350
    - Nicholson–Bailey model of, 324–5, 350
    - type 1 functional response in, 325–6, 326f
    - type 2 functional response in, 326–7, 326f, 337, 338f
    - type 3 functional response in, 327, 327f, 335–6, 336f
  - ratio dependent, 331
  - realised niche and, 294
  - of seeds, 310
  - specialist model for, 470–1, 472f, 473
  - on weakest prey, 317
- Predator-mediated coexistence, 535–7
- Predator–prey relationship
  - coevolution of, 292, 295
  - coupled oscillations of, 206f
  - cycles, 325
  - heterogeneity in, 350–1
  - pesticide biomagnification in, 480–1
  - population cycles for, 467–74, 467f–9f, 472f, 473f
  - zero isocline for, 474, 474f
- Predators
  - aggregated distributions of, density of prey and, 349–50
  - apparent competition of, 270–1, 271f
  - community size spectra and, 577
  - endotherms and, 44
  - fish reproductive allocation and, 220, 221f
  - food preferences of, 286–8, 287f
  - food web trophic cascades for, 544–7, 545f–7f
  - foraging in presence of, 292–4, 293f
  - intimidation of, 318–19, 319f

- Predators (*cont'd*)
- isoclines of, with mutual interference, 334
  - learning by, 349
  - in mesopredator release, 547, 547f
  - migration of, 349
  - optimal foraging theory for, 289–92, 290f
  - parasite host as, 384
  - polymorphism and, 5
  - population cycles and, 466
  - resources of, 91–2
  - saprotrophs and, 65
  - of seeds, 310
    - satiation of, 327–30, 328f, 329f
  - selfish herd and, 181
  - specialisation of, 292
  - switching of food of, 288–9, 289f
  - true, 284
  - types of, 284–5
- Pre-emptive competition, 263–9
- Prey
- apparent competition of, 270
  - biomass of, 160f, 670, 670f
  - density of, predator aggregate distributions and, 349–50
  - endotherms and, 44
  - enemy-free space of, 607
  - isoclines of, with intraspecific competition, 332–4, 333f
  - population of, predation and, 316–19, 317f–19f
  - weakness of, 317
- Primary productivity. *See also* Gross primary productivity; Net primary productivity
- in aquatic biomes, 664–7, 664f–8f
  - energy flux and, 649–56, 649t, 650f–5f
  - grazers and, 670–1
  - by herbivores, 669, 670f
  - ocean acidification and, 723f
  - in oceans, SST and, 666
  - secondary productivity and, 668, 669f
  - in terrestrial biomes, 656–63, 656f–63f
- Primary succession, 582–5, 583f
- phosphorus in, 680–1, 681f
- Production efficiency, 673
- Productive space, food chain length and, 564, 568, 568f
- Productivity. *See also* Primary productivity
- beta ( $\beta$ ) diversity and, 612
  - EEH and, 550
  - energy flux and, 648
  - food chain length and, 564, 565–6, 566f
  - of food webs, 551, 552, 557
  - gamma ( $\gamma$ ) diversity and, 612, 613f
  - latitude and, 651
  - of oceans, 651
  - secondary, 668, 669, 669f, 723f
  - species richness and, 608–15, 609f, 610f, 612f–15f, 661–3, 661f, 662f
- Profit sharing, in harvest management, 493, 493f
- Protected areas (PAs)
- agriculture and, 611
  - for biodiversity conservation, 604–5, 604f
  - for corals, 604f
  - establishing and managing, 724–5, 725f
  - IPs in, 737, 737f
  - MPAs, 637–8, 637f
- Protected landscapes/seascapes, 605
- Protists, 4, 41f, 42, 100, 100f, 361
- Protozoans, 371–2, 379
- beetles and, 411f
  - as detritivores, 361
  - in gut mutualism, 433
  - infections, prevalence of, 405
  - interspecific competition of, 282–3, 283f
- Provisioning services, 58, 502
- Pseudo-interference, 350
- PVAs. *See* Population viability analyses
- Quantitative trait loci (QTLs), 47
- life history trade-offs and, 213
- R. *See* Fundamental net reproductive rate
- RA. *See* Autotrophic respiration
- Rabbits
- as keystone species, 553
  - myxomatosis and, 393–4, 394f
- Rabies, 384
- immunisation for, 406–7
- Radiative forcing, 711–12, 712f
- Ramets, 106
- Random dispersal, 180, 180f
- Random encounters, in predation population dynamics, 324
- Random fraction (RF), 576, 576f
- Rank–abundance diagrams, 501f, 530, 575–7, 576f, 601
- Rarefaction, for species richness, 606, 607f
- Rates of increase, 121–7
- Ratio-dependent isoclines, 334
- Ratio-dependent predation, 331
- Raunkiaer's life forms, 28–9, 28f
- RBP. *See* Rights-based fishery
- RCMs. *See* Regional climate models
- RCPs. *See* Representative concentration pathways
- RDZ. *See* Resource-depletion zone
- RE. *See* Ecosystem respiration

- Reaction–diffusion modelling, 193–4, 193f
- Realised niche, 36, 244, 294
- Recolonisation, metapopulations and, 197–203
- Redfield ratio, 95, 665
- Refection, 433
- Refuges, 350
- Regime shifts, in food webs, 568–71, 568f–71f
- Regional climate models (RCMs), 711
- Regular dispersal, 180, 180f
- Regulating services, 58, 502
- Regulation
  - of abundance, 454–5, 455f, 477–8
  - in predation population dynamics, 324
- Relative growth rate (RGR), 80f
- Remoteness of islands, 624, 624f
- Reorganisation algorithms, 529–30
- Representative concentration pathways (RCPs)
  - carrying capacity and, 711–12, 712f
  - for temperature, 712, 713f
- Reproduction rates, 121–7
  - of helminths, 407
  - population cycles and, 466
- Reproductive allocation
  - age of maturity and, 220
  - in fish, predators and, 220, 221f
  - life histories and, 208, 208f, 218–20, 219f
- Reproductive investment, 218
  - semelparity and, 220–1, 222f
- Reproductive strategy axis, 228
- Reproductive value, 208–10, 209f
- Reptiles. *See also* Lizards
  - biodiversity conservation of, 15
  - dispersal of, by sex, 185–6, 186f
  - in extreme temperatures, 53f
  - as invasive species, 24f
  - population size of, 709, 710f
- Residual reproductive value (RRV), 208, 209
  - clutch size and, 224
  - CR and, 213, 217
  - options sets and, 216
- Resilience
  - of food webs, 552
  - of populations, 29
- Resistance
  - of food webs, 552
  - mean, virulence and, 395f
  - of parasite hosts, 388, 389f
  - to pesticides, 481–2, 482f
- Resistant diapause, 112
- Resource availability hypothesis, 304
- Resource-depletion zone (RDZ), 67, 76
  - exploitation and, 133
  - nitrogen and, 88
  - phosphorus and, 89
- Resources, 65–101. *See also* Carbon dioxide; Food; Minerals; Productivity; Solar radiation; Water
  - of animals, 92–3
  - classification of, 95–7
  - competition for, of parasite host, 388–92, 389f–92f
  - consumer–resource models, of interspecific competition, 253–61, 254f, 256f, 258f–60f
  - of decomposers, 357
  - of detritivores, 357
  - detritivores and, 371–7, 371f–5f, 376t
  - environmental space and, 169
  - of grazers, 92
  - growth and, niche differentiation and, 255
  - life histories and, 210–14, 210f–12f, 214f
  - lifespan and, 93
  - migration between, 172
  - movement and, 169
  - of niches, 97
  - oxygen as, 90–1, 91f
  - parasite host as, 384
  - of parasites, 92
  - of predators, 91–2
  - self-thinning and, 166–7, 167f
  - species richness and, 607
  - transport of, 98
- Resource-transport network theories (RTN), 99, 101
- Resource utilisation curves, 262–4, 263f
- RF. *See* Random fraction
- RGR. *See* Relative growth rate
- RH. *See* Heterotrophic respiration
- Rights-based fishery (RBP), 735, 735f
- Risk. *See* Aggregation of risk
- Rivers
  - acid rain and, 60
  - nitrogen in, 727–8
  - phosphorus in, 727–8
- Robustness, of food webs, 552, 559–60, 560f
  - parasites in, 567, 568f
- Roots
  - competition of, 158, 158f
  - fungi and, 88–90
    - mutualism of, 440–3, 441f, 441t
  - interspecific competition of, 260, 260f
  - minerals and, 87–8
  - plant defences of, 300, 301f
  - water and, 76–7, 78f

- RRV. *See* Residual reproductive value
- r*-selection, 226–7, 228f, 238
- RTN. *See* Resource-transport network theories
- Ruderal strategy, 232
- Ruminants, gut mutualism in, 432–3, 433f, 434t
- Rusts, 381, 395f
- SA. *See* Salicylic acid; Surface area theories
- Salicylic acid (SA), 299–300, 300f
- Salinity, 56–8, 56f, 57f
- Salt marshes
  - decomposition in, 365–6, 366f
  - NPP of, 649t
  - oxygen in, 90–1
  - plant parasite competition in, 402, 402f
- Sample-based data, on species richness, 606
- Saprotrophs. *See also* Decomposition and decomposers; Detritivores
  - predators and, 65
- Satiation of seed predators, 327–30, 328f, 329f
- Scaling coefficient, 556
- SCFAs. *See* Short-chain fatty acids
- Schistosomiasis, 378, 407–8
  - macroparasites and, 380
- Scramble, 140–1
  - in population models, 151
- SDMs. *See* Species distribution models
- Searchers
  - foragers as, 290
  - parasitoids as, 324
- Seasonal environments
  - GPP in, 652
  - niche differentiation in, 618
  - NPP in, 659, 659f
- Sea surface temperature (SST), 42f
  - fish and, 735
  - NPP and, 666, 666f
- Secondary metabolites, of plants, 295–7
- Secondary pest outbreaks, 479–80, 480f
- Secondary productivity, 668, 669, 669f, 723f
- Secondary succession, 585–6
- Seed banks, 110–11, 111f
  - for forests in wetlands, 112
  - in succession, 585
- Seeds
  - bats and, 588
  - birds and, 93
  - dispersal of, 170, 175–6, 175f, 177, 177f
    - mutualism of, 428–9
  - dormancy of, 112–14, 113f
  - life cycles and, 110
  - life histories and, 206
  - movement of, 169
  - parasites on, 286
  - predators of, 310
    - satiation of, 327–30, 328f, 329f
  - species richness and, 609
  - in succession, 584–5, 584f
  - temperature and, 48–9
- Selection. *See also* Natural selection
  - in communities, 572
  - complementarity, 640–1, 641f
  - in food webs, 557
  - K*-selection, 226–7, 228f, 238
  - in metacommunities, 600
  - r*-selection, 226–7, 228f, 238
  - species richness and, 662–3, 662f
- Selfish herd, 181
- Self-thinning, 161–8, 163f, 164f, 166f–8f
- Semelparity, 108–11, 220–1, 222f
- Senescence, 104–6, 105f
- Series replacement model, of succession, 586, 586f
- SEs. *See* Standardised effect sizes
- Sexually transmitted diseases, 404, 411, 412f
- Shade plants, 73–4, 73f, 74f
- Shannon's diversity index, 575
- Shoots
  - interspecific competition of, 260, 260f
  - plant defences of, 300, 301f
- Short-chain fatty acids (SCFAs), 432
- Short-lived agents, 396
- Shredders, 364
- Sickle-cell anaemia, 5
- Sigmoid growth curves, 143, 144f
- Simpson's diversity index, 574
- Single populations, dispersal of, 186–95, 187f–190
- S-I-R model, for acquired immunity, 385–6
- S-I-S model, for acquired immunity, 385–6
- Size. *See also* Population size
  - asymmetric competition and, 155–8, 156f–8f
  - community size spectra and, 577
  - of decomposers, 361, 362f
  - of ecosystem, food chain length and, 564, 568f
  - life histories and, 207, 207f
  - metabolic scaling and, 99f
  - spectra, for communities, 577–8, 577f
  - temperature and, 39–41, 42f
- SLA. *See* Specific leaf area
- Small populations, 502–5
- Smuts, 381
  - as sexually transmitted disease, 411, 412f

- Snails  
 character displacement of, 280, 280f  
 physical defences of, 315–16, 315f  
 plant defences and, 301–2, 301f  
 polymorphism of, 6, 6f
- Sociality, dispersal and, 181
- Social parasitism, 381
- Sodium chloride, 32
- Soil  
 pH of, 54–6, 55f  
 texture, NPP and, 658–9  
 water in, 55, 55f, 76–7, 77f
- Solar radiation, 66–74, 66f, 68f, 69f. *See also* Photosynthetically active radiation  
 in aquatic biomes, 27  
 carbon dioxide and, 62  
 in lakes, 71f  
 photoautotrophs and, 65  
 in tropical rainforest, 27
- Spatial heterogeneity, species richness and, 616, 617f
- Specialisation  
 community size spectra and, 577  
 in non-seasonal environments, 618  
 of predators, 292  
 within species, 2–8
- Specialist predation model, 470–1, 472f, 473
- Speciation  
 allopatric, 8–14, 9f, 10f, 12f–13f  
 in communities, 572  
 on islands, 18–19, 18f
- Species. *See also specific topics*  
 defined, 8  
 environmental conditions and, 32–64  
 geographic variation within, 3–5, 3f, 5f  
 number of, 501–2, 501f  
 reintroduction of, species distribution modelling for, 195, 196f  
 specialisation within, 2–8  
 turnover of, species richness and, 625–6, 625f, 626f
- Species–abundance distributions, 530
- Species accumulation curve, 606, 607f
- Species–area relationship, 619, 619f
- Species distribution. *See also* Aggregated distributions  
 of animals, temperature and, 49–54, 50f–3f  
 global warming and, 22, 64  
 historical factors in, 16–24, 16f–23f, 24t  
 modelling of, for invasion dispersal, 195, 196f  
 of plants, temperature and, 49–54, 50f–3f
- Species distribution models (SDMs), 714
- Species richness  
 biodiversity and, 603–5  
 in communities, 574–5, 574f, 592  
 complementarity effects and, 662–3, 662f  
 data on, 606  
 disturbance and, 608  
 energy and, 613–16, 614f–16f  
 evolutionary time and, 618  
 extrapolation of, 606–7, 607f  
 facilitation and, 608  
 factors in, 605–6  
 of fish, 559, 559f  
 gradients of  
 elevation as, 633–6, 634f–6f  
 latitude as, 631–3, 632f, 633f  
 succession as, 638–9, 639f  
 in grasslands, 558f  
 habitat area and, 619, 619f  
 harsh environments and, 417, 616  
 hot spots of, 603, 640, 735  
 on islands, 619–31, 619f–31f  
 macroecology and, 605  
 in metacommunities, 601  
 model of, 607–8, 608f  
 niche overlap in, 607  
 pH and, 616, 617  
 of plants, grazers and, 536–8, 536f–8f  
 population stability and, 560, 561f  
 productivity and, 608–15, 609f, 610f, 612f–15f, 661–3, 661f, 662f  
 rarefaction for, 606, 607f  
 remoteness of islands and, 624, 624f  
 resources and, 607  
 selection and, 662–3, 662f  
 spatial heterogeneity and, 616, 617f  
 species turnover and, 625–6, 625f, 626f  
 of vertebrates, 710f
- Species-sorting metacommunity, 600
- Specific leaf area (SLA), 232
- Spiders  
 dispersal of, 175  
 food chain length of, 566  
 in four-trophic-level system, 545  
 frequency-dependent predation of, 539f
- Sponges  
 chemical defences of, 315f, 316  
 as modular organisms, 102  
 specialisation of, 3
- Squirrels  
 apparent competition of, 273–4, 273f  
 habitat restoration for, 188–9, 189f  
 predator intimidation of, 318–19, 319f
- SST. *See* Sea surface temperature

- Stability
- of abundance, 454–6, 454f
  - alternative states of, for abundance, 476
  - food chain length and, 564–5
  - of food webs, 552–7
    - biodiversity and, 401
    - complementarity effects and, 559, 559f
    - from fish, 559, 559f
  - of forests, 21f
  - generalist predators switching and, 336–7, 337f, 337t
  - heterogeneity and, 268
  - of population
    - in food webs, 555–6
    - species richness and, 560, 561f
  - in predation population dynamics, 324
  - suppression and, trade-off of, 334
- Standardised effect sizes (SEs), 533
- Static life tables, 118–20, 120f
- Stochastic colonisation, of patches, 543
- Stochastic models
- for communities, 541–3
  - of intraspecific competition, 153–4, 154f
  - of patch occupancy, 515
- Stochastic processes, in metacommunities, 600
- Strange attractors, 153
- Streams
- decomposers in, 365
  - detritivores in, 364–5, 365f
  - invertebrates in, agricultural pollution and, 29, 29f
  - nutrient loss in, 685
  - nutrient spiraling in, 689–90, 690f
  - PAR in, 664, 664f
  - velocity of, in niches, 33
- Stress
- as abiotic factor, 419
  - on habitats, 232
  - plant defences and, 305–6, 306f
- Stress-gradient hypothesis, 419, 540–1, 542f
- Strict nature reserves, 604
- Subpopulations, 192–5
- of bees, 197, 198f
  - of plants, 200, 200f
- Succession
- animals in, 587–8
  - to climax, 588–9
  - competition in colonisation and, 582
  - facilitation in, 587, 590–1
  - in grasslands, 585, 585f
  - nitrogen-fixing bacteria and, 450–1
  - NPP in, 652, 661
  - primary, 582–5, 583f
  - phosphorus in, 680–1, 681f
  - restoration and, 589–91, 590f, 591f
  - secondary, 585–6
  - seed banks in, 585
  - seeds in, 584–5, 584f
  - series replacement model of, 586, 586f
  - as species richness gradient, 638–9, 639f
  - vital attributes in, 568
- Successional mosaics, 592, 598, 598f
- Successional niche, 587, 587f
- Sulphur, 87, 91
- cycle, 703f, 704–5
- Sulphur dioxide
- in acid rain, 60, 60f
  - industrial melanism and, 8
- Superspreaders, 398, 398f
- Supply point resources, coexistence and, 255
- Supporting services, 58, 502
- Surface area theories (SA), 99, 101
- Survivorship, 114
- CR habitats and, 217
  - curves, 117–18, 118f
    - of mammals, 119–20, 119f
  - of grasshoppers, 317
  - growth and, life history trade-offs for, 212, 212f
  - interspecific competition and, 240
  - of parasite hosts, 399–401
  - of plants, herbivores and, 308–10
  - in population projection matrices, 124
- Sustainability, 477–8
- biodiversity conservation and, 642
  - in harvest management, 491, 497–8, 498f
  - of IFS, 485–6, 487f
- Symbiosis, 416. *See also* Commensalism; Mutualism; Parasites
- fitness of host and, 421–2
  - of insects and bacteria, 436–7
  - of photosynthesis, of invertebrates and algae, 438f
- Taiga, 25f, 26
- Tannins, 298
- Target pest resurgence, 479, 480f
- Temperate forests, 25f, 26
- carbon dioxide in, 81–2
  - NPP of, 649t
- Temperature. *See also* Global warming; Sea surface temperature
- carbon dioxide and, 81–2, 81f
  - crops and, 55, 55f
  - development and, 39–41, 41f
  - extremes of, 51–4, 52f, 53f
  - growth and, 39–41, 41f

- high, 46–8  
 low, 44–5, 45f  
 metabolism and, 39–41, 40f  
 in niches, 33, 39–56, 40f–3f, 45f–7f, 50f–3f, 55f  
 normalisation constant and, 98  
 NPP and, 658, 658f  
 photosynthetic capacity and, 71  
 RCPs for, 712, 713f  
 size and, 39–41, 42f  
 species distribution and, 49–54, 50f–3f  
 as stimulus, 48–9
- Terminal investment hypothesis, 218–20, 219f
- Terrestrial biomes/communities/environments, 24–7, 25f  
 detritivores in, 363–4  
 nutrient budgets in, 680f, 682–9, 683f–6f, 689f  
 primary productivity in, 656–63, 656f–63f
- Terrestrial landscape planning, 644–6, 645f
- Territoriality  
 asymmetric competition for, 159–61, 160f, 161f  
 benefits and costs of, 160  
 of vultures, 162, 162f
- Territorial use rights in fisheries (TURFs), 736
- Thermoneutral zone, 43–4
- Time lags, in population models, 151
- Time series, of abundance, 460–5, 462f–5f
- Toads  
 allelopathy of, 248  
 migration of, 172
- Tolerance  
 to cold, 45–7, 46f, 47f  
 of parasite hosts, 388, 389f  
 of plants, to herbivores, 307–14, 308f–13f
- Top-down food webs, 548–9, 549f
- Trade-offs  
 with biodiversity conservation, 642  
 with competition  
 colonisation and, 586–7, 599, 600f  
 commensalism and, 418  
 with costs of reproduction, 210f  
 in grasses, 257, 259f  
 in life histories, 210–14, 210f–12f, 214f  
 with CR, 210f, 213  
 fitness contours and, 216, 217f  
 genotypes and, 212–13  
 with offspring size and number, 213–16, 214f, 215f, 221–6, 222f–5f  
 options sets and, 214–15, 217f  
 for survivorship and growth, 212, 212f  
 with parasite host defences, 386f, 387f  
 with suppression and stability, 334  
 with transmission and virulence, 394
- Transgressive overyielding, 557, 662
- Transmission  
 density-dependent, 396, 397f  
 frequency-dependent, 397, 397f, 404  
 in parasite hosts, 396–402, 397f–401f  
 of parasites, trade-offs with, 394  
 threshold, of infection, 403
- Transpiration, 74
- Transport, of resources, 98
- Trophic cascades  
 in food web, in mesopredator release, 547, 547f  
 in food webs, 544–51, 545f–51f  
 keystone species and, 553–4, 554f
- Trophic compartments, for energy flux, 668–71, 671f
- Trophic loops, in food webs, 561–4, 563f
- Trophic transfer efficiency, 673–4, 673f
- Tropical rainforest, 25f, 27  
 NPP of, 649t  
 PAR in, 67
- Tropics  
 climate change in, 53  
 species richness in, 632–3, 633f
- True predators, 284, 285
- Tsunamis, 58
- Tuberculosis, 378  
 BTB, 387, 387f
- Tundra, 25, 25f  
 niche complementarity in, 527–8, 528f  
 NPP of, 649t  
 species richness in, 616, 617f
- TURFs. *See* Territorial use rights in fisheries
- Turnover dispersal, 180
- Type 1 functional response, in predation population dynamics, 325–6, 326f
- Type 2 functional response, in predation population dynamics, 326–7, 326f, 337, 338f
- Type 3 functional response, in predation population dynamics, 327, 327f, 335–6, 336f
- Typhoid Mary, 398
- Ultraviolet radiation, 730, 730f
- Undercompensating density dependence, 135
- Underexploitation, 487–8
- Unified Neutral Theory of Biodiversity and Biogeography, 530
- Uninhabited habitable patches, metapopulations and, 195–7
- Unitary organisms, 102, 109f
- Upwelling, in oceans, 665–6, 668f
- Vertebrates  
 as detritivores, 376–7

- Vertebrates (*cont'd*)
- gut of
    - metagenome in, 433–4, 435f
    - mutualism in, 433–4, 433f, 434t
  - parasite host defences of, 385
  - SEs of, 533
  - species richness of, 710f
- Virulence
- of bacteria, 395, 395f
  - evolution of, 392–5
  - mean resistance and, 395f
  - parasite hosts and, 388, 389f
  - of parasites, trade-offs with, 394
- Viruses
- as microparasites, 379
  - on plants
    - from aphids, 310, 310f
    - from herbivores, 310
- Vital attributes, in succession, 568
- Vital rates, 102
- VORTEX model, 514, 514f
- Vulnerable species, 711
- Wallace, Alfred Russell, 1–2, 2f
- WASH. *See* Water, sanitation, and hygiene education
- Water
- conservation of, photosynthesis and, 74–5, 76f
  - high temperature and, 47
  - hydrological cycle for, 699–701, 699f, 701f
  - as inhibitory resource, 97
  - NPP and, 657, 657f, 658f, 660f
  - photoautotrophs and, 65
  - roots and, 76–7, 78f
  - in soil, 55, 55f, 76–7, 77f
- Water, sanitation, and hygiene education (WASH), 408
- Weather, density dependence and, 465
- Wetfall, 683–4, 684f
- Wetlands
- catchment management in, 729, 729f
  - forests in, seed banks for, 112
  - nitrates and, 695–6, 696f
- White syndromes, 731
- Wilderness areas, 604–5
- Wind, dispersal by, 176–7
- Yield–effort curve, 490, 490f
- Yield-per-recruitment, 496, 496f
- Y model, for life history trade-offs, 211, 211f
- Zero-deforestation policy, 720
- Zero isoclines, 250, 250f, 474, 474f
- Zero net growth isoclines (ZNGI), 95–96, 96f, 253–4, 254f
- Zeta ( $\zeta$ ) diversity, 575
- Zinc, 60, 87
- ZNGI. *See* Zero net growth isoclines
- Zonation, 57–8
- of marine sediments, 359, 359f
- Zoonotic infections, 384











