

1

Historical Background of Atmospheric Secondary Aerosol Research

1.1 Introduction

Trace components in the tropospheric atmosphere consist of gaseous molecules and particulate matters. Most of gaseous molecules in the atmosphere do not have absorption bands in the visible region. Some species such as ozone and nitrogen dioxide have the absorption, but they are invisible to the naked eye under the normal atmospheric conditions because their absorbance are small. In contrast, since the particulate matters intercept sunlight and small particles scatter strongly the solar radiation, they are captured easily by the naked eye as haze. Thus, particulate matters in the atmosphere called atmospheric aerosols have been studied from relatively early days in relation to air pollution historically.

These atmospheric aerosols are divided broadly into the primary species released directly from emission sources and the secondary compounds formed by chemical reactions in the atmosphere. Further, secondary particulate matter can be classified into secondary inorganic aerosol and secondary organic aerosol (SOA).

This book aims at the understanding of chemical reactions forming secondary aerosols in the gas phase, in the liquid phase, and at their interface, particularly focusing on organic aerosols. Therefore, most of the descriptions are focused on organic species, and inorganic species are addressed whenever necessary. As for the formation of secondary inorganic aerosols, detailed discussion has been given by the textbook of Seinfeld and Pandis (2016).

In this chapter, historical background of research on atmospheric secondary aerosols, including inorganic aerosols, is described looking back before 1980s, when the atmospheric chemistry was founded as one of the academic fields of the global environmental sciences.

1.2 Secondary Inorganic Aerosols


The first capture of atmospheric secondary inorganic aerosol such as nitrate and sulphate was in the form of precipitation component, and their historical reviews are available by Eriksson (1952a, 1952b) and Möller (2008). First discovery of nitrate in precipitation was made by Marggraf (1751), a German chemist, and mineral species (silica and lime), sea salt component (sodium and chloride), ammonium, and organics as brown residue were also detected together with nitrate. It was the earlier half of

nineteenth century when Liebig (1835) advocated a theory that atmospheric nitrogen compounds deposited on ground are essential to plant growth as nutrient salt absorbed by roots, leading to a revolution of agricultural chemistry. Thus, atmospheric nitrate, the main component of the plant nutrient, had been discovered from long ago as a precipitation constituent (Miller 1905; Eriksson 1952a; Möller 2008). On the other hand, the discovery of sulphate was delayed nearly 100 years after that of nitrate. From the view point of air pollution in Manchester, UK, Smith (1852) described based on the analysis of precipitation that three kinds of air can be found: (i) with carbonate of ammonia in the remote field; (ii) with sulphate of ammonia in the suburbs; and (iii) with sulfuric acid in the urban area (Cowling 1982). The described ammonium carbonate ((NH₄)₂CO₃), ammonium sulfate ((NH₄)₂SO₄), and sulfuric acid (H₂SO₄) are formed secondarily by the chemical reactions in the gas phase or in the fog water from atmospheric trace gaseous species, CO₂, NH₃, and SO₂. These aerosols are water-soluble, and recognized as major components of “acid rain” after taken into precipitation. Incidentally, the term of *acid rain* was used for the first time in the monograph of Smith (1872) as accredited by Cowling (1982). Since then, the measurement of nitrate and ammonium had been made in many places in Europe in the latter half of nineteenth century from the interest of agricultural chemistry, while sulfate had been measured in the eastern part of United States since the 1910s (Cowling 1982).

Hydrogen ion concentration (pH) has been measured since the 1950s, started in Europe and United States, over a wide area. Owing to these wide-area observations, spatial distribution and temporal trends of pH and chemical components of precipitation became to be known well in Europe (Emanuelsson et al. 1954; Barrett and Brodin 1955; Odén 1976) and North America (Junge and Werby 1958; Gorham and Gordon 1960; Cogbill 1976). The acid rain causing acidification of lakes and rivers and their impact on fishery was then brought up as a social problem internationally. The quantitative research on the formation of sulfate and nitrate as secondary inorganic aerosol had been developed rapidly as “acid rain” became social concern.

1.2.1 Sulfate

In the earlier studies on acid rain, it was thought that sulfur dioxide (SO₂), primary air pollutants whose atmospheric concentration had increased rapidly after the Industrial Revolution, was taken up into fog water droplets and converted to sulfate by oxidation in the aqueous phase (Junge and Ryan 1958; Junge 1963):



The rate limiting stage of this process is the oxidation step of SO₃²⁻ to SO₄²⁻, and the oxidation by O₂ had been studied for a long time (Fudakowski 1873; Backstrom 1934). However, the oxidation rate of SO₃²⁻ by O₂ was found to be very slow (Fuller and Crist 1941; Brimblecombe and Spedding 1974). Therefore, this reaction is not important for O₂ alone as the oxidation reaction of SO₂ in the atmosphere, but it was found that the reaction is accelerated by the coexistence of trace metal ions such as Fe³⁺, Cu²⁺, and Mn²⁺ (Reinders and Vles 1925; Junge and Ryan 1958; Brimblecombe and Spedding 1974; Hegg and Hobbs 1978). The effects of transition metal ions on the SO₂ oxidation in the aqueous phase still leaves a lot of unknowns, and the studies are ongoing (Deguillaume et al. 2005; Harris et al. 2013; Herrmann et al. 2015).

In 1970s, the importance of the reaction of O_3 and H_2O_2 , formed secondarily in the photochemically polluted atmosphere, was pointed out. The pioneering studies were made by Penkett and Garland (1974), Erickson et al. (1977), and Larson et al. (1978) for O_3 , and by Mader (1958), Hoffmann and Edwards (1975), and Penkett et al. (1979) for H_2O_2 . Later studies on these aqueous phase reactions revealed that the oxidation by H_2O_2 is more important at lower pH than 7, and those of O_3 become important in the higher pH region. Details of these aqueous-phase reactions are summarized in the textbooks by Akimoto (2016, pp. 363–372), and Seinfeld and Pandis (2016).

Atmospheric oxidation reactions of SO_2 to SO_4^{2-} were studied earlier for the aqueous-phase reactions, and the gas-phase reactions was noted later by Cox and Penkett (1972). The years of 1970s are the era that OH radical chain reactions were proposed and demonstrated to cause photochemical air pollution (Akimoto 2016, pp. 288–290). The importance of the reaction of SO_2 and OH for the oxidation of SO_2 was deduced based on the measured rate constant of the reaction (Eggleton and Cox 1978; Davis et al. 1979). Later, Stockwell and Calvert (1983) showed the oxidation process of SO_2 with OH as



This implies that the $HOSO_2$ forms H_2SO_4 without terminating the OH chain reaction. The SO_2 oxidation mechanism in the gas-phase has thus been established. Although the relative importance of gas- and aqueous-phase reactions varies widely, depending on meteorological conditions. It is thought in general that both processes are important (Barrie et al. 2001). Most of sulfate in particles exist as ammonium sulfate ($(NH_4)_2SO_4$) or ammonium bisulfate (NH_4HSO_4), and a part of them exists as sulfuric acid (H_2SO_4) when NH_3 is in short stoichiometrically as observed in the sub-micron particles in many urban samples (van den Heuvel and Mason 1963; Ludwig and Robinson 1965; Wagman, Lee, and Axt 1967).

1.2.2 Nitrate

The measurement of nitrate (NO_3^-) in precipitation has been reported in United States early in 1920s from the interest in agricultural chemistry (Wilson 1926). Its atmospheric concentrations increased rapidly, accompanying with the rapid increase of fossil fuel combustion. It has been monitored since the 1950s as an important secondary inorganic aerosol next to SO_4^{2-} (Junge 1954; Lee and Patterson 1969). For example, the equivalent-basis fractions of SO_4^{2-} and NO_3^- in precipitation in Eastern United States in early 1960s are reported as ca. 60% and ca. 20%, respectively (Likens and Bormann 1974). Particularly, large amounts of nitrates were reported, together with sulfate and organic aerosols existing in photochemical smog mentioned in the next section (Renzetti and Doyle 1959; Lundgren 1970; Appel et al. 1978).

Since the rate constant of the reaction:



is one order of magnitude larger than the reaction, $\text{OH} + \text{SO}_2 + \text{M}$, under the atmospheric conditions, and the Henry's law constant of NO_2 is two orders of magnitude smaller than SO_2 (Table 2.2), nitric acid (HNO_3) in the atmosphere is thought to be formed in the gas phase and then taken into the aqueous phase (Orel and Seinfeld 1977). Meanwhile, a formation pathway other than (1.6) is considered to be the hydrolysis of N_2O_5 formed via NO_3 by the reaction of O_3 and NO_2 (Orel and Seinfeld 1977):



The rate constant of Reaction (1.9) in the gas phase as a homogeneous reaction is very small, $<2.0 \times 10^{-21} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Burkholder et al. 2015), and the heterogeneous reaction on the particle surface is thought to be more important (Mozurkewich and Calvert 1988). The NO_3 radical involved in this reaction process has absorption bands in the visible region and photolysed easily by sunlight, so that the formation of HNO_3 by this heterogeneous reaction process is thought to be important in the nighttime (Richards 1983; Heikes and Thompson 1983).

The gaseous nitric acid, ammonia, and ammonium nitrate formed from them are thought to be in equilibrium:



and comparison between model estimate based on the thermodynamic parameters (Stelson et al. 1979) and field observation for the formation of nitrate have been made (Harrison and Pio 1983; Hildemann et al. 1984). Reaction (1.10) is reversible reaction, and the particulate NH_4NO_3 increases with the decrease of temperature, and thus the concentration ratio of nitrate is known to increase in winter and at dawn. Multiphase models that treat sulfuric and nitric acid simultaneously have been developed in 1980s (Bassett and Seinfeld 1983; Saxena et al. 1983).

Gaseous HNO_3 reacts with sea salt (NaCl) on the surface to give sodium nitrate by releasing HCl :



Although it has been presumed that the reaction causes the decrease of chlorine to sodium ratio in the sea salt in the vicinity of continents and brings the nitrate in coarse particles (Robbins et al. 1959), the reaction has been validated by laboratory experiments only after the latter half of 1990s (De Haan and Finlayson-Pitts 1997; Wahner et al. 1998). Thus, nitrates have the characteristics that they exist as NH_4NO_3 in submicron particles in the inland and as NaNO_3 in coarse particles (2–8 μm) in the coastal urban area (Lee and Patterson 1969; Cronn et al. 1977).

1.3 Secondary Organic Aerosols

Existence of organic materials in the precipitation was noted by Marggraf (1751) in the middle of eighteenth century, and they were more clearly identified as humic acid-like substances in the first half of the nineteenth century (Lampadius 1837; Möller 2008).

However, the trigger to wide concern on the particulate organic compounds was the discovery of carcinogenic polyaromatic hydrocarbons (PAHs) in the diesel exhaust and urban atmosphere in the middle of twentieth century (e.g. Waller 1952; Kotin et al. 1954; Stocks and Campbell 1955; Wynder and Hoffmann 1965). Further findings of many oxygenated compounds in the atmospheric aerosols were made in the photochemical smog.

1.3.1 Photochemical Smog

In the middle of the 1940s, new type of smog totally different from conventional air pollution due to SO₂, sulfate, and coal fly ash, spread in Los Angeles basin, and became a social problem by causing visibility reduction, eye and throat irritation, and particularly big damage to agriculture (Middleton et al. 1950; Finlayson-Pitts and Pitts 2000). Although the cause of so-named Los Angeles smog was unexplained at the beginning, Haagen-Smit (1952), and Haagen-Smit, Bradley and Fox (1956) elucidated for the first time that it is ascribed to the toxic substances, including ozone and other strongly oxidizing compounds so-named *photochemical oxidants*, formed by the solar irradiation to the mixtures of nitrogen oxides and non-methane hydrocarbons (NMHCs) emitted from automobile exhaust. Such atmospheric photochemical processes were systematized by Leighton (1961), and his book, *Photochemistry of Air Pollution* is now a classic of atmospheric photochemistry. Los Angeles smog was later called *photochemical smog* (e.g. Rogers 1958), and the term, *photochemical air pollution*, is now widely used including more general concept (e.g. Robinson 1972).

In photochemical smog, other than gaseous oxidants, many kinds of organic aerosols together with sulfate and nitrate were found as particulate matter, and these were shown to be SOAs, formed by the photo-irradiation of the mixtures of NO_x and NMHCs such as auto exhaust (Mader et al. 1952; Renzetti and Doyle 1959). Incidentally, the NMHC was used as a general term for the collectives of short-lived hydrocarbons, excluding methane, which has a longer atmospheric lifetime of nearly 10 years and does not contribute to urban photochemical air pollution directly. Recently, instead of NMHC, the term nonmethane volatile organic compounds (NMVOC) has been more widely used to include oxygen-containing organic compounds other than hydrocarbons. From the early days of the study on the formation mechanism of photochemical smog, there was interest in what kinds of hydrocarbons generate SOAs more effectively. Studies of Haagen-Smit (1952) showed that cyclic hydrocarbons with double bonds such as cyclohexene, indene, and cyclopentadiene easily form low-volatile oxygenated compounds with higher yields of aerosols, since the multiple functional groups are introduced by the ring-opening reactions. This was later confirmed by O'Brien, Holmes, and Bokian (1975), and they reported α -pinene having a cyclic double bond and dialkenes such as isoprene form aerosols with the higher yields, and monocyclic aromatic hydrocarbons such as xylenes also form aerosols with the lower yields.

In the organic aerosols, particulate alkanes, alkenes, alkylbenzenes, naphthalene, etc. were detected as primary organic aerosols (POAs) released directly from emission sources, and pinonic acid, adipic acid, phenols, alkyl nitrates, etc. as SOA formed in the atmosphere. Primary aerosols are not correlated with ozone, but the secondary aerosols have a high correlation with ozone and have peak concentration in early afternoon. Such clear distinction between the POA and SOA was made in the middle of 1970s (Appel, Colodny, and Wesolowski 1976; Cronn et al. 1977). These early studies

showed that alcohols, carboxylic acids, and carbonyl compounds are included in the aerosols by use of infrared absorption spectroscopy and mass spectrometry (Cukor et al. 1972; Ciaccio et al. 1974; Cronn et al. 1977). Also, it was revealed that atmospheric carbonaceous aerosol sampled in California consisted of elemental carbon (EC) and organic carbon (OC) (Appel, Colodny and Wesolowski 1976).

1.3.2 Blue Haze

It is experienced for a long time that the atmosphere in the boundary layer over forests is covered by blue haze after the air is cleaned by rain in summer. In 1950s, a botanist, Went (1960a) addressed the interest to this phenomenon from the viewpoint of atmospheric aerosols. In the “Blue Mountains” near Sydney, Australia, and “Blue Ridges” near the Smoky Mountains in Tennessee, United States, such blue haze are often visible (Ferman, Wolff, and Kelly 1981). Went (1960a) mentioned that a similar phenomenon in Tuscany, Italy, was described in a note by Leonardo da Vinci a long time ago in sixteenth century. This kind of haze is not natural dust or mist, nor the effects of biomass burning or air pollution, and can be seen in dry air irrelevant to water vapor. From these considerations Went (1960a) concluded that blue haze is due to the Tyndall effect described by Tyndall (1869) in the middle of nineteenth century, a phenomenon that a light path can be seen bright from an oblique due to scatter of light by fine particles in air. He proposed that when the fine particles with a diameter of the order of $0.1 \mu\text{m}$ exist in the atmosphere, blue light in the solar light is effectively scattered and blue haze is visible.

As for the possibility of formation of such sub-micron fine particles over the clean forests, Went (1960b) suggested that the photochemical smog reaction of hydrocarbons such as terpenes and isoprene emitted by plants. However, atmospheric concentrations of such biogenic hydrocarbons were not known in early 1960s. They were measured for the first time by Rasmussen and Went (1965), and total concentration of isoprene and terpenes up to ~ 10 ppbv was reported in several forest highlands in United States.

Later measurement of chemical composition of field aerosols at Great Smoky Mountains showed that the main component of fine particles is sulfate and the fraction of organic compounds are relatively small (Stevens et al. 1980; Ferman, Wolff, and Kelly 1981) in those days. Seasonality of the aerosol components were shown that concentrations of sulfate and organic aerosols are high in summer and that of nitrate is high in winter, reflecting the photochemical formation of SO_4^{2-} and OA, and temperature-dependent gas-solid equilibrium of NH_4NO_3 (Day, Malm, and Kreidenweis 1997). Meanwhile, although the measurement of chemical analysis of aerosols in Blue Mountains in Australia is scarce, solvent extracted organics contains n-alkane, n-alkanoic acid, and n-alcohol, which compose lipids contained in plant wax (Simoneit et al. 1991).

It is interesting to note that the prototype of SOA formation from biogenic and anthropogenic hydrocarbons was thus shown in early studies more than 50 years ago. The research on SOA has been developed extensively after the year of 2000 being related to the interests in the impact on human health of $\text{PM}_{2.5}$ and in the climate impact of aerosols.

References

- Akimoto, H. (2016). *Atmospheric Reaction Chemistry*. Tokyo: Springer Japan.
- Appel, B.R., Colodny, P., and Wesolowski, J.L. (1976). Analysis of carbonaceous materials in southern California atmospheric aerosols. *Environ. Sci. Technol.* 10: 359–363.
- Appel, B.R., Kothny, E.L., Hoffer, E.M. et al. (1978). Sulfate and nitrate data from the California aerosol characterization experiment (AICHEX). *Environ. Sci. Technol.* 12: 418–425.
- Backstrom, H. (1934). The chain reaction mechanism in the autoxidation of sodium sulfite solution. *Z. Physik. Chem.* B25: 99–121.
- Barrett, E. and Brodin, G. (1955). The acidity of Scandinavian precipitation. *Tellus* 2: 251–257.
- Barrie, L.A., Yi, Y., Leaitch, W.R. et al. (2001). A comparison of large-scale atmospheric sulphate aerosol models (COSAM): overview and highlights. *Tellus* 53B: 615–645.
- Bassett, M. and Seinfeld, J.H. (1983). Atmospheric equilibrium model of sulfate and nitrate aerosols. *Atmos. Environ.* 17: 2237–2252.
- Brimblecombe, P. and Spedding, D.J. (1974). The catalytic oxidation of micromolar aqueous sulphur dioxide. I. Oxidation in dilute solutions containing iron (III). *Atmos. Environ.* 8: 937–945.
- Burkholder, J.B., Sander, S.P., Abatt, J.P.D. et al. (2015). *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation Number 18*, JPL Publication 15-10. Pasadena, California: NASA and the Jet Propulsion Laboratory, California Institute of Technology, <http://jpldataeval.jpl.nasa.gov>.
- Ciaccio, L.L., Rubino, R.L., and Flores, J. (1974). Composition of organic constituents in breathable airborne particulate matter near a highway. *Environ. Sci. Technol.* 8: 935–942.
- Cogbill, C.V. (1976). The history and character of acid precipitation in eastern North America. *Water Air Soil Pollut.* 6: 407–413.
- Cowling, E.B. (1982). Acid precipitation in historical perspective. *Environ. Sci. Technol.* 16: 110A–123A.
- Cox, R.A. and Penkett, S.A. (1972). Aerosol formation from sulphur dioxide in the presence of ozone and olefinic hydrocarbons. *J. Chem. Soc., Faraday Trans.* 68: 1735–1753.
- Cronn, D.R., Charlson, R.J., Knights, R.L. et al. (1977). A survey of the molecular nature of primary and secondary components of particles in urban air by high-resolution mass spectrometry. *Atmos. Environ.* 11: 929–937.
- Cukor, P., Ciaccio, L.L., Lanning, E.W., and Rubino, R.L. (1972). Some chemical and physical characteristics of organic fractions in airborne particulate matter. *Environ. Sci. Technol.* 6: 633–637.
- Davis, D.D., Ravishankara, A.R., and Fischer, S. (1979). SO₂ oxidation via the hydroxyl radical: atmospheric fate of HSO_x radicals. *Geophys. Res. Lett.* 6: 113–116.
- Day, D.E., Malm, W.C., and Kreidenweis, S.M. (1997). Seasonal variations in aerosol composition and acidity at Shenandoah and Great Smoky Mountains National Parks. *J. Air Waste Manage. Assoc.* 47: 411–418.

- De Haan, D.O. and Finlayson-Pitts, B.J. (1997). Knudsen cell studies of the reaction of gaseous nitric acid with synthetic sea salt at 298 K. *J. Phys. Chem. A* 101: 9993–9999.
- Deguillaume, L., Leriche, M., and Desboe, K. (2005). Transition metals in atmospheric liquid phases: sources, reactivity, and sensitive parameters. *Chem. Rev.* 105: 3388–3431.
- Eggleton, A.E.J. and Cox, R.A. (1978). Homogeneous oxidation of sulphur compounds in the atmosphere. *Atmos. Environ.* 12: 227–230.
- Emanuelsson, A., Eriksson, E., and Egnér, H. (1954). Composition of atmospheric precipitation in Sweden. *Tellus* 6: 261–267.
- Erickson, R.E., M, L., Yates, R.L.C., and McEwen, D. (1977). The reaction of sulfur dioxide with ozone in water and its possible atmospheric significance. *Atmos. Environ.* 11: 813–817.
- Eriksson, E. (1952a). Composition of atmospheric precipitation I. nitrogen compounds. *Tellus* 4: 215–232.
- Eriksson, E. (1952b). Composition of atmospheric precipitation II. Sulfur, chloride, iodine compounds. *Tellus* 4: 280–303.
- Ferman, M.A., Wolff, G.T., and Kelly, N.A. (1981). The nature and sources of haze in the Shenandoah Valley/Blue Ridge Mountains area. *J. Air Pollut. Control Assoc.* 31: 1074–1082.
- Finlayson-Pitts, B.J. and Pitts, J.N. Jr. (2000). *Chemistry of the Upper and Lower Atmosphere*. Academic Press.
- Fudakowski, H. (1873). Zur Lehre von dem Activwerden des Sauerstoffs bei langsamen Oxydationen. *Ber. Dtsch. Chem. Ges.* 6: 106–109. (in German) <http://doi.org/10.1002/cber.18730060141>.
- Fuller, E.C. and Crist, R.H. (1941). The rate of oxidation of sulfite ions by oxygen. *J. Am. Chem. Soc.* 63: 1644–1650.
- Gorham, E. and Gordon, A.G. (1960). The influence of smelter fumes upon the chemical composition of lake waters near Sudbury, Ontario, and upon the surrounding vegetation. *Can. J. Bot.* 38: 477–487.
- Haagen-Smit, A.J. (1952). Chemistry and physiology of Los Angeles smog. *Ind. Eng. Chem.* 44: 1342–1346.
- Haagen-Smit, A.J., Bradley, C.E., and Fox, M.M. (1956). Ozone formation in photochemical oxidation of organic substances. *Ind. Eng. Chem.* 45: 2086–2089.
- Harris, E., Sinha, B., van Pinxteren, D. et al. (2013). Enhanced role of transition metal ion catalysis during in-cloud oxidation of SO₂. *Science* 340: 727–730.
- Harrison, R.M. and Pio, C.A. (1983). An investigation of the atmospheric HNO₃-NH₃-NH₄NO₃ equilibrium relationship in a cool, humid climate. *Tellus* 35B: 155–159.
- Hegg, D.A. and Hobbs, P.V. (1978). Oxidation of sulfur dioxide in aqueous systems with particular reference to the atmosphere. *Atmos. Environ.* 12: 241–253.
- Heikes, B. and Thompson, A.M. (1983). Effects of heterogeneous processes on NO₃, HONO, and HNO₃ chemistry in the troposphere. *J. Geophys. Res.* 88: 10883–10895. (Correction, *J. Geophys. Res.*, 89, 11829, 1984.)
- Herrmann, H., Schaefer, T., Tilgner, A. et al. (2015). Tropospheric aqueous-phase chemistry: kinetics, mechanisms, and its coupling to a changing gas phase. *Chem. Rev.* 115: 4259–4334.
- Hildemann, L.M., Russell, A.G., and Cass, G.R. (1984). Ammonia and nitric acid concentrations in equilibrium with atmospheric aerosols. *Atmos. Environ.* 18: 1737–1750.

- Hoffmann, M.R. and Edwards, J.O. (1975). Kinetics of the oxidation of sulfite by hydrogen peroxide in acidic solution. *J. Phys. Chem.* 79: 2096–2098.
- Junge, C.E. (1954). The chemical composition of atmospheric aerosols, I: measurements at Round Hill field station, June–July 1953. *J. Meteorol.* 11: 323–333.
- Junge, C.E. (1963). *Air Chemistry and Radioactivity*. New York/London: Academic Press.
- Junge, C.E. and Ryan, T.G. (1958). Study of the SO₂ oxidation in solution and its role in atmospheric chemistry. *Q. J. R. Meteorolog. Soc.* 84: 46–55.
- Junge, C.E. and Werby, R.T. (1958). The concentration of chloride, sodium, potassium, calcium, and sulfate in rain water over the United States. *J. Meteorol.* 15: 417–425.
- Kotin, P., Falk, H.I., Mader, P., and Thomas, M. (1954). Aromatic hydrocarbons. I. Presence in the Los Angeles atmosphere and the carcinogenicity of atmospheric extracts. *Arch. Ind. Hyg. Occup. Med.* 9: 153–163.
- Lampadius, W.A. (1837). Fortgesetzte Beiträge zur Kenntnis verschiedener Wasser. *J. Prakt. Chem.* 10: 78–88. (in German).
- Larson, T.V., Horike, N.H., and Harrison, H. (1978). Oxidation of sulfur dioxide by oxygen and ozone in aqueous solution: a kinetic study with significance to atmospheric rate processes. *Atmos. Environ.* 12: 1597–1611.
- Lee, R.E. and Patterson, R.K. (1969). Size determination of atmospheric phosphate, nitrate, chloride, and ammonium particulate in several urban areas. *Atmos. Environ.* 3: 249–255.
- Leighton, P.A. (1961). *Photochemistry of Air Pollution*. Academic Press.
- Liebig, J.V. (1835). Über einige Stickstoffverbindungen. – Poggendorfs. *Ann. Phys. Chem.* 34: 570–613. (in German).
- Likens, G.E. and Bormann, F.H. (1974). Acid rain: a serious regional environmental problem. *Science* 184: 1176–1179.
- Ludwig, F.L. and Robinson, E. (1965). Size distribution of sulfur-containing compounds in urban aerosols. *J. Colloid Sci.* 20: 571–584.
- Lundgren, D.A. (1970). Atmospheric aerosol composition and concentration as a function of particle size and of time. *J. Air Pollut. Control Assoc.* 20: 603–608.
- Mader, P.M. (1958). Kinetics of the hydrogen peroxide-sulfite reaction in alkaline solution. *J. Am. Chem. Soc.* 80: 2634–2639.
- Mader, P.P., MacPhee, R.D., Lofberg, R.T., and Larson, G.P. (1952). Composition of organic portion of atmospheric aerosols in the Los Angeles area. *Ind. Eng. Chem.* 44: 1352–1355.
- Marggraf, A.S. (1751). Examen chymique de l'eau. *Histoire de l'Académie Royale des Sciences et des Belles-Lettres* 7: 131–157. (in French).
- Middleton, J.T., Kendrick, J.B. Jr., and Schwalm, H.W. (1950). Injury to herbaceous plants by smog or air pollution, U.S.D.A. *Plant Dis. Rep.* 34: 249–252.
- Miller, N.H.J. (1905). The amounts of nitrogen as ammonia and as nitric acid and chlorine in the rainwater collected at Rothamsted. *J. Agric. Sci.* 1: 280–303.
- Möller, D. (2008). On the history of the scientific exploration of fog, dew, rain and other atmospheric water. *Die Erde* 139: 11–44.
- Mozurkewich, M. and Calvert, J.G. (1988). Reaction probability of N₂O₅ on aqueous aerosols. *J. Geophys. Res.* 93: 15889–15896.
- O'Brien, R.J., Holmes, J.R., and Bockian, A.H. (1975). Formation of photochemical aerosol from hydrocarbons: chemical reactivity and products. *Environ. Sci. Technol.* 9: 568–576.
- Odén, S. (1976). The acidity problem – an outline of concepts. *Water Air Soil Pollut.* 6: 137–166.
- Orel, A.E. and Seinfeld, J.H. (1977). Nitrate formation in atmospheric aerosols. *Environ. Sci. Technol.* 11: 1000–1007.

- Penkett, S.A. and Garland, J.A. (1974). Oxidation of sulphur dioxide in artificial fogs by ozone. *Tellus* 26: 284–290.
- Penkett, S.A., Jones, B.M.R., Brice, K.A., and Eggleton, A.E.J. (1979). The importance of atmospheric ozone and hydrogen peroxide in oxidizing sulphur dioxide in cloud and rainwater. *Atmos. Environ.* 13: 123–137.
- Rasmussen, R.A. and Went, F.W. (1965). Volatile organic material of plant origin in the atmosphere. *Proc. Natl. Acad. Sci. U.S.A.* 53: 215–220.
- Reinders, W. and Vles, S.I. (1925). Reaction velocity of oxygen with solutions of some inorganic salts: III. The catalytic oxidation of sulphites. *Recl. Trav. Chim. Pays-Bas* 44: 249–268.
- Renzetti, N.A. and Doyle, G.J. (1959). The chemical nature of the particulate in irradiated automobile exhaust. *J. Air Pollut. Control Assoc.* 8: 293–296.
- Richards, L.W. (1983). Comments on the oxidation of NO₂ to nitrate—day and night. *Atmos. Environ.* 17: 397–402.
- Robbins, R.C., Cadle, R.D., and Eckhardt, D.L. (1959). The conversion of sodium chloride to hydrogen chloride in the atmosphere. *J. Meteorol.* 16: 53–56.
- Robinson, E. (1972). Global aspects of photochemical air pollution. In: *Photochemical Smog and Ozone Reaction*, Advances in Chemistry Series, vol. 113 (ed. R.F. Gould), 1–15. American Chemical Society.
- Rogers, L.H. (1958). Report on photochemical smog. *J. Chem. Educ.* 35: 310.
- Saxena, P., Seigneur, C., and Peterson, T.W. (1983). Modeling of multiphase atmospheric aerosols. *Atmos. Environ.* 17: 1315–1329.
- Seinfeld, J.H. and Pandis, S.N. (2016). *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 3e. Hoboken, NJ: Wiley.
- Simoneit, B.R.T., Crisp, P.T., Mazurek, M.A., and Standley, L.J. (1991). Composition of extractable organic matter of aerosols from the Blue Mountains and southeast coast of Australia. *Environ. Int.* 17: 405–419.
- Smith, R.A. (1852). On the air and rain of Manchester. *Mem. Manchester Lit. Philos. Soc.* 2: 207–217.
- Smith, R.A. (1872). *Air and Rain: The Beginning of a Chemical Climatology*. Green, London: Longmans.
- Stelson, A.B.W., Friedlander, S.K., and Seinfeld, J.H. (1979). A note on the equilibrium relationship between ammonia and nitric acid and particulate ammonium nitrate. *Atmos. Environ.* 13: 369–371.
- Stevens, R.K., Dzubay, T.G., Shaw, R.W. Jr. et al. (1980). Characterization of the aerosol in the Great Smoky Mountains. *Environ. Sci. Technol.* 14: 1491–1498.
- Stocks, P. and Campbell, M.J. (1955). Lung cancer death rates among non-smokers and pipe and cigarette smokers: evaluation in relation to air pollution by benzo[a]pyrene and other substances. *BMJ* 2: 923–929.
- Stockwell, W.R. and Calvert, J.G. (1983). The mechanism of the HO-SO₂ reaction. *Atmos. Environ.* 17: 2231–2235.
- Tyndall, J. (1869). On a new series of chemical reactions produced by light. *Proc. R. Soc. London* 17: 92–102.
- Van den Heuvel, A.P. and Mason, B.J. (1963). The formation of ammonium sulfate in water droplets exposed to gaseous sulfur dioxide and ammonia. *Q. J. R. Meteorolog. Soc.* 89: 271–275.

- Wagman, J., Lee, R.E., and Axt, C.J. (1967). Influence of some atmospheric variables on the concentration and particle size distribution of sulfate in urban air. *Atmos. Environ.* 1: 479–489.
- Wahner, A., Mentel, T.F., Sohn, M., and Stier, J. (1998). Heterogeneous reaction of N_2O_5 on sodium nitrate aerosol. *J. Geophys. Res.* 103: 31103–31112.
- Waller, R.E. (1952). The benzpyrene content of town air. *Br. J. Cancer* 6: 8–21.
- Went, F.W. (1960a). Blue hazes in the atmosphere. *Nature* 187: 641–643.
- Went, F.W. (1960b). Organic matter in the atmosphere, and its possible relation to petroleum formation. *Proc. Natl. Acad. Sci. U.S.A.* 46: 212–221.
- Wilson, B.D. (1926). Nitrogen and sulfur in rainwater in New York. *J. Am. Soc. Agric.* 18: 1108–1112.
- Wynder, E.L. and Hoffmann, D. (1965). Some laboratory and epidemiological aspects of air pollution carcinogenesis. *J. Air Pollut. Control Assoc.* 15: 155–159.

