


# I: CELLULAR PHYSIOLOGY

- Transmembrane solute transport, 3
  - Ion channels, 4
  - Ion ATPases, 6
- Resting membrane potential ( $E_m$ ), 8
  - Action potential, 10
  - Second messenger pathways, 12

COPYRIGHTED MATERIAL



Transmembrane solute transport



**Fig. 1**  
A, B and C are different molecules.

## Ion channels

Ion channels are protein tunnels spanning the cell membrane. Channel opening results in a current of the order of a few picoamps generated by the flow of highly specific ions.

### Potassium channels

#### (a) Outward or delayed rectifier K<sup>+</sup> channel (K<sub>v</sub>)

Activated by membrane depolarization  
Produces an outward K<sup>+</sup> current  
Responsible for the repolarization of the cardiac action potential

#### (b) ATP-sensitive K<sup>+</sup> channel (K<sup>+</sup>-ATP)

Accelerates repolarization  
Shortens the cardiac action potential

*Prostacyclin, vasoactive intestinal peptide (VIP), nitric oxide (NO) and adenosine act in part via K<sup>+</sup>-ATP opening*

*K<sup>+</sup>-ATP channels open during ischaemia in response to a fall in intracellular ATP, acidosis, a rise in ADP and GDP, and the accumulation of extracellular adenosine*

*Antianginal (nicorandil) and vasodilator agents (diazoxide and minoxidil) act via myocyte K<sup>+</sup>-ATP opening. Sulphonylureas such as glibenclamide are selective K<sup>+</sup>-ATP blockers*

#### (c) G-protein-activated K<sup>+</sup> channel (K-ACh)

Opened by vagally secreted acetylcholine (ACh)  
Decreases spontaneous depolarization in the sinus node  
Slows atrioventricular (AV) node conduction, underlying the vagal slowing of heart rate

#### (d) Inwardly rectifying K<sup>+</sup> channel

Opens at very negative potentials (less than -40 mV)  
Shows a reduced K<sup>+</sup> conductance at positive membrane potentials (opposite to normal outward rectification seen in delayed rectifier channels)  
K<sup>+</sup>-ATP and K<sup>+</sup>-ACh display some inward rectification

### Calcium channels

#### (a) L-type Ca<sup>2+</sup> channel (long lasting)

High voltage activated  
Expressed in vascular and cardiac tissue  
Generates a slow inward current  
*Blocked by dihydropyridines (nifedipine, amlodipine)*

#### (b) T-type Ca<sup>2+</sup> channel (transient)

Low voltage activated  
Rapidly inactivated  
High expression in the sinus node—possible role in pacemaking  
*Blocked by verapamil, diltiazem*

#### (c) N-, P-, Q- and R-type Ca<sup>2+</sup> channels

Found in neuronal cells

## Ion channel disorders

Disorder	Channel	Clinical notes
Bartter's syndrome	AR Bumetanide-sensitive Na <sup>+</sup> K <sup>+</sup> Cl <sup>-</sup> cotransporter (NKCC2)	<i>Hypokalaemia, alkalosis, renal salt wasting, hypotension, hyperreninaemia, hyperaldosteronism</i>
Liddle's syndrome (hereditary hypertension)	AR ENaC (epithelial Na channel)	
Hyperkalaemic periodic paralysis	AD Skeletal muscle Na channel	
Hypokalaemic periodic paralysis	AD L-type Ca <sup>2+</sup> channel	
Becker's generalized myotonia	AR Skeletal muscle Cl channel	
Long QT syndrome	AD Type 1, KVLQT1 (cardiac K <sup>+</sup> channel) Type 2, HERG (cardiac K <sup>+</sup> channel) Type 3, SCN5A (cardiac Na <sup>+</sup> channel)	<i>Characterized by prolonged and abnormal ventricular repolarization and risk of life-threatening arrhythmias (particularly torsades de pointes)</i>

AD, autosomal dominant; AR, autosomal recessive.


## Cystic fibrosis (CF)

- The CF transmembrane conductance regulator (*CFTR*) gene is defective in CF.
- CFTR is a cAMP-regulated Cl channel found in the apical membrane of epithelial cells.
- CFTR also downregulates Na absorption via the amiloride-sensitive ENaC.
- Reduced Cl transport is thought to reduce Cl and water secretion into the airway lumen.

## Ion ATPases

### Na<sup>+</sup>/K<sup>+</sup>-ATPase

The chemical energy of ATP hydrolysis is used to extrude three Na<sup>+</sup> ions for every two K<sup>+</sup> ions entering the cell and every ATP molecule hydrolysed.



**Fig. 2 Na<sup>+</sup>/K<sup>+</sup>-ATPase function**

d, digoxin-binding site; ECF, extracellular fluid; ICF, intracellular fluid.

There is a net export of one third of a positive charge per Na<sup>+</sup> ion transported. Intracellular Na<sup>+</sup> is the substrate of the pump and a rise in intracellular Na<sup>+</sup> concentration favours Na<sup>+</sup>/K<sup>+</sup> exchange.

Na<sup>+</sup>/K<sup>+</sup>-ATPase maintains intracellular and extracellular Na<sup>+</sup> and K<sup>+</sup> concentrations and is thus responsible for maintaining the resting mem-

brane potential. The active transport of  $\text{Na}^+$  is also coupled to the transport of other substances (secondary active transport, counter transport and cotransport).

Magnesium is a cofactor of  $\text{Na}^+/\text{K}^+$ -ATPase and thus helps to maintain intracellular  $\text{K}^+$ .

*Digoxin is an  $\text{Na}^+/\text{K}^+$ -ATPase inhibitor and thus produces a rise in intracellular  $\text{Na}^+$  as well as a fall in intracellular  $\text{K}^+$ .*

### Other ATPases

#### *Gastric $\text{H}^+/\text{K}^+$ -ATPase*

- Responsible for hydrogen ion secretion.
- *Antigen recognized by parietal cell autoantibodies in pernicious anaemia.*

#### *$\text{Ca}^{2+}/\text{Mg}^{2+}$ -ATPase*

- Actively pumps  $\text{Ca}^{2+}$  into the sarcoplasmic reticulum during muscular relaxation (see 'Excitation–contraction coupling', p. 176).

#### *$\text{H}^+$ -ATPase*

- Responsible for acid secretion in the distal convoluted tubule and collecting duct of the kidney.
- *A deficiency of this active proton pump (as in Sjögren's syndrome) results in distal (type 1) renal tubular acidosis (see 'Renal', p. 97).*

## Resting membrane potential ( $E_m$ )

The Nernst potential ( $E_{rev}$ ) for an ion is the point at which chemical and electrical driving forces across the cell membrane (occurring in opposite directions) are in equilibrium. At this potential, there is no net flow of that specific ion.

Ion	Extracellular concentration (mmol L <sup>-1</sup> )	Intracellular concentration (mmol L <sup>-1</sup> )	Nernst potential (mV)
Na <sup>+</sup>	142	10	+70 ( $E_{Na}$ )
K <sup>+</sup>	4	155	-98 ( $E_K$ )
Ca <sup>2+</sup>	2.5	0.0001	+150 ( $E_{Ca}$ )
Cl <sup>-</sup>	101	5–30	+30 to -65 ( $E_{Cl}$ )

Under physiological conditions, Na<sup>+</sup>, Ca<sup>2+</sup> and Cl<sup>-</sup> flow into cells to depolarize the cell towards  $E_{Na}$ ,  $E_{Ca}$  and  $E_{Cl}$  respectively. Similarly, K<sup>+</sup> flows out of the cell to repolarize the cell towards  $E_K$ .  $E_m$  depends on the distribution of Na<sup>+</sup>, Ca<sup>2+</sup>, Cl<sup>-</sup> and K<sup>+</sup> as well as membrane permeability to these ions.





Fig. 3

## Action potential




**Fig. 4 Axonal action potential**

(From Schmidt, R.F. & Thews, G. (eds) (1983) *Human Physiology*. Springer-Verlag, Berlin.)

The action potential is an all or nothing event triggered by the arrival of a depolarizing stimulus when  $\text{Na}^+$  influx ( $g_{\text{Na}}$ ) exceeds  $\text{K}^+$  efflux ( $g_{\text{K}}$ ).

Depolarization	When a critical threshold ( $-55 \text{ mV}$ ) is reached, all voltage-gated $\text{Na}^+$ channels open, causing $E_m$ to approach $E_{\text{Na}}$ ( $+55 \text{ mV}$ ) rapidly
Repolarization	A delayed voltage-dependent $\text{Na}^+$ channel inactivation and $\text{K}^+$ channel activation causes $E_m$ to fall, exceeding the resting potential briefly (hyperpolarization) before returning to the starting point


The action potential is followed by an absolute and then relative refractory period.



**Fig. 5 Cardiac action potential**

SA, sinoatrial; AV, atrioventricular.

## Second messenger pathways



**Fig. 6**

AC, adenyl cyclase;  $Ca^{2+}$ ; CaMK, calmodulin-dependent kinase; DAG, 1,2-diaclyglycerol; Gs, Gi, Gq, G proteins;  $IP_3$ , inositol 1,4,5-triphosphate; PC, phospholipase C;  $PIP_2$ , phosphatidylinositol 4,5-biphosphate; PKA, protein kinase A; PKC, protein kinase C. Receptors: DA, dopamine; M, muscarinic.

### cAMP pathway

Activated  $\beta_1$  and  $\alpha_2$  adrenergic receptors, for example, act via Gs or Gi proteins to stimulate or inhibit AC respectively

AC induces cAMP synthesis

cAMP stimulates target gene expression (tyrosine hydroxylase, somatostatin) via:

1 PKA induction

2 phosphorylation of transcription factors (cAMP-responsive element (CRE)-binding protein, CREB)

### $IP_3$ pathway

Activated  $\alpha_1$  adrenergic receptors, for example, act via G proteins to stimulate PC

PC cleaves phosphoinositide to give  $IP_3$  and DAG

$IP_3$  mobilizes  $Ca^{2+}$  from intracellular stores

$Ca^{2+}$  and DAG activate calmodulin kinases and PKC

These in turn phosphorylate a number of important proteins (epidermal growth factor receptor (EGFR), glycogen synthase)

### Notes:

Ca may modulate CREB activity via calmodulin kinases but also induces target gene expression via the cAMP pathway.

Other second messengers include cGMP (atrial natriuretic peptide (ANP), NO, phototransduction).

Use of second messenger pathways by various agonists

Agonist	cAMP raised	cAMP reduced	IP <sub>3</sub> /DAG
ACh		M <sub>2</sub>	M <sub>1</sub>
Epinephrine	β <sub>1</sub>	α <sub>2</sub>	α <sub>1</sub>
Dopamine	DA <sub>1</sub>	DA <sub>2</sub>	
ADH	VP <sub>2</sub>		VP <sub>1</sub>
Histamine	H <sub>2</sub>		H <sub>1</sub>
Adenosine	A <sub>2</sub>	A <sub>1</sub>	
Other	TSH	Somatostatin	Gastrin
	LH	All	CCK
	FSH	5HT	GABA

ADH, antidiuretic hormone; CCK, cholecystokinin; FSH, follicle-stimulating hormone; GABA, γ-aminobutyric acid; 5HT, 5-hydroxytryptamine; LH, luteinizing hormone; TSH, thyroid-stimulating hormone; VP, vasopressin.

G proteins




Fig. 7 G-protein function

E, effector molecule; F, first messenger; R, receptor. G protein: A, α subunit; B, β and γ subunits.

G proteins consist of three subunits (A, B, C).

- 1 In the resting state, GDP is bound to the A subunit which is a GTPase.
- 2 On hormone binding, GDP is displaced by GTP which activates the G protein.
- 3 The A subunit and BC complex dissociate to interact with effectors.
- 4 GTP is then rapidly hydrolysed to GDP.

G-protein abnormalities are implicated in human disease:

- 1 continued Gs activation is a pathophysiological mechanism in acromegaly, McCune–Albright syndrome and *Vibrio cholerae* infection;
- 2 the oncogene *ras* encodes p21 which is a G protein;
- 3 Gs activity is reduced by 50% in pseudohypoparathyroidism.

