CHAPTER 1 Introduction

1.1 A historical background to metal aquatic chemistry

In terrestrial waters, much of the initial work by environmental engineers and scientists was aimed at understanding the processes of waste water treatment, and of the reactions and impact of human-released chemicals on the environment, and on the transport and fate of radioactive chemicals. This initial interest was driven by the need to understand industrial processes and the consequences of these activities, and the recognition of the potential toxicity and environmental impact of trace metals released during extraction. from the Earth as well as during refining and use. Initial interest in the chemistry of the ocean was driven by the key question of why the sea is salty. Of course, it is now known that freshwaters, and even rainwater, contain a small amount of dissolved salts; and that the high salt content of the ocean, and some terrestrial lakes, is due to the buildup of salts as a result of continued input of material to a relatively enclosed system where the major loss of water is due to evaporation rather than outflow. This is succinctly stated by Broecker and Peng [1]: "The composition of sea salt reflects not only the relative abundance of the dissolved substances in river water but also the ease with which a given substance is entrapped in the sediments".

The constituents in rivers are derived from the dissolution of rocks and other terrestrial material by precipitation and more recently, through the addition of chemicals from anthropogenic activities, and the enhanced release of particulate through human activity. The dissolution of carbon dioxide (CO_2) in rainwater results in an acidic solution and this solution subsequently dissolves the mostly basic constituents that form the Earth's crust [2]. The natural acidity, and the recently enhanced acidity of precipitation due to human-derived atmospheric inputs, results in the dissolution (weathering) of the terrestrial crust and the flow of freshwater to the ocean transports these dissolved salts, as well as suspended particulate matter, entrained and resuspended during transport. However, the ratio of river to ocean concentration is not fixed for all dissolved ions [3, 4] as the concentration in the ocean is determined primarily by the ratio of the rate of input compared to the rate of removal, which will be equivalent at steady state. Thus, the ocean concentration of a constituent is related to its reactivity, solubility or other property that may control its rates of input or removal [1, 5].

Many of the early investigations looked at the aquatic ecosystem as an inorganic entity or reactor, primarily based on the assumption that aquatic chemistry was driven by abiotic chemical processes and reactions, and that the impact of organisms on their environment was relatively minor. In 1967, Sillen [6], a physical chemist, published a paper *The Ocean as a Chemical System*, in which he aimed at explaining the composition of the ocean in terms of various equilibrium processes, being an equilibrium mixture of so-called "volatile components" (e.g., H₂O and HCl) and "igneous rock" (primarily KOH, Al(OH)₃ and SiO₂), and other components, such as CO₂, NaOH, CaO and MgO. His analysis was a follow-up of the initial proposed weathering reaction of Goldschmidt in 1933:

0.6 kg igneous rock + \sim 1 kg volatiles = 1 L sea water + \sim 0.6 kg sediments + 3 L air (1.1)

Sillen stated that "the composition may . . . be given by well-defined equilibria, and that deviations from equilibrium may be explainable by well-defined processes", and proposed the box model shown in Fig. 1.1 [6]. However, he also stated that this did not "mean that I suggest that there would be true equilibrium in the real system". In reality, the ocean composition is a steady state system where the

Trace Metals in Aquatic Systems, First Edition. Robert P. Mason.

^{© 2013} Robert P. Mason. Published 2013 by Blackwell Publishing Ltd.



Fig. 1.1 A representation of the global material cycle. The reservoirs are indicated as symbols C = continental rock; B: basaltic rock; O: ocean; S_c: continental sediments; S_B: basaltic sediments. The fluxes into (+) and out of (–) each reservoir are indicated by appropriate symbols that represent the flow paths. Redrawn from Sillen (1967) *Science* **156**: 1189–97. Reprinted with permission of AAAS [6].

concentrations are determined primarily by the relative rates of addition and removal, and that reversible equilibrium reactions are not the primary control over ocean composition [1]; something Sillen [6] alluded to. The major difference between these two conceptualizations is that the equilibrium situation would lead to a constancy of composition over geological time while the steady state model accommodates variation due to changes in the rate of input of chemicals to the ocean. The Sillen paper established the idea that the ocean was a system that could be described as being at geological equilibrium in terms of the major reactions of the primary chemicals at the Earth's surface, and that the average composition of the overall system could be described. The composition of seawater, for example, was due to a series of reversible equilibrium reactions between the ocean waters, sediments and the atmosphere [6]. Similarly, it was proposed that the composition of the biosphere was determined by a complex series of equilibrium reactions - acid-base and oxidation-reduction reactions - by which the reduced volatile acids released from the depths of the Earth by volcanoes and other sources, reacted with the basic rocks of the Earth and with the oxygen in the atmosphere.

A follow-up paper in 1980 by McDuff and Morel [7] entitled *The Geochemical Control of Seawater (Sillen Revisited)*, discussed in detail these approaches and contrasted them in terms of explaining the composition of the major ions in seawater. This paper focused on the controls over alkalinity in the ocean and the recycled source of carbon that is required to balance the removal of CaCO₃ via precipitation in the ocean, and its burial in sediments. These authors concluded that "while seawater alkalinity is directly controlled by the formation of calcium carbonate as its major sedimentary sink, it is also controlled indirectly by carbonate metamorphism which buffers the CO₂ content of the atmosphere" [7]. They concluded that the "ocean composition [is] dominated by geophysical rather than geochemical processes. The acid-base chemistry reflects, however, a fundamental control by heterogeneous chemical processes."

McDuff and Morel [7] also focused attention on the importance of biological processes (photosynthesis and respiration) on the carbon balance, something that the earlier chemists did not consider [6]. These biological processes result in large fluxes via carbon fixation in the surface ocean and through organic matter degradation at depth, but overall most of this material is recycled within the ocean so that little organic carbon is removed from the system through sediment burial. Thus, the primary removal process for carbon is through burial of inorganic material, primarily as Ca and Mg carbonates. Recently, the short-term impact of increasing atmospheric CO₂ on ocean chemistry has been vigorously debated, and is a topic of recent research focus [8, 9] because of the resultant impact of pH change due to higher dissolved CO₂ in ocean waters on the formation of insoluble carbonate materials. Carbonate formation is either biotic (shell formation by phytoplankton and other organisms) or abiotic. Thus, there has been a transition from an initial conceptualization of the ocean and the biosphere in general, as a physical chemical system to one where the biogeochemical processes and cycles are all seen to be important in determining the overall composition. This is true for saline waters as well as large freshwaters, such as the Great Lakes of North America, and small freshwater ecosystems, and even more so for dynamic systems, such as rivers and the coastal zone [2, 4, 10].

It has also become apparent that most chemical (mostly redox) reactions in the environment that are a source of energy are used by microorganisms for their biochemical survival and that, for example, much of the environmental oxidation and reduction reactions of Fe and Mn are mediated by microbes, even in environments that were previously deemed unsuitable for life, because of high temperature and/or acidity [2, 10]. Overall, microorganisms specifically, and biology in general, impact aquatic trace metal(loid) concentrations and fate and their presence and reactivity play multiple roles in the biogeochemistry of aquatic systems. Understanding their environmental concentration, reactivity, bioavailability and mobility are therefore of high importance to environmental scientists and managers. Trace metals are essential to life as they form the basis of many important biochemicals, such as enzymes [10, 11], but they are also toxic and can cause both human and environmental damage. As an example, it is known that mercury (Hg) is an element that is toxic to organisms and bioaccumulates into aquatic food chains, while the other elements in Group 12 of the periodic table - zinc (Zn) and cadmium (Cd) - can play a biochemical role [11]. Until recently, Cd was thought to be a toxic metal only but the

demonstration of its substitution for Zn in the important enzyme, carbonic anhydrase, in marine phytoplankton has demonstrated one important tenet of the trace metal(loid)s – that while they may be essential elements for organisms, at high concentrations they can also be toxic [11, 12]. Copper (Cu) is another element that is both required for some enzymes but can also be toxic to aquatic organisms, especially cyanobacteria, at higher concentrations. This dichotomy is illustrated in Fig. 1.2, and is true not only for metals but for non-metals and organic compounds, many of which also show a requirement at low concentration and a toxicity at high exposure.

This dual role is also apparent for the group of elements, termed the metalloids, which occupy the bottom right region



Fig. 1.2 Concentration-response curve for a trace element that is both required by organisms for survival but that can be toxic to the same organism at high concentration.

of the periodic table (Groups 13-17, from gallium (Ga) to astatine (As); see Fig. 1.3) [12]. They do not fit neatly into the definition of "trace metals," as these elements are transitional between metals and non-metals in chemical character, but their behavior and importance to aquatic biogeochemistry argue for their inclusion in this book. Indeed, selenium (Se) is another example of an element that is essential to living organisms, being found in selenoproteins which have a vital biochemical role, but it can also be toxic if present at high concentrations [13]. As noted previously, the chemistry of both metals and microorganisms are strongly linked and many microbes can transform metals from a benign to a more toxic form, or vice versa; for example, bacteria convert inorganic Hg into a much more toxic and bioaccumulative methylmercury, while methylation of arsenic (As) leads to a less toxic product [14].

Therefore, in this book the term "trace metal(loid)" is used in a relatively expansive manner to encompass: (1) the transition metals, such as Fe and Mn, that are relatively abundant in the Earth's crust but are found in solution at relatively low concentrations due to their insolubility, and because they play an important role in biogeochemical cycles [2, 10]; (2) the so-called "heavy metals," such as Hg, Cd, lead (Pb) and Zn that are often the source of environmental concern, although some can have a biochemical role [4, 10]; (3) the metalloids, such as As and Se, which can be toxic and/or required elements in organisms; and (4) includes the lanthanides and actinides, which exist at low concentration and/or are radioactive [15]. Many of the lanthanides (so-called "rare earth elements") are now being

hydrogen											e gases							
H	1 □ alkali earth metals □ nonmeta										als			2 He				
Li 3	Be ⁴	,									Б Б	C	7 N	08	F ⁹	¹⁰ Ne		
11 Na	12 Mg	🔳 tra	transition metals										18 Ar					
19 K	20 Ca	21 Sc	22 Ti	23 V	24 Cr	²⁵ Mn	26 Fe	27 Co	28 Ni	29 Cu	³⁰ Zn	³¹ Ga	³² Ge	33 As	³⁴ Se	35 Br	36 Kr	
37 Rb	³⁸ Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	⁵¹ Sb	⁵² Te	53 	54 Xe	
55 Cs	56 Ba	57 La	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 TI	82 Pb	83 Bi	⁸⁴ Po	At 85	⁸⁶ Rn	
87 Fr	⁸⁸ Ra	⁸⁹ Ac	¹⁰⁴ Unq	¹⁰⁵ Unp	106 Unh	¹⁰⁷ Uns	¹⁰⁸ Uno	¹⁰⁹ Une	110 Unn	■ poor metals								
alkali metals																		
			58 Ce	59 Pr	60 Nd	⁶¹ Pm	62 Sm	Eu	64 Gd	⁶⁵ Tb	66 Dy	67 Ho	68 Er	69 Tm	⁷⁰ Yb	71 Lu		
			90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr		

rare earth metals

Fig. 1.3 The Periodic Table of the elements.

actively mined given their heightened use in modern technology [16]. Many of the actinides, including uranium (U), plutonium (Pu), are used or produced in the nuclear industry, while others, such as thorium (Th) and protactinium (Pa) are formed naturally from the decay of U in the environment and have been used as geochemical tracers. The elements that are the primary focus of the book are contained in Table 1.1, along with some key details about each element. The Periodic Table (Fig. 1.3) and the information in Table 1.1 provide an indication of the differences between these elements. While Fe is not truly a trace element, being one of the most abundant elements on Earth, it is a metal that has been shown to be a limiting nutrient to oceanic phytoplankton [17]. This reason, and the fact that its chemistry plays a vital role in the overall fate and transport of many trace metals and metalloids argues for its inclusion in this book, along with Mn. The focus of the book will be on the metal(loid)s for which the most information is available, or which are the most environmentally relevant. While there will be a focus on metals of the first transition series, and all the metals of Groups 11 and 12, as they are either important

Table 1.1 Characteristics of the elements that will be principally discussed in this book. The average ocean and river concentrations for unpolluted waters are taken from various sources [4, 5, 17, 18].

Element	Symbol, Atomic Number	Atomic Mass	Abundance (μmol kg ⁻¹ or as noted)	Main Oxidation States	Average Ocean Concentration (nM)	Average River Concentration* (nM)	
Aluminum	Al 13	26.982	Most	III	20	1850	
Titanium	Ti 22	47.88	0.6%	IV	<20		
Vanadium	V 23	50.942	0.02%	II–V	30		
Chromium	Cr 24	51.996	2700	III, VI	4	19	
Manganese	Mn 25	54.938	0.085%	II, IV, VII	0.3	145	
Iron	Fe 26	55.847	2nd most	11, 111	0.5	720	
Cobalt	Co 27	58.933	500	11, 111	0.02	3.4	
Nickel	Ni 28	58.69	1500	I	8	8.5	
Copper	Cu 29	63.546	1100	II, I	4	24	
Zinc	Zn 30	65.39	1200	II	5	460	
Gallium	Ga 31	69.72	270	111	0.3	1.3	
Germanium	Ge 32	72.59	20	IV	0.07	0.1	
Arsenic	As 33	74.913	28	III, V	23	23	
Selenium	Se 34	78.96	0.6	–II, IV, VI	1.7		
Molybdenum	Mo 42	95.94	11	VI	110	5.2	
Silver	Ag 47	107.868	0.7	I	0.025	2.8	
Cadmium	Cd 48	112.41	1.3	II	0.6	0.2	
Tin	Sn 50	118.71	18	II, IV	0.004	0.01	
Antimony	Sb 51	121.75	1.7	III, V	1.6	8.2	
Tellurium	Te 52	127.60	0.008	-II, IV, VI	<0.001		
Cerium	Ce 58	140.12	430	III, IV	0.02		
Other Lanthanides	59–71	140.91-		II, III	La 0.03	La 0.4	
		174.97					
Tungsten	W 74	183.85	0.6	IV-VI	0.06		
Rhenium	Re 75	186.207	0.016	–I, VI, VII	0.04		
Osmium	OS 76	190.2	0.011	II, III, IV	0.05		
Platinum	Pt 78	195.08	0.19	II, IV	0.003		
Gold	Au 79	196.967	0.015	1. 111	<<0.001	0.01	
Mercury	Hg 80	200.59	0.33	0, 11	0.002	0.01	
Lead	Pb 82	207.2	50	II.	0.01	0.48	
Bismuth	Bi 83	208.98	0.12	III, V	<0.025		
Polonium	Po 84	(209)	No data	II, IV, –II	<<0.001		
Thorium	Th 90	232.038	26	IV	<0.001	0.4	
Protactinium	Pa 91	231.036			-		
Uranium	U 92	238.029	7.6	VI	14	1.0	
Plutonium	Pu 94	244.064					

*Average for unpolluted rivers. For some elements, variability in river concentrations is high.

commercially, or are toxic/required elements, it is also necessary to discuss the radioactive and rare earth elements. Few of the elements from the second and third transition series have important biochemical or biogeochemical reactions, besides molybdenum (Mo), and possibly tungsten (W), but some of these elements, such as rhenium (Re) and osmium (Os) have been used as tracers of environmental processes. Overall, the elements of the second and third transition series have been little studied. The lanthanides, as a group, have been used by aquatic chemists as chemical tracers, and have similar chemistry, because, while the inner 4f orbital is being filled through the series, the electrons lost are the outer orbital electrons and they mostly behave as +3 cations in solution. For the actinides, which are often radioactive, the focus will be on the chemistry of U, Th, Pa, Pu as well as the products of nuclear fission reactions, which are often radioactive [4].

There are many texts that cover one or more aspects of the topic of this book, which are referred to throughout, such as Stumm and Morgan [2], Morel and Hering [10], Cotton and Wilkinson [18], Drever [3], Buffle [19], Langmuir [4] and Wilkinson and Leads [20]. A compilation by Holland and Turekian [21] and authors therein, *Treatise on Geochemistry*, synthesizes in the various volumes many of the topics that are contained in this book. Some books are specific to ocean metal chemistry [1, 5].

1.2 Historical problems with metal measurements in environmental media

While the concentrations of metals in solid phase media rocks, sediments - and in biota are mostly sufficiently high that they can be analyzed using "typical" conditions found in a geochemical laboratory, this is not always the case, and it is certainly not true for the more trace constituents such as Cd, Hg, and Ag, and especially when making measurements in natural waters. Additionally, many other metals, such as Fe, Zn, and Cu, whose concentration in dust and in general laboratory materials (e.g., acids, plastics) is elevated, are difficult to accurately quantify without the proper scientific care. The potential for sample contamination during collection and analysis in many early environmental studies was not initially appreciated prior to the 1980s, and subsequently the techniques and approaches applied to the analysis of rocks were found not to be suitable for the analysis of natural water samples [22]. The difficulty in making measurements of the actual concentration of a metal in the water samples when collected, compared with the concentration in the sample bottle at the time of analysis, was not initially appreciated by many environmental scientists [23-25], and this still could cause erroneous results. Differences in concentration could result from any inadvertent addition of metal during sampling, handling, and analysis, and due to contaminants in the bottle plastics and in the reagents used.

Losses, due to adsorption to container walls are a problem for some metals in water samples, and the subsequent leaching of these metals back into solution can produce sampling artifacts. The impact of all these factors on the measurement of the dilute metal concentrations in natural solutions (Table 1.1) was not fully appreciated in the early environmental studies of natural waters.

The first convincing demonstration of the errors in many of the measurements of trace metals in environmental waters was made by Patterson and Settle [22]. Their landmark paper, The Reduction of Orders of Magnitude Errors in Lead Analyses of Biological Materials and Natural Waters by Evaluating and Controlling the Extent and Sources of Industrial Lead Contamination Introduced during Sample Collection and Analyses, [22] fundamentally changed the approach to the collection of samples of natural waters for trace element analysis, and lead to the development of protocols, so-called "clean techniques", for the analysis of low level trace metal samples. These investigators, and the scientists who adopted their approaches, clearly demonstrated the potential for contamination of samples due to traces of metals in the plastic materials, the acids and other reagents used in storage and analysis, and the importance of excluding dust in laboratory settings and the potential for introduction of metals from humans during handling. They showed the extent of preparation and care that was needed at all stages of the process: in sample container cleaning and preparation, during sample collection and manipulation, and during analysis. Such precautions that are now routinely used by environmental scientists (acid cleaning of glassware and plastics, use of gloves, cleaning of sampling devices and sampling lines, use of specific materials for the contaminant prone metals (e.g., use of Teflon for Hg sampling), and use of non-metallic materials where possible) were not done routinely in the early studies.

Patterson and Settle [22] showed that their measurements of Pb in ocean waters were orders of magnitude lower than those reported by other investigators and furthermore, that their results were "environmentally and geochemically consistent". This realization of the magnitude and extent of the contamination of samples subsequently lead to a revolution in the way that samples were collected and analyzed, and, as a result, in the reported concentrations of metals in environmental waters. This is illustrated in Fig. 1.4 for a number of metals. In Fig. 1.4(a), a compilation of data from various sources shows the dramatic decrease in the reported concentration for the metals Cu, Fe, Zn, and Hg in open ocean seawater over time [49-52]. In all cases, the change between the reported concentrations in 1983 and 1990 are relatively small compared to the dramatic decrease between 1965 and 1983, especially for Fe, Hg, and Zn. For Cu, the reported concentration decrease is about a factor of 50, while is it more than two orders of magnitude for the other metals. Similar decreases in the reported metal concentrations for river water are illustrated by a comparison over time [23] (see Fig. 1.4b). Again, the use of clean approaches to



Fig. 1.4 (a) Graph showing the trend in the reported mean concentration of several metals in ocean waters. Early data are taken from reviews by Brewer [24]; Bruland [49]; Bruland et al. [40] and Goldberg [25]. The 1990 data are taken from Refs [50, 52] and are for the Pacific Ocean. Figure redrawn using this information and Mason [26]; (b) reported mercury concentrations for Vandercook Lake in Wisconsin over time. Figure reprinted

sampling and analysis lead to more than an order of magnitude decrease in the measured concentrations of Cd and Pb in river water compared to the values reported previously. Finally, data from samples collected in a lake in Wisconsin show a similar trend for Hg [27] (Fig. 1.4c). The decrease in reported concentration was again more than two orders of magnitude between 1983–6. As noted earlier, the changes in the reported concentration in all these instances was not primarily the result of a "cleanup" of the environment, although there is evidence for changing concentrations over recent decades, but rather a realization and appreciation that the investigators themselves, and the plastics and other materials used for sampling contain significant traces of metals, and that these were being leached into the samples during handling and storage.

For the ocean, sampling methodologies that used metal wire cables led to contamination, as did the use of untreated

from Fitzgerald, W. and Watras, C.J. (1989) *Science of the Total Environment* **87/88**: 223–32 with permission of Elsevier [27]. (c) Figure showing the difference between the reported concentrations of lead and cadmium for the Quinnipiac River. Figure reprinted with permission from Benoit, G. (1994) *Environmental Science and Technology* **28**: 1987–91, Copyright (1994), American Chemical Society [23].

plastic sampling bottles used to collect samples at depth (e.g., Niskin bottles). Metal inputs from the vessels themselves can also be a source of contamination. These sources of contamination were most severe for those waters that have low concentration due to their remoteness (e.g., the middle of the Pacific Ocean). A comparison by a reader of the reported concentrations of metals in the ocean in the recent literature would further illustrate the point. Such an examination indicates that while the reported concentrations of Al (~40nM), Cr (40nM) and other metals that are present in seawater at relatively high concentrations are similar to those in Table 1.1, many of the concentrations reported in various books are 2-100 times higher, especially for the heavy metals (e.g., Hg, Pb) and even for some of the transition metals (Ni, Co, Zn). So, there is still a need for caution when reading the literature and certainly much care and application is needed when doing field studies.

The extent of potential sample contamination is obviously, but not exclusively, a function of the concentration, with potentially more contamination likely for metals that are present at low concentrations. Cadmium provides one example where this is not true as, even though it exists in natural waters at low concentrations, there has been less evidence of its contamination in earlier studies. Sampling and analytical difficulties are most severe for those metals that are present in low concentration and which are routinely used in materials, or are part of typical sampling equipment and vessels, or occur in high abundance in atmospheric aerosols, such as Fe, Cu, Zn, Hg, and Pb. One example is the use of metal-containing dyes in plastics, such as in the coloring of pipette tips. Use of these tips with acidic solutions can lead to metal leaching and contamination. One recent example of contamination for Zn that has been highlighted in ocean studies is the use of Zn as "sacrificial anodes" on metal sampling devices to prevent corrosion - dissolution of Zn was found to be sufficient to contaminate samples during a recent international intercalibration study as part of the GEOTRACERS program [28].

The introduction of these approaches has led to a wealth of information on metal distributions in the ocean and in freshwaters that has resulted in an enhanced understanding of their toxicity to aquatic organisms, and their requirement as essential nutrients, and to the factors that control their cycling. Accurate measurements also led to a clearer and better understanding of the role of human inputs in environmental contamination, and the need to curb industrial and other emissions of metals to the atmosphere and aquatic systems. The importance of this development and the result-

ant explosion of high quality research can be illustrated by the papers presented at a special symposium held in honor of Claire Patterson on his retirement, in December 1992. The papers presented at the meeting were published in a special issue of Geochimica et Cosmochimica Acta in 1994 and many papers cover topics related to Pb and other metals, and their geochemical cycle, such as using ice core records and ocean measurements and Pb isotope ratios to examine recent changes and those over geological timescales [29-37]. The paper by Boutron et al. [29] illustrates the difficulty in making measurements of trace concentrations of metals in a situation where the ice core needed to be drilled and recovered using contaminating techniques. These authors measured the concentration of the metals in successive lavers of ice shaved from the outside to the inside of the ice core and showed that while the outer layers were highly contaminated, the inner core was not (Fig. 1.5). This analysis further provided a clear record of the impact of human use and release of Pb, primarily from its use as a gasoline additive, on the concentrations of Pb in remote precipitation (Fig. 1.6) [29].

Other papers presented at the symposium, such as the review paper of Bruland et al. [40], detail measurements of reactive trace metals (Al, Fe, Mn, Zn and Cd) in ocean waters. These authors discussed the notion, first proposed by Whitefield and Turner [38], that the profiles of such elements reflect their biogeochemistry. Some metals such as Zn and Cd have so-called "nutrient (recycled) profiles" in the ocean as their concentration is depleted in surface waters and higher at depth, while others, such as Al, reflect their primary atmospheric input and their high association with

Fig. 1.5 The measured concentrations of lead and cadmium in successive layers of ice shaved from an ice core recovered from Vostok Station in East Antarctica. Figure reprinted from Boutron et al. (1994) *Geochimica et Cosmochimica Acta* **58**: 3217–25 with permission of Elsevier [29].

Fig. 1.6 Lead profiles for ice cores collected in polar region. Figure reprinted from Boutron et al. (1994) *Geochimica et Cosmochimica Acta* 58: 3217–25 with permission of Elsevier [29].

the particulate phase (so-called "scavenged profiles"; high surface concentrations and low deep water values). Other metals, such as Hg, have a more complex ocean distribution due to their complex chemistry and cycling [39].

The notion that a metal, such as Zn, could have an ocean profile that is similar to that of the major nutrient elements [40] (i.e., nitrogen (N), phosphorous (P), and silica (Si) – required by diatoms), and that the concentration of Zn and Si appear closely correlated for open ocean waters [17] was reconciled with the knowledge that Zn has a biological role and forms part of many enzymes important to cellular biochemistry [11]. At the time, the correlation of Cd and P profiles was more of a "mystery" as Cd was considered a toxic heavy metal with no biological function and there was "no biological or chemical explanation for the (relationship)" [1]. It has since been shown that Cd can substitute for Zn in one important enzyme for ocean photosynthetic organisms (carbonic anhydrase) and therefore its role as a nutrient has been established [11].

It is worth noting that contamination is not only a problem in trace metal research where quantification of *in situ* concentrations is the principal focus. Fitzwater et al. [41] demonstrated, for example, that the introduction of metals into solutions from container vessels and reagents could lead to substantial under-estimation of the primary productivity when using ¹⁴C-incubation approaches. They demonstrated the potential for leaching of Cu from glass bottles even after acid washing, and the converse, the adsorption of Cu onto glass containers in unacidified solutions. High metal levels were found in the standard ¹⁴C solutions and in other solutions used in productivity studies. As a result, more recent techniques have been devised that use trace metal free "clean techniques" [42] for productivity and other studies with phytoplankton and other microbes, if experiments are performed under natural conditions.

The environmental scientist is therefore strongly cautioned to check the reported concentrations in any publication against other sources as, while there is obviously variability between ecosystems and especially for systems with high or variable TSS (total suspended solids) loads, there is still the possibility that reported concentrations in the literature are not reflective of actual concentrations. However, on the contrary, while it is often considered that the lowest concentration must be correct, this is not always the case, as losses and incomplete recovery of the metal during sample analysis is a real possibility, and therefore this metric should not be used exclusively to gauge whether a reported value is correct or not.

Technique	Linear Range	Precision	DL Cd (nM)	DL Cu (nM)	DL Zn (nM)	Interferences
GFAA	~10 ²	0.5–5 %	0.02	0.2	0.3	Relatively few
ICP-AE	~105	<2%	0.9	6	3	Spectral; many lines overlap
ICP-MS	10 ⁵ -10 ⁷	1–3%	10 ⁻³	$3 imes 10^{-3}$	$4.5 imes 10^{-3}$	Many; isotopes, oxides and other molecular mass

Table 1.2 Comparison of techniques for determining metal concentrations in water samples showing relative precision and other analytical metrics [46].

1.3 Recent advances in aquatic metal analysis

While the previous section outlined the historical problems associated with sample contamination, this issue has been mostly addressed and most recent studies have demonstrated the ability to accurately quantify the total concentration of an element in environmental waters. Some recent intercalibration exercises in open ocean seawater indicate that contamination is always a possibility, and care and application of "clean techniques" is necessary as the concentrations are low (nM or pM; Table 1.1). Comparison and confirmation of accuracy can be achieved through the continued use and further development of low level reference materials. Standard (or Certified) Reference Materials (SRMs or CRMs) are environmental materials that have been repeatedly measured, often by more than one technique, so that their concentration is well-defined by an accepted value and a standard deviation which reflects the overall precision of the measurement. Such materials allow an analyst to confirm that the method and approach used is providing an accurate answer. The US National Institute for Standards and Technology (NIST), the European Institute for Reference Materials and Measures (IRMM), and other organizations in other countries around the world, provide an impressive array of SRMs in many different media. However, it is often difficult to provide adequate SRMs for the low levels of many metals in seawater or freshwaters (nM or lower concentrations), and the number of SRMs for water are limited. There are recent concerted efforts to develop such SRMs, and the international GEOTRACERS Program [43] is one avenue through which such reference materials are being developed.

While accurate measurement of the total concentration is a goal, there is an acknowledgement that the total concentration of an element in solution is not often an accurate predictor of the bioavailability, toxicity, or reactivity of the element [2–4, 10], as most metals do not exist in solution purely as the free metal ion, and are often associated with dissolved organic matter or can be in colloidal phases in the filtered fraction, as discussed in Chapter 2. Therefore, protocols have been developed to examine the distribution and speciation (chemical form) of the metal(loid) in solution, as well as isotope ratios and other more detailed analysis [2, 10, 40, 44, 45]. Analytical advances and improvements in sensitivity have also improved the amount of information that is collected. For example, graphite furnace atomic adsorption spectroscopy (GFAAS) was often the preferred method for trace metal analysis in the early 1980s but this method is slow as each element requires a different lamp, and the linear calibration range is small, often requiring reanalysis of samples [45]. The analysis is also time-consuming although there is the potential to use an autosampler. Additionally, extensive pretreatment is required to concentrate samples to levels sufficient for analysis, and to remove interferences from the matrix, although techniques, such as Zeeman correction, have evolved to reduce these issues. As shown in Table 1.2 [46], the detection limits (DL) of GFAAS for the metals shown are within an order of magnitude of their concentrations in seawater (Table 1.1), and so sample preconcentration and/or matrix elimination is often required. The development of ICP (inductively coupled plasma) as an atomization technique has allowed for the more rapid analysis of samples although when coupled with atomic emission quantification (ICP-AES) results in a higher detection limit which is unsuitable for many metals in environmental waters, or even in sediment extracts (Table 1.2). However, coupling to a mass spectrometer (ICP-MS) provides high throughput, and low DL with high precision. However, ICP-MS is prone to substantial interferences such as different elements with the same atomic mass, or interferences due to the formation of oxides and other molecules that form after atomization (e.g., ⁴⁰Ar¹⁶O and ⁵⁶Fe; ⁴⁰Ar³⁵Cl and ⁷⁵As) but there are a variety of approaches to circumvent these interferences [47]. The severity of the interferences depends on the resolution of the MS, which is relatively low (0.8 amu) for a quadrupole ICP-MS, which are the most robust and stable instruments. Therefore, analysis using these instruments needs to contend with interferences. One approach that has been used to circumvent molecular interferences is the use of a reaction cell, which is designed to break up diatomic and polyatomic molecules prior to the MS, thereby removing the interferences [47]. High resolution instruments are also available that are capable of resolving these interferences because of their much higher resolving power, such as that with ⁴⁰Ar¹⁶O and ⁵⁶Fe, and these are used in many instances. However, these instruments are more delicate and require more maintenance than a quadrupole [47]. More recently, multicollector ICP-MS instruments have been more widely used in aquatic metal analysis as these instruments are able to quantify the isotope ratios of specific elements and can be used to examine isotope fractionation and the factors causing such fractionation. Such isotope fraction can be used to examine environmental processes in a similar fashion to the more widely used fractionation of the major organic elements (C, N, S). Isotope analysis provides a new avenue for research and exploration as many abiotic and biotic processes can cause isotope fractionation in the environment. This is discussed further in Section 7.5. Additionally, ICP-MS is now being used to analyze radioisotopes in natural waters (e.g., Th, Pa, U, Pu) as this can be done with more accuracy and precision than can be obtained with conventional counting techniques.

Therefore, the amount of information has expanded rapidly with the development of more accurate and precise measurement techniques and with the ability to sample at higher resolution. Many of the factors that caused contamination in early studies have been overcome and sampling approaches that were not feasible 20 years ago, such as sampling for metals in seawater with an automated sampling system (rosette sampler), are now being used [43, 48], resulting in a higher throughput of sample collection. In freshwater systems, automated water samplers, such as Isco devices (www.isco.com), are now used to collect samples without contamination. In addition to intercalibration studies, which have allowed for more detailed understanding of sources of sampling and analytical error, methods have been developed to examine the distribution of metals between dissolved, colloidal and particulate fractions without contamination, using ultrafiltration and other approaches. Analytical class-100 clean rooms are now routinely used for most metal analysis and high precision instrumentation has also allowed for a better characterization of blanks and sources of error.

As concentrations in water samples are generally too low for direct analysis, sample preconcentration often occurs. This can additionally remove potential matrix effects and other interferences. Preconcentration techniques typically rely on [44]: (1) co-precipitation of insoluble (hydr)oxides (e.g., Fe, Mn, Mg); or (2) solvent or solid-phase extraction of the analyte from the matrix using organic ligands and organic solvents or onto chelating resins, with subsequent elution into a small volume. Analysis on board ship or in the field often relies on techniques other than ICP-MS as it is not transportable. Many of these analyses rely on flow injection approaches, which allow for preconcentration prior to detection using spectrometry of a metal-ligand complex, or after reaction of the metal with a complex that changes the signal, as occurs, for example, with the reaction of Fe^{II} or Mn^{II} with luminol [44]. Alternatively, as discussed in Chapter 6, many metals can be determined using electrochemical techniques using mercury drop electrodes.

One example of a detailed approach to standardizing methods and analysis is the GEOTRACERS Program intercalibration study [28, 43]. This program has provided a detailed manual on how to collect ocean water samples for the determination of a suite of metals, metalloids, radionuclides and other tracer compounds (TEIs). The manual that has been developed covers the sampling, filtration and storage with details on which type of equipment, filters and other methods to be used, and the best container and sample preservation protocols. These methods are for both radioac-tive elements as well as stable trace metal(loid)s and other elements, and their stable isotopes; see Section 6.4.2 for further details.

Finally, the development of *in situ* detection methods for the analysis of metal(loid)s in environmental waters is in its infancy but there is the promise of a further explosion of information as such techniques become available. It will be apparent through the discussions in the later chapters of the book, especially Chapters 6 and 7, that there has been enormous progress in recent decades in the understanding of the fate, transport and transformation of metal(loid)s in aquatic systems and the dynamics of the interactions between metals in solution and the solid phase, and with microorganisms that inhabit these waters. Many of the recent discoveries would not be possible without the development of new techniques and approaches, including the methods to quantify different fractions of an element in aqueous systems.

References

- 1. Broecker, W. and Peng, T.-H. (1982) *Tracers in the Sea*. Columbia University Press, Palisades, NY.
- Stumm, W. and Morgan, J.J. (1996) Aquatic Chemistry. John Wiley & Sons, Inc., New York.
- 3. Drever, J. (1997) *The Geochemistry of Natural Waters*. Prentice Hall, Upper Saddle River.
- Langmuir, D. (1997) Aqueous Environmental Geochemistry. Prentice Hall, Upper Saddle River.
- 5. Chester, R. (2003) *Marine Geochemistry*. Blackwell Science, Oxford.
- Sillen, L. (1967) The ocean as a chemical system. *Science*, **156**, 1189–1197.
- McDuff, R. and Morel, F.M.M. (1980) The geochemical control of seawater (Sillen revisited). *Environmental Science & Technology*, 14, 1182–1186.
- 8. Doney, S.C. (2006) The dangers of ocean acidification. *Scientific American*, **294**, 58–65.
- Orr, J.C., Fabry, V.J., Aumont, O. et al. (2005) Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*, **437**, 681–686.

- 10. Morel, F.M.M. and Hering, J.G. (1993) *Principles and Applications* of Aquatic Chemistry. John Wiley & Sons, New York.
- 11. Morel, F., Milligan, A.J. and Saito, M.A. (2004) Marine Bioinorganic chemistry: The role of trace metals in the oceanic cycles of major nutrients. In: Elderfield, H. (ed.) *The Oceans and Marine Geochemistry*. in Holland, H.D. and Turekian, K.K. (Exec eds) Treatise on Geochemistry. Elsevier Pergamon, Oxford, pp. 113–143.
- Lindh, U. (2005) Biological functions of the elements. In: Selinus, O., Alloway, B., Centeno, J.A. et al. (eds) Chapter 5, *Essentials of Medical Geology*. Elsevier, Amsterdam, pp. 115–160.
- Fordyce, F. (2005) Selenium deficiency and toxicity in the environment. In: Selinus, O., Alloway, B., Centeno, J.A. et al. (eds) Chapter 15, *Essentials of Medical Geology*. Elsevier, Amsterdam, pp. 373–416.
- 14. Craig, P.J. (ed.) (2003) Organometallic Compounds in the Environment. John Wiley & Sons, Ltd, Chichester.
- Nozaki, Y. (2010) Rare earth elements and their isotopes in the ocean. In: Steele, J.H., Thorpe, S.A. and Turekian, K.K. (eds) *Marine Chemistry and Geochemistry*. Elsevier, Amsterdam, pp. 39–51.
- USGS (2011) Rare earth elements end use and recyclability, USGS Scientific Investigations Report 2011–5094. http:// pubs.usgs.gov/sir/2011/5094/pdf/sir2011-5094.pdf (accessed April 16, 2011).
- Bruland, K. and Lohan, M.C. (2004) Controls on trace metals in seawater. In: Elderfield, H. (ed.) *The Oceans and Marine Geochemistry*. in Holland, H.D. and Turekian, K.K. (Exec eds) Treatise on Geochemistry. Elsevier Pergamon, Oxford, pp. 23–47.
- Cotton, F.A. and Wilkinson, G. (1972) Advanced Inorganic Chemistry: A Comprehensive Treatise. John Wiley & Sons, Inc., New York, 1355 pp.
- Buffle, J. (1988) Complexation Reactions in Aquatic Systems. An Analytical Approach. Ellis Horwood Pub., Chichester, 700 pp. (downloadable at http://www.unige.ch/cabe/ (accessed April 16, 2011).
- Wilkinson, K.J. and Lead, J.R. (eds) (2007) Environmental Colloids and Particles. Behaviour, Separation and Characterization. IUPAC Series in Analytical and Physical Chemistry of Environmental Systems, Buffle, J. and van Leeuwen, H.P., Series Editors. John Wiley & Sons, Ltd, Chichester.
- Holland, H.D. and Turekian, K.K. (2004) Treatise on Geochemistry, in 10 Volumes. Elsevier, Oxford.
- 22. Patterson, C.C. and Settle, D.M. (1976) The reduction in orders of magnitude errors in lead analysis of biological materials and natural waters by evaluating and controlling the extent and sources of industrial lead contaminantion introduced during sample collection and analysis. In: LaFleur, P. (ed.) *Accuracy in Trace Analysis: Sampling, Sample Handling and Analysis.* US National Bureau of Standards, Washington, D.C., pp. 321–351.
- Benoit, G. (1994) Clean technique measurement of Pb, Ag and Cd in freshwater: A redefinition of metal pollution. *Environmental Science and Technology*, 28, 1987–1991.
- Brewer, P. (1975) Minor elements in seawater. In: Riley, J. and Skirrow, G. (eds) *Chemical Oceanography*. Academic Press, London, pp. 415–496.

- Goldberg, E. (1965) Minor elements in seawater. In: Riley, J. and Skirrow, G. (eds.) *Chemical Oceanography*. Academic Press, London, pp. 163–196.
- 26. Mason, R. (1991) The Chemistry of Mercury in the Equatorial Pacific Ocean. University of Connecticut, Storrs-Mansfield, CT.
- Fitzgerald, W. and Watras, C.J. (1989) Mercury in surficial waters of rural Wisconsin lakes. *Science of the Total Environment*, 87/88, 223–232.
- Geotracers (2011) The GEOTRACERS Cookbook. http:// www.geotraces.org/science/intercalibration/222-samplingand-sample-handling-protocols-for-geotraces- (September 25, 2012).
- 29. Boutron, C.F., Candelone, J. and Hong, S.M. (1994) Past and recent changes in the large-scale tropospheric cycles of lead and other heavy-metals as documented in Antarctic and Greenland snow and ice a review. *Geochimica et Cosmochimica Acta*, **58**, 3217–3225.
- Rosman, K.J.R., Chisholm, W., Boutron, C.F., Candelone, J. and Hong, S. (1994) Isotopic evidence to account for changes in the concentration of lead in Greenland snow between 1960 and 1988. *Geochimica et Cosmochimica Acta*, 58, 3265– 3269.
- 31. Erel, Y. and Patterson, C.C. (1994) Leakage of industrial lead into the hydrocycle. *Geochimica et Cosmochimica Acta*, **58**, 3289–3296.
- Hirao, Y., Matsumoto, A., Yamakawa, H., Maeda, M. and Kimura, K. (1994) Lead behavior in abalone shell. *Geochimica et Cosmochimica Acta*, 58, 3183–3189.
- Veron, A.J., Church, T.M., Patterson, C.C. and Flegal, A.R. (1994) Use of stable lead isotopes to characterize the sources of anthropogenic lead in North-Atlantic surface waters. *Geochimica et Cosmochimica Acta*, **58**, 3199–3206.
- Boyle, E.A., Sherrell, R.M. and Bacon, M. (1994) Lead variability in the western North-Atlantic Ocean and central Greenland ice – implications for the search for decadal trends in anthropogenic emissions. *Geochimica et Cosmochimica Acta*, 58, 3227– 3238.
- Ritson, P.I., Esser, B.K., Niemeyer, S. and Flegal, A.R. (1994) Lead isotopic determination of historical sources of lead to Lake Erie, North-America. *Geochimica et Cosmochimica Acta*, 58, 3297– 3305.
- Rivera-Duarte, I. and Flegal, A.R. (1994) Benthic lead fluxes in San-Francisco Bay, California, USA. *Geochimica et Cosmochimica Acta*, 58, 3307–3313.
- Sanudo-Wilhelmy, S.A. and Flegal, A.R. (1994) Temporal variations in lead concentrations and isotopic composition in the southern California Bight. *Geochimica et Cosmochimica Acta*, 58, 3315–3320.
- Whitefield, M. and Turner, D.R. (1987) The role of particles in regulating the composition of seawater. In: Stumm, W. (ed.) *Aquatic Surface Chemistry*. John Wiley & Sons, Inc., New York, pp. 457–493.
- Mason, R.P., Fitzgerald, W.F. and Morel, F.M.M. (1994) The biogeochemical cycling of elemental mercury – Anthropogenic influences. *Geochimica et Cosmochimica Acta*, 58(15), 3191– 3198.
- 40. Bruland, K.W., Orians, K.J. and Cowen, J. (1994) Reactive trace-metals in the stratified central North Pacific. *Geochimica et Cosmochimica Acta*, **58**, 3171–3182.

- 41. Fitzwater, S., Knauer, G.A. and Martin, J.H. (1982) Metal contamination and its effects on primary production measurements. *Limnology and Oceanography*, **27**, 544–551.
- Price, N., Harrison, G.I., Hering, J.G., Hudson, R.J., Nirel, P., Palenik, B. and Morel, F.M.M. (1988) Preparation and chemistry of the artificial algal medium Aquil. *Biological Oceanography*, 5, 43–46.
- 43. Henderson, G..M., Anderson, R.F., Adkins, J., et al. (SCOR Working Group) (2007) GEOTRACES – An international study of the global marine biogeochemical cycles of trace elements and their isotopes. *Chemie der Erde-Geochemistry*, **67**, 85–131.
- Sohrin, Y. and Bruland, K.W. (2011) Global status of trace elements in the ocean. *Trends in Analytical Chemistry*, **30**, 1291–1307.
- 45. Harrington, C.F., Clough, R., Drennan-Harris, L.R., Hill, S.J. and Tyson, J.F. (2011) Atomic spectrometry update: Elemental speciation. *Journal of Analytical Atomic Spectrometry*, **26**, 1561– 1595.
- 46. Thermo (2012) ICP or ICP-MS? Which technique should I use? An elementary overview of Elemental Aalysis. Available at http://www.thermo.com/eThermo/CMA/PDFs/Articles/ articlesFile_18407.pdf (accessed October 29, 2012).
- 47. Linge, K.L. and Kym, J.K. (2009) Quadrupole ICP-MS: Introduction to instrumentation: Measurement techniques and analytical capabilities. *Geostandards and Geoanalytical Research*, **33**, 445–467.
- Measures, C.I., Landing, W.M., Brown, M.T. and Buck, C.S. (2008) A commercially available rosette system for trace metalclean sampling. *Limnology & Oceanography-Methods*, 6, 384–394.
- Bruland, K. (1983) Trace elements in seawater. In: Riley, J. and Chester, R. (eds) *Chemical Oceanography*. Academic Press, London, pp. 157–215.
- 50. Coale, K. and Bruland, K.H. (1990) Spatial and temporal variability in copper complexation in the North Pacific. *Deep-Sea Research Part I*, **37A**, 809–832.
- 51. Landing, W. and Bruland, K.H. (1987) The contrasting biogeochemistry of iron and manganese in the Pacific Ocean. *Geochimica et Cosmochimica Acta*, **51**, 29–43.
- 52. Mason, R.P. and Fitzgerald, W.F. (1990) Alkylmercury species in the equatorial Pacific. *Nature*, **347**, 457–459.

Problems

1.1. Discuss briefly the chemistry of the elements that form the "rare earths" and provide brief details of the use of four of these elements in industrial applications.

- **1.2.** A chemist makes a standard solution for the analysis of metals by ICP-MS. She wants the final standards to cover the range from one tenth to 10 times the average concentration of the element in seawater. To do this she will make a composite stock solution that is 1μ M in the following metals (Co, Cu, Zn, Cd, Ag) and 100 nM in Pb and Hg. How much metal does she need to weigh out for each element, using the anhydrous chloride salt (e.g., CuCl₂) given that the final solution will be made in 11 of acidified water?
- **1.3.** Examine the concentrations of the elements in Table 1.1 and summarize any evident trends between the concentrations in seawater versus those in river water. Discuss why this may be so, and comment on the elements that do not fit the trend and why you may suspect they do not.
- 1.4.
 - **a.** If the pH of seawater is 8.1 and the pH is mostly controlled by the acid-base chemistry of the carbonate system, calculate the concentration of dissolved carbonate assuming the total carbonate concentration (all species) is 2 mM. The following acid-base constants are appropriate for this problem:

 $H_2CO_3 = H^+ + HCO_3^ K = 10^{-6.1}$ $HCO_3^- = H^+ + CO_3^{2-}$ $K = 10^{-9.3}$

Equilibrium constants have been adjusted to account for ionic strength effects.

(Hint: given the pH, it is reasonable to assume that the bicarbonate concentration is approximately equivalent to the total carbonate concentration.)

- **b.** If the pH decreased to 7.9, how much would the carbonate concentration change?
- **1.5.** It has been stated that the concentrations of Fe and Mn in seawater are controlled by the precipitation of their solid phases. Estimate the concentrations based on the precipitation of Fe(OH)₃ and MnO₂ and compare these values to those in Table 1.1. If the concentrations are different, provide a reasonable explanation for the differences.