

LECTURE 1

An Overview

Immunology is a difficult subject to study for several reasons. First, there are lots of details, and sometimes these details get in the way of understanding the concepts. To get around this problem, we're going to concentrate on the big picture – it will be easy for you to find the details somewhere else. A second difficulty in learning immunology is that there is an exception to every rule. Immunologists love these exceptions, because they give clues as to how the immune system functions. But for now, we're just going to learn the rules. Oh, sure, we'll come upon exceptions from time to time, but we won't dwell on them. Our goal will be to examine the immune system, stripped to its essence. The third difficulty in studying immunology is that our knowledge of the immune system is still evolving. As you'll see, there are many unanswered questions, and some of the things that seem true today will be proven false tomorrow. I'll try to give you a feeling for the way things stand now, and from time to time I'll discuss what immunologists speculate may be true. But keep in mind that although I'll try to be straight with you, some of the things I'll tell you will change in the future (maybe even by the time you read this!).

I think the main reason immunology is such a tough subject, however, is that the immune system is really a "network" that involves many different players interacting with each other. Imagine you're watching a football game on TV, and the camera is isolated on one player, say, the tight end. You see him run full speed down the field, and then stop. It doesn't seem to make any sense. Later, however, you see the same play on the big screen, and now you understand: That tight end took two defenders with him down the field, leaving the running back uncovered to catch the pass and run for a touchdown. The immune system is just like a football team. It's a network of players who cooperate to get things done, and just looking at one player doesn't make much sense. You need an overall

view. That's the purpose of this first lecture, which you might call "turbo immunology." Here, I'm going to take you on a quick tour of the immune system, so you can get a feeling for how it all fits together. Then in the next lectures, we'll go back and take a closer look at the players and their interactions.

PHYSICAL BARRIERS

Our first line of defense against invaders consists of physical barriers, and to cause trouble, viruses, bacteria, and parasites must first penetrate these shields. Although we tend to think of our skin as the main barrier, the area covered by our skin is only about two square meters. In contrast, the area covered by the mucous membranes that line our digestive, respiratory, and reproductive tracts measures about 400 square meters – an area about as big as two singles-tennis courts. The point here is that there is a large perimeter which must be defended.


THE INNATE IMMUNE SYSTEM

Any invader that breaches the physical barrier of skin or mucosa is greeted by the innate immune system – our second line of defense. Immunologists call this system "innate" because it is a defense that all animals just naturally seem to have. Indeed, some of the weapons of the innate immune system have been around for more than 500 million years. The way the innate system works is pretty amazing.

Imagine you are getting out of your hot tub, and as you step onto the deck, you get a large splinter in your big toe. On that splinter are many bacteria, and within a few hours you'll notice (unless you had a lot to drink in that hot tub!)

2 LECTURE 1 An Overview


that the area around where the splinter entered is red and swollen. These are indications that your innate immune system has kicked in. In your tissues are roving bands of white blood cells that defend you against attack. To us, tissue looks pretty solid, but that's because we're so big. To a cell, tissue looks somewhat like a sponge with holes through which individual cells can move rather freely. One of the defender cells that is stationed in your tissues is the most famous innate immune system player of them all: the macrophage. If you are a bacterium, a macrophage is the last cell you want to see after your ride on that splinter. Here is an electron micrograph showing a macrophage about to devour a bacterium:



You will notice that this macrophage isn't just waiting until it bumps into the bacterium, purely by chance. No, this macrophage actually has sensed the presence of the bacterium, and is reaching out a "foot" to grab it. But how does a macrophage know that a bacterium is out there? The answer is that macrophages have antennae (receptors) on their surfaces which are tuned to recognize "danger molecules" characteristic of common microbial invaders. For example, the walls that surround bacteria are made up of fats and carbohydrates that normally are not found in the human body, and macrophages have receptors which can detect these "foreign" molecules. Not only do these antennae alert macrophages to danger, they also can cause a macrophage to "crawl" toward the microbe which is emitting the foreign molecule.

When it encounters a bacterium, a macrophage first engulfs it in a pouch (vesicle) called a phagosome. This vesicle is then taken inside the macrophage, where it fuses with another vesicle termed a lysosome. Lysosomes contain powerful chemicals and enzymes which can destroy bacteria. In fact, these agents are so destructive that they would kill the macrophage itself if they were released inside it. That's why they are kept in pouches. Using this clever strategy, the macrophage can destroy an invader

without "shooting itself in the foot." This whole process is called phagocytosis, and this series of snapshots shows how it happens:




Macrophages have been around for a very long time. In fact, the ingestion technique macrophages employ is simply a refinement of the strategy that amoebas use to feed themselves – and amoebas have roamed the earth for about 2.5 billion years.

Why is this creature called a macrophage, you may be wondering. *Macro*, of course, means "large" – and a macrophage is a large cell. *Phage* comes from a Greek word meaning "to eat." So a macrophage is a big eater. In fact, in addition to defending against invaders, the macrophage functions as a garbage collector. It will eat almost anything. Immunologists take advantage of this appetite by feeding macrophages iron filings. Then, using a small magnet, they can separate macrophages from other cells in a cell mixture. Really!

Where do macrophages come from? Macrophages and all the other blood cells in your body are made in the bone marrow, where they descend from self-renewing cells called stem cells – the cells from which all the blood cells "stem." By self-renewing, I mean that when a stem cell grows and divides into two daughter cells, it does a "one for me, one for you" thing in which some of the daughter cells go back to being stem cells, and some of the daughters go on to become mature blood cells. This strategy of continuous self-renewal insures that there will always be blood stem cells "in reserve" to carry on the process of making mature blood cells.

As each daughter cell matures, it has to make choices that determine which type of blood cell it will be when

it grows up. As you can imagine, these choices are not random, but are carefully controlled to make sure you have enough of each kind of blood cell. For example, some daughter cells become red blood cells, which capture oxygen in the lungs and transport it to all parts of the body. Indeed, our stem cell “factories” must turn out more than two million new red blood cells each second to replace those lost due to normal wear and tear. Other descendants of a stem cell may become macrophages, neutrophils, or other types of “white” blood cells. Just as white wine really isn’t white, these cells aren’t white either. They are colorless, but biologists use the term *white* just to indicate that they lack hemoglobin, and therefore are not red. Here is a figure showing some of the many different kinds of blood cells a stem cell can become:



When the cells which will mature into macrophages first exit the bone marrow and enter the bloodstream, they are called monocytes. All in all you have about two billion of these cells circulating in your blood at any one time. This may seem a little creepy, but you can be very glad they are there. Without them, you’d be in deep trouble. Monocytes remain in the blood for an average of about three days. During this time they travel to the capillaries, which represent the “end of the line” as far as blood vessels go, looking for a crack between the endothelial cells that line the capillaries. These cells are shaped like shingles, and by sticking a foot between them, a monocyte can leave the blood, enter the tissues, and mature

into a macrophage. Once in the tissues, most macrophages just hang out, do their garbage collecting thing, and wait for you to get that splinter so they can do some real work.

When macrophages eat the bacteria on that splinter, they give off chemicals which increase the flow of blood to the vicinity of the wound. The build-up of blood in this area is what makes your toe red. Some of these chemicals also cause the cells that line the blood vessels to contract, leaving spaces between them so that fluid from the capillaries can leak out into the tissues. It is this fluid which causes the swelling. In addition, chemicals released by macrophages can stimulate nerves in the tissues that surround the splinter, sending pain signals to your brain to alert you that something isn’t quite right in the area of your big toe.

During their battle with bacteria, macrophages also produce proteins called cytokines. These are hormone-like messengers which facilitate communication between cells of the immune system. Some of the cytokines alert monocytes and other immune system cells traveling in nearby capillaries that the battle is on, and influence these cells to exit the blood to help fight the rapidly multiplying bacteria. I should also point out that macrophages are not very tidy eaters. They frequently burp some of their meal back out into the tissues, and this debris also serves as a signal to recruit more defenders from the blood. Pretty soon you have a vigorous “inflammatory” response going on in your toe, as the innate immune system battles to eliminate the invaders.

When you think about it, this is a great strategy. You have a large perimeter to defend, so you station sentinels (macrophages) to check for invaders. When these sentinels encounter the enemy, they send out signals that recruit more defenders to the site of the battle. The macrophages then do their best to hold off these invaders until reinforcements arrive. Because the innate response involves warriors like macrophages that are programmed to recognize many of the most common invaders, your innate immune system usually responds so quickly that the battle is over in just a few days.

There are other players on the innate team, and we will talk about them at length in the next lecture. For example, in addition to cells like macrophages (the “professional phagocytes”), which make it their business to eat invaders, the innate system includes the complement proteins that can punch holes in bacteria, and natural killer cells which are able to destroy bacteria, parasites, virus-infected cells, and some cancer cells.

4 LECTURE 1 An Overview

THE ADAPTIVE IMMUNE SYSTEM

About 99% of all animals get along just fine with only natural barriers and the innate immune system to defend them. However, for vertebrates like us, Mother Nature laid on a third level of defense: the adaptive immune system. This is a defense system that actually can adapt to protect us against almost any invader. It is most likely that the adaptive immune system was designed to protect us and the other vertebrates against viruses, because as you will see, the innate immune system is not terribly effective against viruses.


One of the first clues that the adaptive immune system existed came back in the 1790s when Edward Jenner began vaccinating the English against smallpox virus. In those days, smallpox was a major health problem. Hundreds of thousands of people died from this disease, and many more were horribly disfigured. What Jenner observed was that milkmaids frequently contracted a disease called cowpox which caused lesions on their hands that looked similar to the sores caused by the smallpox virus. Jenner also noted that milkmaids who had had cowpox almost never got smallpox (which, it turns out, is caused by a close relative of the cowpox virus).

So Jenner decided to conduct a daring experiment. He collected pus from the sores of a milkmaid who had cowpox, and used this to inoculate a little boy named James Phipps. Later, when Phipps was reinoculated with pus from the sores of a person infected with smallpox, he did not contract the disease. In Latin, the word for cow is *vacca* – which explains where we get the word vaccine. History makes out the hero in this affair to be Edward Jenner, but I think the real hero that day was the young boy. Imagine having this big man approach you with a large needle and a tube full of pus! Although this isn't the sort of thing that could be done today, we can be thankful that Jenner's experiment was a success, because it paved the way for vaccinations that have saved countless lives.

Smallpox virus is not something humans encounter regularly. So Jenner's experiment showed that if the human immune system were given time to prepare, it could produce weapons which could provide protection against an invader which it never had seen before. Importantly, the smallpox vaccination only protected against smallpox or closely related viruses like cowpox. Phipps was still able to get mumps, measles, and the rest. This is one of the hallmarks of the adaptive immune system: It adapts to defend against specific invaders.

Antibodies and B cells

Eventually, immunologists determined that immunity to smallpox was conferred by special proteins that circulated in the blood of immunized individuals. These proteins were named antibodies, and the agent that caused the antibodies to be made was called an antigen – in this case, the cowpox virus. Here's a sketch that shows the prototype antibody, immunoglobulin G (IgG):



As you can see, an IgG antibody molecule is made up of two pairs of two different proteins, the heavy chain (Hc) and the light chain (Lc). Because of this structure, each molecule has two identical “hands” (Fab regions) that can bind to antigens. Proteins are the ideal molecules to use for constructing antibodies that can grasp attackers, because different proteins can fold up into a myriad of complex shapes.

IgG makes up about 75% of the antibodies in the blood, but there are four other classes of antibodies: IgA, IgD, IgE, and IgM. All of these antibodies are produced by “B cells” – white blood cells that are born in the bone marrow, and which then mature to become antibody factories called plasma B cells.

In addition to having hands that can bind to an antigen, an antibody molecule also has a constant region (Fc) “tail” which can bind to receptors (Fc receptors) on the surface of cells like macrophages. In fact, it is the special structure of the antibody Fc region that determines its class (e.g., IgG vs. IgA), which immune system cells it will bind to, and how it will function.

The hands of each antibody bind to a specific antigen (e.g., a protein on the surface of the smallpox virus), so in order to have antibodies available that can bind to many different antigens, many different antibody molecules are required. Now, if we want antibodies to protect us from every possible invader (and we do!), how many different antibodies would we need? Well, immunologists have


made rough estimates that about 100 million should do the trick. Since each antigen-binding region of an antibody is composed of a heavy chain and a light chain, we could mix and match about 10,000 different heavy chains and 10,000 different light chains to get the 100 million different antibodies we need. However, human cells only have about 25,000 genes in all, so if each heavy or light chain protein were encoded by a different gene, most of the B cell's genetic information would be used up just to make antibodies. You see the problem.

Generating antibody diversity by modular design

The riddle of how B cells could produce the 100 million different antibodies required to protect us was solved in 1977 by Susumu Tonegawa, who received the Nobel Prize for his discovery. When Tonegawa started working on this problem, the dogma was that the DNA in every cell in the body was the same. This made perfect sense, because after an egg is fertilized, the DNA in the egg is copied. These copies are then passed down to the daughter cells, where they are copied again, and passed down to their daughters, and so on. Therefore, barring errors in copying, each of our cells should end up with the same DNA as the original, fertilized egg. Tonegawa, however, hypothesized that although this is probably true in general, there might be exceptions. His idea was that all of our B cells might start out with the same DNA, but that as these cells mature, the DNA that makes up the antibody genes might change – and these changes might be enough to generate the 100 million different antibodies we need.

Tonegawa decided to test this hypothesis by comparing the DNA sequence of the light chain from a mature B cell with the DNA sequence of the light chain from an immature B cell. Sure enough, he found that they were different, and that they were different in a very interesting way. What Tonegawa and others discovered was that the mature antibody genes are made by modular design!

In every B cell, on the chromosomes that encode the antibody heavy chain, there are multiple copies of four types of DNA modules (gene segments) called V, D, J, and C. Each copy of a given module is slightly different from the other copies of that module. For example, in humans there are about forty different V segments, about twenty-five different D segments, six different J segments, and so on. To assemble a mature heavy chain gene, each B cell chooses (more or less at random) one of each kind of segment and pastes them together like this:



You have seen this kind of mix-and-match strategy used before to create diversity. For example, twenty different amino acids are mixed and matched to create the huge number of different proteins that our cells produce. And to create genetic diversity, the chromosomes you inherited from your mother and father are mixed and matched to make the set of chromosomes that goes into your egg or sperm cells. Once Mother Nature gets a good idea, she uses it over and over – and modular design is one of her very best ideas.

The DNA that encodes the light chain of the antibody molecule is also assembled by picking gene segments and pasting them together. Because there are so many different gene segments that can be mixed and matched, this scheme can be used to create about 10 million different antibodies – not quite enough. So, to make things even more diverse, when the gene segments are joined together, additional DNA bases are added or deleted. When this “junctional diversity” is included, there is no problem creating 100 million B cells, each with the ability to make a different antibody. The magic of this scheme is that by using modular design and junctional diversity, only a small amount of genetic information is required to create incredible antibody diversity.

Clonal selection

In the human bloodstream, there is a total of about three billion B cells. This seems like a lot, but if there are 100 million different kinds of B cells (to produce the 100 million different kinds of antibodies we need for protection), this means that, on average, there will only be about thirty B cells in the blood that can produce an antibody which will bind to a given antigen (e.g., a protein on the surface of a virus). Said another way, although we have B cells in our arsenal that can deal with essentially any invader, we don't have a lot of any one kind of B cell. As a result, when we are attacked, more of the appropriate B cells must be made.

6 LECTURE 1 An Overview

Indeed, B cells are made “on demand.” But how does the immune system know which B cells to make more of? The solution to this problem is one of the most elegant in all of immunology: the principle of clonal selection.

After B cells do their mix-and-match thing and paste together the modules required to form the complete recipes for their heavy and light chain antibody proteins, a relatively small number of these proteins is made – a “test batch” of antibody molecules, if you will. These tester antibodies, called B cell receptors (BCRs), are transported to the surface of the B cell and are tethered there with their hands facing out. Although each B cell has thousands of BCRs anchored on its surface, all the BCRs on a given B cell recognize the same antigen.

The B cell receptors on the surface of a B cell act like “bait,” and what they are “fishing for” is the molecule which their hands have the right shape to grasp – their “cognate” antigen. Sadly, the vast majority of B cells fish in vain. For example, most of us will never be infected with the SARS virus or the AIDS virus, so B cells that could make antibodies that recognize these viruses never find their match. It must be very frustrating for most B cells. They fish all their lives, and never catch anything!

On occasion, however, a B cell does make a catch. When a B cell’s receptors bind to its cognate antigen, that B cell is triggered to double in size and divide into two daughter cells – a process immunologists call proliferation. Both daughter cells then double in size and divide to produce a total of four cells, and so forth. Each cycle of cell growth and division takes about twelve hours to complete, and this period of proliferation usually lasts about a week. At the end of this time, a “clone” of roughly 20,000 identical B cells will have been produced, all of which have receptors on their surface that can recognize the same antigen. Now there are enough for a real defense!

After the selected B cells proliferate to form this large clone, most of them begin to make antibodies in earnest. The antibodies produced by these selected B cells are slightly different from the antibody molecules displayed on their surface in that there is no “anchor” to attach them to the B cell’s surface. As a result, these antibodies are transported out of the B cell and into the bloodstream. One B cell, working at full capacity, can pump out about 2,000 antibody molecules per second! After this heroic effort, most of these B cells die, having worked for only about a week as antibody factories.

When you think about it, this is an amazing strategy. First, because they employ modular design, B cells use


relatively few genes to create enough different antibody molecules to recognize any possible invader. Second, B cells are made on demand. So instead of filling up our bodies with a huge number of B cells which may never be used, we begin with a relatively small number of B cells, and then select the particular B cells that will be useful against the “invader *du jour*.” Once selected, the B cells proliferate rapidly to produce a large clone of B cells whose antibodies are guaranteed to be useful against the invader. Third, after the clone of B cells has grown sufficiently, most of these cells become antibody factories which manufacture huge quantities of the very antibodies that are right to defend against the invader. Finally, when the invader has been conquered, most of the B cells die. As a result, we don’t fill up with B cells that are appropriate to defend against yesterday’s invader, but which would be useless against the enemy that attacks us tomorrow. I love this system!

What antibodies do

Interestingly, although antibodies are very important in the defense against invaders, they really don’t kill anything. Their job is to plant the “kiss of death” on an invader – to tag it for destruction. If you go to a fancy wedding, you’ll usually pass through a receiving line before you are allowed to enjoy the champagne and cake. Of course, one of the functions of this receiving line is to introduce everyone to the bride and groom. But the other function is to be sure no outsiders are admitted to the celebration. As you pass through the line, you will be screened by someone who is familiar with all the invited guests. If she finds that you don’t belong there, she will call the bouncer and have you removed. She doesn’t do it herself – certainly not. Her role is to identify undesirables, not to show them the door. And it’s the same with antibodies: They identify invaders, and let other players do the dirty work.

In developed countries, the invaders we encounter most frequently are bacteria and viruses. Antibodies can bind to both types of invaders and tag them for destruction. Immunologists like to say that antibodies can “opsonize” these invaders. This term comes from a German word that means “to prepare for eating.” I like to equate *opsonize* with *decorate*, because I picture these bacteria and viruses with antibodies hanging all over them, decorating their surfaces. Anyway, when antibodies opsonize bacteria or viruses, they do so by binding to the invader with their Fab regions, leaving their Fc tails available to bind to Fc

receptors on the surface of cells such as macrophages. In this way, the antibodies form a bridge between the invader and the phagocyte (e.g., a macrophage), bringing the invader in close, and preparing it for eating (phagocytosis).



In fact, it's even better than this. When a phagocyte's Fc receptors bind to antibodies that are opsonizing an invader, the appetite of the phagocyte increases, making it even more phagocytic. Macrophages have proteins on their surface that can bind directly to many common invaders. However, the ability of antibodies to form a bridge between a macrophage and an invader allows a macrophage to "extend its grasp" to any invader to which an antibody can bind, common or uncommon. In effect, antibodies focus a macrophage's attention on invaders, some of which (the uncommon ones) a macrophage would otherwise ignore.

During a viral attack, antibodies can do something else that is very important. Viruses are parasites that enter our cells by binding to certain receptor molecules on a cell's surface. Of course, these receptors are not placed there for the convenience of the virus. They are normal receptors, like the Fc receptor, that have quite legitimate functions, but which the virus has learned to use to its own advantage. Once inside a cell, a virus uses the cell's machinery to make many copies of itself. These newly made viruses then burst out of the cell, sometimes killing it, and go on to infect neighboring cells. Now for the neat part: Antibodies can actually bind to a virus while it is still outside of a cell, and can keep the virus either from entering the cell or from replicating once it has entered. Antibodies with these properties are called "neutralizing" antibodies. For example, some neutralizing antibodies can prevent a virus from "docking" on the surface of the cell by binding to the part of the virus that would plug into the cellular receptor. When this happens, the virus is "hung out to dry," opsonized and ready to be eaten by phagocytes!

T cells

Although antibodies can tag viruses for phagocytic ingestion, and can help keep viruses from infecting cells, there is a flaw in the antibody defense against viruses: Once a virus gets into a cell, antibodies can't get to it, so the virus is safe to make thousands of copies of itself. Mother Nature recognized this problem, and to deal with it, she invented the famous "killer T cell," another member of the adaptive immune system team.

The importance of T cells is suggested by the fact that in a human, there are about a trillion of them. T cells are very similar to B cells in appearance. In fact, even under an ordinary microscope, an immunologist can't tell them apart. Like B cells, T cells are produced in the bone marrow, and on their surface they display antibody-like molecules called T cell receptors (TCRs). Like the B cell's receptors (the antibody molecules attached to its surface), TCRs are also made by a mix-and-match, modular design strategy. As a result, TCRs are about as diverse as BCRs. T cells also obey the principle of clonal selection: When a T cell's receptors bind to their cognate antigen, the T cell proliferates to build up a clone of T cells with the same specificity. This proliferation stage takes about a week to complete, so like the antibody response, the T cell response is slow and specific.


Although they are similar in many ways, there are also important differences between B cells and T cells. Whereas B cells mature in the bone marrow, T cells mature in the thymus (that's why they're called *T* cells). Further, although B cells make antibodies that can recognize any organic molecule, T cells specialize in recognizing protein antigens. In addition, a B cell can export (secrete) its receptors in the form of antibodies, but a T cell's receptors remain tightly glued to its surface. Perhaps most importantly, a B cell can recognize an antigen "by itself," whereas a T cell, like an old English gentleman, will only recognize an antigen if it is "properly presented" by another cell. I'll explain what this means in a bit.

There are actually three main types of T cells: killer T cells (frequently called cytotoxic lymphocytes – CTLs for short), helper T cells, and regulatory T cells. The killer T cell is a potent weapon that can destroy virus-infected cells. Indeed, by recognizing and killing virus-infected cells, the killer T cell solves the "hiding virus" problem – the flaw I mentioned in the antibody defense against viruses. The way the killer T cell does this is by making contact with its target cell and then triggering it to commit suicide! This "assisted suicide" is a great way to deal with viruses that

8 LECTURE 1 An Overview

have infected cells, because when a virus-infected cell dies, the viruses within the cell die also.

The second type of T cell is the helper T cell (Th cell). As you will see, this cell serves as the quarterback of the immune system team. It directs the action by secreting protein molecules called cytokines that have dramatic effects on other immune system cells. These cytokines have names like interleukin 2 (IL-2) and interferon gamma (IFN- γ), and we will discuss what they do in later lectures. For now, it is only important to realize that helper T cells are basically cytokine factories.



The third type of T cell, the regulatory T cell, is still quite mysterious. Although it is believed that regulatory T cells help keep other T cells under control, it still isn't clear exactly how they do this or what "under control" means in this context.

Antigen presentation


One thing I need to clear up is exactly how antigen is presented to T cells. It turns out that special proteins called major histocompatibility complex proteins (MHC for short) actually do the "presenting," and that T cells use their receptors to "view" this presented antigen. As you may know, *histo-* means "tissue," and these major histocompatibility proteins, in addition to being presentation molecules, also are involved in the rejection of transplanted organs. In fact, when you hear that someone is waiting for a "matched" kidney, it's the MHC molecules of the donor and the recipient that the transplant surgeon is trying to match.

There are two types of MHC molecules, called class I and class II. Class I MHC molecules are found in varying

amounts on the surface of most cells in the body, and they function as "billboards" that inform killer T cells about what is going on inside other cells. For example, when a human cell is infected by a virus, fragments of viral proteins (called peptides) are loaded onto class I MHC molecules, and transported to the surface of the infected cell. By inspecting these protein fragments displayed by class I MHC molecules, killer T cells can use their receptors to "look into" the cell to determine that it has been infected and that it should be destroyed.

Class II MHC molecules also function as billboards, but this display is intended for the enlightenment of helper T cells. Only certain cells in the body make class II MHC molecules, and these are called antigen presenting cells (APCs for short). Macrophages, for example, are excellent antigen presenting cells. During a bacterial infection, a macrophage will "eat" the bacteria, and will load fragments of ingested bacterial proteins onto class II MHC molecules for display on the surface of the macrophage. By using their T cell receptors, helper T cells can scan the macrophage's class II MHC billboards for news of the bacterial infection. So class I MHC molecules alert killer T cells when something isn't right inside a cell, and class II MHC molecules displayed on APCs inform helper T cells that problems exist outside of cells.

Although a class I MHC molecule is made up of one long chain (the heavy chain) plus a short chain (β 2-microglobulin), and a class II MHC molecule has two long chains (α and β), you'll notice that these molecules look rather similar.




I know it's hard to visualize the real shapes of molecules from drawings like this, so I thought I'd show you a few pictures that may make this more real. Here's what an


empty MHC molecule might look like from the viewpoint of the T cell receptors. Right away you see the groove into which the protein fragment would fit.



Next, let's look at a fully loaded, class I molecule:



I can tell it's a class I MHC molecule because the peptide is contained nicely within the groove. It turns out that the ends of the groove of a class I molecule are closed, so a protein fragment must be about nine amino acids in length to fit in properly. Class II MHC molecules are slightly different:




Here you see that the peptide overflows the groove. This works fine for class II, because the ends of the groove are open, so protein fragments as large as about twenty amino acids fit nicely.

So MHC molecules resemble buns, and the protein fragments they present resemble wieners. And if you imagine that the cells in our bodies have hot dogs on their surfaces, you won't be far wrong about antigen presentation. That's certainly the way I picture it!

Activation of the adaptive immune system

Because B and T cells are such potent weapons, Mother Nature put into place the requirement that cells of the adaptive immune system must be activated before they can function. Collectively, B and T cells are called lymphocytes, and how they are activated is one of the key issues in immunology. To introduce this concept, I will sketch how helper T cells are activated.

The first step in the activation of a helper T cell is recognition of its cognate antigen (e.g., a fragment of a bacterial protein) displayed by class II MHC molecules on the surface of an antigen presenting cell. But seeing its cognate antigen on that billboard isn't enough – a second signal or "key" is also required for activation. This second signal is nonspecific (it's the same for any antigen), and it involves a protein (B7 in this drawing) on the surface of an APC that plugs into its receptor (CD28 in this drawing) on the surface of the helper T cell.



You see an example of this kind of two-key system when you visit your safe deposit box. You bring with you a key that is specific for your box – it won't fit any other. The bank teller provides a second, nonspecific key that will fit all the boxes. Only when both keys are inserted into the locks on your box can it be opened. Your specific key alone won't do it, and the teller's nonspecific key alone won't either. You need both. Now, why do you suppose helper T cells and other cells of the adaptive immune system require two keys for activation? For safety, of course – just like your bank box. These cells are powerful weapons that must only be activated at the appropriate time and place.

Once a helper T cell has been activated by this two-key system, it proliferates to build up a clone composed

10 LECTURE 1 An Overview


of many helper T cells that recognize the same antigen. These helper cells then mature into cells that can produce the cytokines needed to direct the activities of the immune system. B cells and killer T cells also require a two-key system for their activation.

The secondary lymphoid organs

If you've been thinking about how the adaptive immune system might get turned on during an attack, you've probably begun to wonder whether this could ever happen. After all, there are probably only about 10,000 T cells that will have TCRs specific for a given invader, and for these T cells to be activated, they must come in contact with an antigen presenting cell that also has "seen" the invader. Given that these T cells and APCs are spread all over the body, it would seem very unlikely that this would happen before an invasion got completely out of hand. Fortunately, to make this system work with reasonable probability, Mother Nature invented the "secondary lymphoid organs," the best known of which is the lymph node. You may not be familiar with the lymphatic system, so I'd better say a few words about it.

In your home, you have two plumbing systems. The first supplies the water that comes out of your faucets. This is a pressurized system, with the pressure being provided by a pump. In addition, you also have another plumbing system that includes the drains in your sinks, showers, and toilets. This second system is not under pressure – the water just flows down the drain and out into the sewer. The two systems are connected in the sense that eventually the wastewater is recycled and used again.

The plumbing in a human is very much like this. We have a pressurized system (the cardiovascular system) in which blood is pumped around the body by the heart. Everybody knows about this one. But we also have another plumbing system: the lymphatic system. This system is not under pressure, and it drains the fluid (called lymph) that leaks out of our blood vessels into our tissues. Without this system, our tissues would fill up with fluid and we'd look like the Pillsbury Doughboy. Fortunately, lymph is collected from our tissues into lymphatic vessels, and is transported by these vessels, under the influence of muscular contraction, through a series of one-way valves to the upper torso, where it is recycled back into the blood. From this diagram, you can see that as the lymph winds its way back to empty into the blood, it passes through a series of way stations – the lymph nodes.



There are thousands of lymph nodes that range in size from very small to almost as big as a brussels sprout. Invaders like bacteria and viruses are carried by the lymph to nearby nodes, and antigen presenting cells that have picked up foreign antigens in the tissues travel to lymph nodes to present their cargo. Meanwhile, B cells and T cells circulate from node to node, looking for the antigens for which they are "fated." So lymph nodes really function as "dating bars" – places where T cells, B cells, APCs, and antigen all gather for the purposes of communication and activation. By bringing these cells and antigens together within the small volume of a lymph node, Mother Nature increases the probability that they will interact to efficiently activate the adaptive immune system.

Immunological memory

After B and T cells have been activated, have proliferated to build up clones of cells with identical antigen specificities, and have vanquished the enemy, most of them die off. This is a good idea, because we wouldn't want our immune systems to fill up with old B and T cells. On the other hand, it would be nice if some of these experienced B and T cells would stick around, just in case we are exposed to the same invaders again. That way, the adaptive immune system wouldn't have to start from scratch. And that's just

the way it works. These “leftover” B and T cells are called memory cells, and in addition to being more numerous than the original, inexperienced B and T cells, memory cells are easier to activate. As a result of this immunological memory, the adaptive system usually can spring into action so quickly during a second attack that you never even experience any symptoms.

Tolerance to self

As I mentioned earlier, B cell receptors and T cell receptors are so diverse that they should be able to recognize any potential invader. This raises a problem, however, because if the receptors are this diverse, many of them are certain to recognize our own “self” molecules (e.g., the molecules that make up our cells, or proteins like insulin that circulate in our blood). If this were to happen, our adaptive immune system might attack our own bodies, and we could die from autoimmune disease. Fortunately, Mother Nature has devised ways to educate B cells and T cells to discriminate between ourselves and dangerous invaders. Although the mechanisms responsible for self-tolerance still are not completely understood, the education which B and T cells receive in self-tolerance is sufficiently rigorous that autoimmune disease is relatively rare.

A COMPARISON OF THE INNATE AND ADAPTIVE IMMUNE SYSTEMS

Now that you have met some of the main players, I want to emphasize the differences between the innate and adaptive immune system “teams.” Understanding how they differ is crucial to understanding how the immune system works.

Imagine that you are in the middle of town and someone steals your shoes. You look around for a store where you can buy another pair, and the first store you see is called Charlie’s Custom Shoes. This store has shoes of every style, color, and size, and the salesperson is able to fit you in exactly the shoes you need. However, when it comes time to pay, you are told that you must wait a week or two to get your shoes – they will have to be custom-made for you, and that will take a while. But you need shoes right now! You are barefoot, and you at least need something to put on your feet until those custom shoes arrive. So they send you across the street to Freddie’s Fast Fit – a store that has only a few styles and sizes. Freddie’s wouldn’t be able to fit Shaquille O’Neal, but this store does stock shoes in the

common sizes that fit most people, so you can get a pair right away to tide you over.

This is very similar to the way the innate and adaptive immune systems work. The players of the innate system (like the macrophage) are already in place, and are ready to defend against relatively small quantities of the invaders we are likely to meet on a day-to-day basis. In many instances, the innate system is so effective and so fast that the adaptive immune system never even kicks in. In other cases, the innate system may be insufficient to deal with the invasion, and the adaptive system will need to be mobilized. This takes a while, because although the B and T cells of the adaptive system can deal with huge quantities of almost any invader, these weapons must be custom-made. Meanwhile, the innate immune system must do its best to hold the invaders at bay.

THE INNATE SYSTEM RULES!

Until fairly recently, immunologists thought that the only function of the innate system was to provide a rapid defense which would deal with invaders while the adaptive immune system was getting cranked up. However, it is now clear that the innate system does much more than that.

The adaptive immune system’s antigen receptors (BCRs and TCRs) are so diverse that they can probably recognize any protein molecule in the universe. However, the adaptive system is clueless as to which of these molecules is dangerous and which is not. So how does the adaptive system distinguish friend from foe? The answer is that it relies on the judgment of the innate system.

In contrast to the antigen receptors of the adaptive immune system, which are totally “unfocused,” the receptors of the innate system are precisely tuned to detect the presence of the common pathogens (disease-causing agents) we encounter in daily life – viruses, bacteria, fungi, and parasites. In addition, the innate system has receptors that can detect when “uncommon” pathogens kill human cells. Consequently, it is the innate system which is responsible for sensing danger and for activating the adaptive immune system.

So in a real sense, the innate system gives “permission” to the adaptive system to respond to an invasion. But it’s even better than that, because the innate system does more than just turn the adaptive system on. The innate system actually integrates all the information it collects about an invader, and formulates a plan of action. This “game plan,”

12 LECTURE 1 [An Overview](#)

which the innate system delivers to the adaptive immune system, tells which weapons to mobilize (e.g., B cells or killer T cells) and exactly where in the body these weapons should be deployed. So if we think of the helper T cell as the quarterback of the adaptive immune system team, we should consider the innate immune system to be the “coach” – for it is the innate system which “scouts” the opponents, designs the game plan, and sends in the plays for the quarterback to call.

EPILOGUE

We have come to the end of our turbo overview of the immune system, and by now you should have a rough idea of how the system works. In the next eight lectures, we will focus more sharply on the individual players of the innate and adaptive system teams, paying special attention to how and where these players interact with each other to make the system function efficiently.