

Part 1

Introduction

COPYRIGHTED MATERIAL

Chapter 1

Tectonics of sedimentary basins, with revised nomenclature

RAYMOND V. INGERSOLL

Department of Earth and Space Sciences, University of California, Los Angeles, California

ABSTRACT

Actualistic plate-tectonic models are the best framework within which to understand the tectonics of sedimentary basins. Sedimentary basins develop in divergent, intra-plate, convergent, transform, hybrid, and miscellaneous settings. Within each setting are several variants, dependent on type(s) of underlying crust, structural position, sediment supply, and inheritance. Subsidence of sedimentary basins results from (1) thinning of crust (2) thickening of mantle lithosphere (3) sedimentary and volcanic loading (4) tectonic loading (5) subcrustal loading (6) asthenospheric flow, and (7) crustal densification. Basins vary greatly in size, life span, and preservation potential, with short-lived basins formed in active tectonic settings, especially on oceanic crust, having low preservation potential, and long-lived basins formed in intraplate settings having the highest preservation potential.

Continental rifts may evolve into nascent ocean basins, which commonly evolve into active ocean basins bordered by intraplate continental margins with three types of configurations: shelf-slope-rise, transform, and embankment. Continental rifts that do not evolve into oceans become fossil rifts, which later become sites for development of intracratonic basins and aulacogens. If all plate boundaries within and around an ocean basin become inactive, a dormant ocean basin develops, underlain by oceanic crust and surrounded by continental crust.

Sites for sedimentary basins in convergent settings include trenches, trench slopes, forearcs, intra-arcs, backarcs, and retroarcs. Complex dynamic behavior of arc-trench systems results in diverse configurations for arc-related basins. Most notable is the overall stress regime of the arc-trench system, with resulting response along and behind the magmatic arc. Intra-arc rifting in highly extensional arcs commonly evolves into backarc spreading to form new oceanic crust. Backarcs of neutral arcs can contain any type of preexisting crust that was trapped there at the time of initiation of the related subduction zone. Highly compressional arcs develop retroarc foldthrust belts and related retroforeland basins, and may develop hinterland basins; in extreme cases, broken retroforelands may develop in former cratonic areas.

As nonsubductable continental or arc crust is carried toward a subduction zone, collision generally initiates at one point and the resulting suture propagates away from this point of initial impact. Remnant ocean basins form on both sides of the initial impact point, and rapidly fill with sediment derived from the suture zone. As collision continues, the flux of sediment into the remnant ocean basin(s) increases concurrently with shrinkage of the basin until final suturing and obduction of the accreted sediment occur. Concurrently with collision, proforeland basins form on continental crust of the subducting plate and collisional retroforeland basins form on the overriding plate. Impactogens, broken forelands, and hinterland basins also may result.

In transform settings and along complex strike-slip fault systems related to convergent settings, changing stress regimes related to irregularities in fault trends, rock types, and plate motions result in transtension, transpression, and transrotation, with associated complex, diverse, and short-lived sedimentary basins.

Two previously unnamed basin types that have received increasing attention recently are halokinetic basins (related to salt tectonics, especially along intraplate margins with embankment configurations) and bolide basins (resulting from extraterrestrial impacts). Sediment accumulates in successor basins following cessation of basin-controlling processes, whether in divergent, convergent, transform, or hybrid settings.

The ultimate goal of classifying and reviewing all types of sedimentary basins is the improvement of paleotectonic and paleogeographic reconstructions through the application of actualistic models for basin evolution. Interdisciplinary studies that test and refine these models will improve our knowledge of Earth history.

Keywords: basin nomenclature; plate-tectonic settings; subsidence mechanisms; preservation potential; paleotectonic reconstruction

INTRODUCTION

It has been more than a decade since I reviewed and revised my original basin classification (i.e., Ingersoll, 1988; Ingersoll and Busby, 1995), which was based primarily on Dickinson's (1974b, 1976a) statement of fundamental principles that should guide discussion of the tectonics of sedimentary basins. Many new insights and models have been developed recently; in addition, nomenclature has evolved in complex ways. Therefore, now is an appropriate time to consolidate, revise, and discuss how to communicate about the tectonics of sedimentary basins.

As in my previous papers on this subject, I follow Dickinson's (1974b, 1976a) suggestions that nomenclature and classification be based on the following actualistic plate-tectonic processes and characteristics, which ultimately control the location, initiation, and evolution of sedimentary basins in diverse tectonic settings. Horizontal motions of plates, thermal changes through time, stretching and shortening of crust, isostatic adjustments, mantle dynamics, surficial processes, and even extraterrestrial events influence sedimentary basins. Additional study of sedimentary basins, inevitably, leads to greater complexity of models to explain them. Although we should search for unifying principles that lead to deeper understanding of processes and results, the complexity of the real world dictates that enhanced knowledge about sedimentary basins results in more complex models. Thus, new types of sedimentary basins are added to the list provided in Ingersoll and Busby (1995) because these are actual features that need to be understood. Gould (1989, 98) stated, "Classifications are theories about the basis of natural order, not dull catalogues compiled

only to avoid chaos." I hope that my discussion serves the dual purposes of reducing nomenclatural chaos and suggesting a framework within which to understand the complex controls on the origin and evolution of sedimentary basins.

NOMENCLATURE

First-order criteria for classifying sedimentary basins (Dickinson, 1974b, 1976a) are (1) type of nearest plate boundary(ies) (2) proximity of plate boundary(ies), and (3) type of substratum. Thus, the first-order classification, based on criteria (1) and (2) is divergent, intraplate, convergent, transform, hybrid, and miscellaneous settings (Table 1.1). Within each of these categories are several variants, dependent on type of substratum (oceanic, transitional, continental, and anomalous crust), as well as structural position, sediment supply, and inheritance.

Basin classification and nomenclature are based on characteristics of a basin *at the time of sedimentation*. Thus, many stratigraphic successions are multidimensional and multigenerational in terms of plate-tectonic controls on their evolution. A single stratigraphic succession may represent several different tectonic settings. "The evolution of a sedimentary basin thus can be viewed as the result of a succession of discrete plate-tectonic settings and plate interactions whose effects blend into a continuum of development" (Dickinson, 1974b, 1).

It is important to realize that "basin," as used herein, refers to any stratigraphic accumulation of sedimentary or volcanic rock; the three-dimensional architecture of basins may approximate saucers, wedges, sheets, and odd shapes.

Table 1.1. Basin classification with modern and ancient examples

Setting	Basin type	Definition	Modern example	Ancient example	Analog model (Figure)
Divergent	Continental rifts	Rifts within continental crust, commonly associated with bimodal magmatism	Rio Grande rift	Proterozoic Keweenawan rift	3B
	Nascent ocean basins and continental margins	Incipient ocean basins floored by new oceanic crust and flanked by young rifted continental margins	Red Sea	Jurassic of East Greenland	3C
Intraplate	Intraplate continental margins				
	<i>Shelf-slope-rise configuration</i>	Mature rifted intraplate continental margins with shelf edge near boundary between continental and oceanic crust	East Coast of USA	Early Paleozoic of USA and Canadian Cordillera	3D
	<i>Transform configuration</i>	Intraplate continental margins that originate along transform plate boundaries	South Coast of West Africa	Precambrian-early Paleozoic Alabama-Oklahoma transform	3E
	<i>Embankment configuration</i>	Progradational intraplate continental margins with shelf edge above oceanic crust	Mississippi River Gulf Coast	Early Paleozoic Meguma terrane of Canadian Appalachians (?)	3F
	Intracratonic basins	Broad cratonic basins underlain by fossil rifts	Chad basin	Paleozoic Michigan basin	3A
	Continental platforms	Stable cratons with thin and laterally extensive sedimentary strata	Barents Sea	Middle Paleozoic, North American	3A
	Active ocean basins	Basins floored by oceanic crust formed at active divergent plate boundaries unrelated to arc-trench systems	Pacific Ocean	midcontinent Various ophiolitic complexes (?)	3G
	Oceanic islands, seamounts, aseismic ridges, and plateaus	Sedimentary aprons and platforms formed in intraoceanic settings other than arc-trench systems	Emperor-Hawaii seamounts	Mesozoic Snow Mountain Volcanic Complex (Franciscan of northern California)	3G
	Dormant ocean basins	Basins floored by oceanic crust, which is neither spreading nor subducting	Gulf of Mexico	Paleozoic Tarim basin (China) (?)	3H
					(continued)

Table 1.1. (*Continued*)

Setting	Basin type	Definition	Modern example	Ancient example	Analog model (Figure)
Convergent	Trenches	Deep troughs formed at oceanic subduction zones	Chile Trench	Cretaceous, Shumagin Island (southern Alaska)	4A
	Trench-slope basins	Local structural depressions on subduction complexes	Central American Trench	Cretaceous Cambria slab (central California)	4B
	Forearc basins	Basins within arc-trench gaps	Offshore Sumatra	Cretaceous Great Valley (California)	4B
	Intraarc basins <i>Oceanic intraarc basins</i>	Basins along intraoceanic arc platforms, which include superposed and overlapping volcanoes	Izu Bonin arc	Copper Hill, Gopher Ridge Complex (Jurassic, California)	4A
	<i>Continental intraarc basins</i>	Basins along continental-margin arc platforms, which include superposed and overlapping volcanoes	Lago de Nicaragua	Early Jurassic Sierra Nevada (eastern California)	4C
	Backarc basins <i>Oceanic backarc basins</i>	Oceanic basins behind intraoceanic magmatic arcs (including interarc basins between active and remnant arcs)	Marianas backarc	Jurassic Josephine ophiolite (northern California)	4A,B
	<i>Continental backarc basins</i>	Continental basins behind continental-margin arcs without foreland foldthrust belts	Sunda Shelf	Late Triassic - Early Jurassic of USA Cordillera	4C
	Retroforeland basins <i>Retroarc foreland basins</i>	Foreland basins on continental sides of continental-margin arc-trench systems	Andes foothills	Cretaceous Sevier foreland of USA Cordillera	4E
	<i>Collisional retroforeland basins</i>	Foreland basins formed on overriding plates during continental collisions (may have retroarc precursors)	Western Tarim basin (China)	Triassic-Jurassic Ordos basin (China)	4F
	<i>Broken-retroforeland basins</i>	Basins formed among basement-cored uplifts in retroforeland setting	Sierras Pampeanas basins (Argentina)	Late Cretaceous-Paleogene Laramide basins of USA Cordillera	4D
	Remnant ocean basins	Shrinking ocean basins between colliding continental margins and/or arc-trench systems, and ultimately subducted or deformed within suture belts	Bay of Bengal	Pennsylvanian-Permian Ouachita basin	4E

Proforeland basins	Foreland basins formed on continental crust that is part of the subducting plate during continental and/or arc collision	Persian Gulf	Mid-Genozoic Swiss Molasse basin	4F
Wedge-top basins	Basins formed and carried on moving thrust sheets	Peshawar basin (Pakistan)	Neogene, Apennines (Italy)	4F
Hinterland basins	Basins formed on thickened continental crust behind foreland fold-thrust belts	Altiplano Plateau (Bolivia)	Neogene Zhada basin (Tibet)	4D
Transform	Transensional basins	Dead Sea	Carboniferous Magdalen basin (Gulf of Saint Lawrence)	5A
	Transpressional basins	Santa Barbara Basin (foreland type) (California)	Miocene Ridge basin (fault-bend type) (California)	5B
	Transrotational basins	Western Aleutian forearc (?)	Miocene Los Angeles basin (Topanga basin) (California)	5C
Miscellaneous and hybrid	Aulacogens	Mississippi embayment	Paleozoic Anadarko aulacogen (Oklahoma)	6A
	Impactogens	Baikal rift (distal) (Siberia)	Rhine Graben (proximal) (Europe)	6B
	Collisional broken foreland	Qaidam basin (China)	Pennsylvanian-Permian Ancestral Rocky Mountain basins of the USA Cordillera	6B
	Halokinetic basins	Mini-basins of deep Gulf of Mexico	Cretaceous-Paleogene La Popa basin (Mexico)	3F
	Bolide basins	Meteor Crater (Arizona)	Cretaceous-Paleogene Chicxulub basin (Mexico)	3E
	Successor basins	Southern Basin and Range (Arizona)	Paleogene Sustut basin (?) (British Columbia)	5C

Modified after Ingersoll and Busby (1995).

Also, basins may form by subsidence of a substrate, development of a barrier to transport of sediment, filling of a preexisting hole, or relative movement of source and sink.

**SUBSIDENCE MECHANISMS
AND PRESERVATION POTENTIAL**

Surfaces of deposition may subside due to the following processes (Dickinson, 1974b, 1976a, 1993; Ingersoll and Busby, 1995) (Table 1.2): (1) thinning of crust due to stretching, erosion, and magmatic withdrawal (2) thickening of mantle lithosphere during cooling (3) sedimentary and volcanic loading (local crustal isostasy or regional lithospheric flexure) (4) tectonic loading of both crust and lithosphere (5) subcrustal loading of both crust and lithosphere (6) dynamic effects of asthenospheric flow, and (7) crustal densification. Figure 1.1 illustrates that crustal thinning dominates during early stages of extension (e.g., rifts and transtensional basins), and mantle-lithospheric thickening dominates following the initiation of seafloor spreading (during the rift-to-drift transition along divergent margins which evolve into intraplate margins). Sedimentary loading dominates along continental-oceanic crustal boundaries which are supplied by major rivers and deltas (e.g., continental embankments and remnant ocean basins). Tectonic loading dominates in settings where crustal shortening dominates (e.g., trenches and foreland basins). The other three types of subsidence mechanisms are generally subordinate.

The diversity of tectonic and structural settings of sedimentary basins dictates that they vary greatly in size, life span, and preservation potential (Fig. 1.2) (Ingersoll, 1988; Ingersoll and Busby, 1995; Woodcock, 2004). Many sediment

accumulations are destined to be destroyed relatively soon after deposition (e.g., most basins residing on oceanic crust or in rapidly uplifting orogenic settings). In contrast, basins formed during and following stretching of continental crust (e.g., continental rifts that either evolve into seafloor spreading or fail to do so) have high preservation potential because they subside and are buried beneath intraplate deposits following rifting. On the other hand, stratigraphic sequences along intraplate continental margins are destined to be partially subducted as they are pulled into trenches, thus preserving them at moderate to deep crustal levels as highly deformed and metamorphosed terranes. Such metasedimentary and metavolcanic terranes, along with voluminous sediments deposited in remnant ocean basins, are major rock bodies involved in the construction of continental crust, although their substrates (oceanic crust) are mostly subducted (e.g., Graham et al., 1975; Ingersoll et al., 1995, 2003).

DIVERGENT SETTINGS

Sequential rift development and continental separation

The relative importance of “active” (mantle-convective-driven) versus “passive” (lithospheric-driven) processes during initiation of continental rifting is debated (e.g., Sengor and Burke, 1978; Ingersoll and Busby, 1995; Sengor, 1995). Regardless of the mechanisms of initiation of rifting, continental rifts may experience two life paths: “successful” rifting that evolves into seafloor spreading to form nascent ocean basins (Ingersoll and Busby, 1995; Leeder, 1995), which then evolve into active ocean basins with paired intraplate margins (Fig. 1.3), or “failed” rifting, which does not evolve into nascent ocean basins,

Table 1.2. Subsidence mechanisms

Crustal thinning	Extensional stretching, erosion during uplift, and magmatic withdrawal
Mantle-lithospheric thickening	Conversion of asthenosphere to mantle lithosphere during cooling following cessation of stretching and/or heating
Sedimentary and volcanic loading	Local isostatic compensation of crust and/or regional lithospheric flexure during sedimentation and volcanism
Tectonic loading	Local isostatic compensation of crust and/or regional lithospheric flexure during overthrusting and/or underpulling
Subcrustal loading	Lithospheric flexure during underplating of dense lithosphere
Asthenospheric flow	Dynamic effects of asthenospheric flow, commonly due to descent or delamination of subducted lithosphere
Crustal densification	Increased density of crust due to changing pressure/temperature conditions and/or emplacement of higher-density melts into lower-density crust




Fig. 1.1. Suggested subsidence mechanisms for all types of sedimentary basins.

instead producing fossil rifts, commonly overlain by intracratonic basins (Sengor, 1995). Ingersoll and Busby (1995), Leeder (1995), and Sengor (1995) reviewed most aspects of continental stretching, basin formation, structural development, and

different life paths during and after continental rifting. Here, I highlight changes in nomenclature and models involved in the evolution from continental rifts to intraplate margins (the rift-drift transition).




Fig. 1.2. Typical life spans for sedimentary basins versus their post-sedimentation preservation potential, which refers to average time interval during which basins will not be uplifted and eroded during and following sedimentation. Sedimentary or volcanic fill may be preserved as accretionary complexes during and after basin destruction (especially true for all strata deposited on oceanic crust). Intraplate continental margins have high preservation potential in the sense of retaining their basement, but are likely to be highly deformed and metamorphosed beneath and within suture belts, and may be difficult to recognize in ancient settings.




Fig. 1.3. True-scale actualistic analog models for sedimentary basins in divergent, intraplate and miscellaneous settings. Mantle lithosphere thins during decompression melting as plates diverge; mantle lithosphere thickens during cooling, following cessation of divergence. Also shown are two miscellaneous basins (bolide and halokinetic). Placement of bolide basin is arbitrary; they may form anywhere on Earth's surface, although preservation is more likely in cratonic areas (as shown in E). Halokinetic basins may form anywhere that salt is deeply buried; however, continental embankments (as shown in F) are the most common locations. Continental crust = jackstraw pattern; oceanic crust = vertical lines; mantle lithosphere and derived igneous rocks = black; asthenosphere and derived melts = orange; salt (halokinetic only) = black.

Continental rifts

The most common basins associated with continental rifts (Fig. 1.3b) (“terrestrial rift valleys” of Dickinson, 1974b; Ingersoll, 1988) are half grabens developed on the hanging walls of normal faults (Leeder and Gawthorpe, 1987; Leeder, 1995; Gawthorpe and Leeder, 2000). Gawthorpe and Leeder (2000) summarized conceptual models for the tectono-sedimentary evolution of continental rift basins, including their three-dimensional development. They discussed structural, geomorphic, climatic, and lake/sea-level influences on basin development.

All of the models presented by Gawthorpe and Leeder (2000) involve high-angle normal faults. In these half grabens, most sediment is derived from the hanging wall, whereas the coarsest material, which is derived primarily from the footwall, is restricted to small steep alluvial fans or fan deltas along the faulted basin boundary. In contrast, supradetachment basins (formed above low-angle normal faults) receive most of their sediment from the breakaway footwall and tend to be dominated by coarse-grained detritus (Friedmann and Burbank, 1995). Additional variants on the Gawthorpe and Leeder (2000) half-graben model include development of accommodation zones, relay ramps, anticlinal-full-graben basins, and synclinal-horst basins (Rosendahl, 1987; Faulds and Varga, 1998; Ingersoll, 2001; Mack et al., 2003).

Nascent ocean basins and continental margins

As continental lithosphere is stretched and thinned, mantle asthenosphere eventually rises close to the surface (Fig. 1.3c). During the transition from continental rifting to seafloor spreading, transitional crust forms, either as stretched continental crust (quasicontinental) or sediment-rich basaltic crust (quasioceanic) (Dickinson, 1974b; Ingersoll, 2008b). Continental rifting evolves into seafloor spreading only in the absence of significant sediment so that oceanic crust is the only solid material with which rising asthenospheric melts can interact (Einsele, 1985; Nicolas, 1985). Thus, a significant width of transitional crust typically forms on the margins of nascent ocean basins prior to initiation of true seafloor spreading.

As these transitional types of crust form and the two continental margins move apart, a nascent ocean basin develops (“proto-oceanic gulf” and “narrow ocean” of Dickinson (1974b);

“proto-oceanic rift trough” of Ingersoll (1988)). The Red Sea is the type nascent ocean basin, with active seafloor spreading, clastic and carbonate sedimentation along the margins, and uplifted rift shoulders along the continental margins (Cochran, 1983; Bohannon, 1986a, 1986b; Coleman, 1993; Leeder, 1995; Purser and Bosence, 1998; Bosworth et al., 2005). Thick evaporite deposits may form during the transition from rift basin to nascent ocean basin, as well as during much of the history of nascent ocean basins, given the right combination of arid climate, limited communication with other marine bodies, and lack of detrital input (Dickinson, 1974b). The Gulf of California is an example of a transtensional nascent ocean basin (e.g., Atwater, 1989; Lonsdale, 1991; Atwater and Stock, 1998; Axen and Fletcher, 1998).

INTRAPLATE SETTINGS

Intraplate continental margins

Nascent ocean basins evolve into wide (Atlantic-type) oceans as two continents diverge along spreading ridges. During this evolutionary process, the newly rifted continental margins with uplifted rift flanks cool and subside as they move away from the spreading ridge. This process is referred to as the “rift-to-drift” transition, as a divergent setting evolves into an intraplate setting (Dickinson, 1974b, 1976a; Ingersoll, 1988; Bond et al., 1995; Ingersoll and Busby, 1995). Withjack et al. (1998) discussed complications in timing and process during this transition.

Subsidence mechanisms evolve from (1) thinning of continental crust by stretching and erosion during doming and rifting, to (2) thermal subsidence following rifting as the intraplate margin moves away from the spreading ridge, to (3) both local crustal and regional lithospheric sediment loading during the later history of the intraplate continental margin (Bond et al., 1995; Ingersoll and Busby, 1995). Lower-crustal and subcrustal flow and densification can locally modify subsidence.

Shelf-slope-rise configuration

Most mature intraplate continental margins consist of a seaward thickening wedge of shelf deposits on top of continental crust, which is thinner seaward (Fig. 1.3d). Transitional crust (both quasicontinental and quasioceanic; Dickinson, 1974b,

1976a) underlies the seaward transition from thick shelf deposits to thin slope deposits, which, in turn, merge into thick turbiditic rise and abyssal-plain deposits on oceanic crust (Bond et al., 1995; Ingersoll and Busby, 1995). Most modern Atlantic continental margins have this configuration, with carbonate environments dominating at lower latitudes devoid of extensive clastic input.

Transform configuration

Intraplate continental margins that originate along transform boundaries rather than rift boundaries have narrower sediment prisms and transitional crust (Fig. 1.3e). Tens of millions of years may pass between the time of initiation of transform motion (coincident with the rift-to-drift transition on adjoining margins) and the time of intraplate sedimentation (following passage of the spreading ridge along the transform boundary) (e.g., Bond et al., 1995; Turner et al., 2003; Wilson et al., 2003). The southern coast of West Africa exemplifies these characteristics; the latest Proterozoic - early Paleozoic Alabama-Oklahoma transform margin is an ancient example (e.g., Thomas, 1991).

Embankment configuration

Major rivers along intraplate continental margins commonly are localized by fossil rifts trending at high angle to the margins (Burke and Dewey, 1973; Dickinson, 1974b; Audley-Charles et al., 1977; Ingersoll and Busby, 1995). The best examples are the Niger Delta (Burke, 1972) and the Mississippi Delta (Worrall and Snelson, 1989; Salvador, 1991; Galloway et al., 2000), where the shelf edge has prograded over oceanic crust because the maximum sediment thickness allowed by isostatic loading (16–18 km; Kinsman, 1975) has been reached inland of the shelf edge (Fig. 1.3f). In the case of the USA Gulf Coast, several rivers in addition to the Mississippi have contributed to considerable progradation of the continental margin over a wide area; this is the type example of a continental embankment, a distinctly different configuration than either the shelf-slope-rise or transform configuration.

Intracratonic basins

Most intracratonic basins (e.g., Michigan basin) overlie fossil rifts (e.g., DeRito et al., 1983; Quinlan, 1987; Klein, 1995; Sengor, 1995; Howell

and van der Pluijm, 1999) (Fig. 1.3a). Renewed periods of subsidence in cratonic basins can generally be correlated with changes in lithospheric stress related to orogenic activity in neighboring orogenic belts (DeRito et al., 1983; Howell and van der Pluijm, 1999). Subsidence occurs when lithospheric rigidity lessens, allowing uncompensated mass in the upper crust (remnants of fossil rifts) to subside over a broad area. Between times of orogenic activity, the lithosphere strengthens so that attainment of local isostatic equilibrium is interrupted. Thus, an intracratonic basin may take hundreds of millions of years to reach full isostatic compensation (DeRito et al., 1983; Ingersoll and Busby, 1995; Howell and van der Pluijm, 1999).

Continental platforms

Cratonic stratigraphic sequences primarily reflect global tectonic events and eustasy (e.g., Sloss, 1988; Bally, 1989), although mantle dynamics, and local and regional events also influence continental platforms (e.g., Cloetingh, 1988; Burgess and Gurnis, 1995; van der Pluijm et al., 1997; Burgess, 2008). In contrast to intracratonic basins, platforms (Fig. 1.3a) accumulate sediment of uniform thickness over continental scales. Platform stratigraphic sequences are transitional into continental margins, intracratonic basins, foreland basins, and other tectonic settings along continental margins (Ingersoll and Busby, 1995; Burgess, 2008). The distinction of distal foreland and platform sequences may be arbitrary, especially during times of high sea level, high carbonate productivity, and broad foreland flexure. Eustatically induced cyclothems are best expressed on platforms (e.g., Heckel, 1984; Klein, 1992; Klein and Kupperman, 1992), and paleolatitude and paleoclimate signals are best isolated in platform sequences (Berry and Wilkinson, 1994). Platforms have generally experienced exposure and erosion during times of supercontinents, and have experienced maximum flooding approximately 100 My after supercontinent breakup (Heller and Angevine, 1985; Cogne et al., 2006).

Active ocean basins

The systematic exponential thermal decay of oceanic lithosphere as it moves away from spreading ridges is expressed by increasing water depth with age of oceanic crust (Sclater et al., 1971; Parsons and Sclater, 1977; Stein and Stein, 1992)

(Fig. 1.3g). As oceanic crust subsides with age and distance from spreading ridges, systematic pelagic and hemipelagic deposits accumulate (Berger, 1973; Heezen et al., 1973; Winterer, 1973; Berger and Winterer, 1974). Carbonate ooze accumulates above the carbonate compensation depth (CCD), which is depressed under areas of high biologic productivity; silica ooze accumulates above the poorly defined silica compensation depth (SCD); and only abyssal clay accumulates below the SCD. The result is a dynamic and predictive stratigraphy relating the age, depth, and paleoaltitude of oceanic crust to oceanic depositional facies. Volcaniclastic and turbidite deposits near magmatic arcs and continental margins complicate predicted stratigraphic sequences on oceanic plates (e.g., Cook, 1975; Ingersoll and Busby, 1995).

Oceanic islands, seamounts, aseismic ridges, and plateaus

Islands, seamounts, ridges, and plateaus thermally subside as oceanic plates migrate away from spreading ridges. Thermal anomalies independent of spreading ridges (e.g., hot spots) create new islands, ridges, and plateaus, which may have complex subsidence histories, dependent on their magmatic histories. Clague (1981) divided the post-volcanic history of seamounts into three sequential stages: subaerial, shallow water, and deep water or bathyal (Ingersoll, 1988; Ingersoll and Busby, 1995). As an island is eroded and subsides, fringing reefs and atolls may form, depending on latitude, climate, and relative sea level (e.g., Jenkyns and Wilson, 1999; Dickinson, 2004). Oceanic features, which may become accreted terranes at convergent margins (e.g., Wrangellia of the North American Cordillera; Ricketts, 2008), range in size from small seamounts to large mafic igneous provinces, such as the Ontong Java Plateau and related features (e.g., Taylor, 2006).

Dormant ocean basins

Dormant ocean basins are floored by oceanic crust, which is neither spreading nor subducting; in other words, there are no active plate margins within or adjoining the basin (Ingersoll and Busby, 1995) (Fig. 1.3h). This setting contrasts with active ocean basins, which include at least one active spreading ridge (e.g., Atlantic, Pacific,

and Indian oceans), and remnant ocean basins, which are small shrinking oceans bounded by at least one subduction zone (e.g., Bay of Bengal and Huon Gulf). The term “dormant” implies that there is no orogenic or taphrogenic activity within or adjacent to the basin; “oceanic” requires that the basin is underlain by oceanic lithosphere, in contrast to intracratonic basins, which are typically underlain by partially rifted continental lithosphere (Ingersoll and Busby, 1995).

Dormant ocean basins are created by two contrasting processes: (1) spreading ridges of nascent ocean basins cease activity (e.g., Gulf of Mexico; Pindell and Dewey, 1982; Pindell, 1985; Dickinson and Lawton, 2001), or (2) backarc basins (either extensional or neutral) are not subducted during suturing of continents and/or arcs (e.g., Black Sea; Okay et al., 1994) or South Caspian basin (Brunet et al., 2003; Vincent et al., 2005). The origin of dormant ocean basins may be difficult to determine because basement and original strata commonly remain deeply buried for hundreds of millions of years following cessation of seafloor spreading (e.g., Tarim and Junggar basins of western China) (e.g., Sengor et al., 1996). Following cessation of plate activity within and around the basin, sediment loading is the dominant subsidence mechanism, although lithospheric thickening due to residual cooling may be important (Ingersoll and Busby, 1995). Dormant ocean basins may have life spans of hundreds of millions of years and may vary considerably in size. The modern Gulf of Mexico, the largest known dormant ocean basin, is filling rapidly along its northern margin (the continental embankment of the Gulf Coast), but still contains oceanic crust with thin sediment cover in the south (e.g., Buffler and Thomas, 1994; Galloway et al., 2000; Dickinson and Lawton, 2001). The South Caspian Basin is small and partially filled with sediment (locally over 20 km thick; Brunet et al., 2003), and yet still is an oceanic basin. In contrast, the Tarim basin has a comparable sediment thickness, but is completely filled. These three basins are likely underlain by oceanic crust, or in the case of Tarim, an oceanic Plateau (Sengor et al., 1996); their long histories of cooling means that they are also underlain by thick and strong mantle lithosphere (Ingersoll and Busby, 1995). When a dormant ocean basin is filled to sea level, it may superficially resemble an intracratonic basin. The former, however, contains 16–20 km of sedimentary strata on top of strong oceanic lithosphere, whereas the latter

contains a few km of sedimentary strata underlain primarily by continental crust, with one or more fossil rifts beneath the basin center. Thus, when in-plate stresses affect dormant ocean basins and their surroundings, deformation usually occurs along their weak boundaries, whereas deformation of intracratonic basins is concentrated along the fossil rifts underlying their interiors. Foreland basins may form above the edges of dormant ocean basins during contractional deformation (e.g., the margins of the modern Tarim basin). Intracratonic basins may experience renewed subsidence or inversion tectonics (e.g., the modern North Sea) (Cooper and Williams, 1989; Cameron et al., 1992).

CONVERGENT SETTINGS

Arc-trench systems

Arc-trench systems may be categorized into three fundamental types: (1) extensional (2) neutral, and (3) compressional (Dickinson and Seely, 1979; Dewey, 1980) (Fig. 1.4). Arc-trench systems with significant strike slip may be considered a fourth type (Dorobek, 2008); strike-slip faults may occur in all types of arc-trench system, but they are especially common in strongly coupled systems experiencing oblique convergence (Beck, 1983). Many parameters determine the behavior of arc-trench systems, but the most important factors appear to be (1) convergence rate (2) slab age, and (3) slab dip (Molnar and Atwater, 1978; Uyeda and Kanamori, 1979; Jarrard, 1986; Kanamori, 1986), based on analyses of modern arc-trench systems (although see Cruciani et al., 2005, for an alternative interpretation). A major question arises from these analyses of contemporary Earth: is the present arrangement of spreading ridges and arc-trench systems typical of Earth history or an unusual configuration? Almost all modern east-facing arcs (e.g., Marianas) are extensional, with subduction of old lithosphere at steep angles. Almost all west-facing arcs (e.g., Andes) are compressional, with subduction of young lithosphere at shallow angles. Most south-facing arcs (e.g., Aleutians) are neutral, with subduction of middle-aged lithosphere at moderate angles. There are no north-facing arcs. Thus, it is very difficult to separate the covarying parameters of slab age, slab dip, facing direction, and type of arc-trench system. There is growing consensus (although see Schellart, 2007, 2008, for a contrary view) that facing direction of arc-trench systems

may be the fundamental determinant of the behavior of arc-trench systems because of westward tidal lag of the eastward rotating planet (e.g., Bostrom, 1971; Moore, 1973; Dickinson, 1978; Doglioni, 1994; Doglioni et al., 1999). If this is the case today, then it should have been the case throughout Earth history because of the constancy of eastward planetary rotation. Therefore, models for ancient arc-trench systems must account for the azimuth of their facing directions when they were active. Lack of recognition of this fundamental characteristic of arc-trench systems has resulted in many invalid analog models of ancient mountain belts (Dickinson, 2008).

Dickinson (1974a, 1974b), Ingersoll (1988), Ingersoll and Busby (1995) and Dorobek (2008) summarized tectonic settings and subsidence mechanisms of the diverse basin types related to arc-trench systems. Ingersoll and Busby (1995), and Smith and Landis (1995) also discussed construction and erosion of arc edifices that provide most sediment to neighboring basins.

The distinction of forearc, intra-arc, and backarc basins is not always clear. Intra-arc basins are defined as thick volcanic-volcaniclastic and other sedimentary accumulations along the arc platform, which is formed of overlapping or superposed volcanoes. The presence of vent-proximal volcanic rocks and related intrusions is critical to the recognition of intra-arc basins in the geologic record, since arc-derived volcaniclastic material may be spread into forearc, backarc, and other basins. A more general term, “arc massif,” refers to crust generated by arc magmatic processes (Dickinson, 1974a, 1974b), and arc crust may underlie a much broader region than the arc platform. The distinction of forearc and intra-arc basins is also discussed by Dickinson (1995). Many backarc basins form by rifting within the arc platform (Marsaglia, 1995), and were intra-arc basins in their early stages. Also, forearc, intra-arc and backarc settings change temporally and are superposed on each other due to both gradual evolution and sudden reorganization of arc-trench systems resulting from collisional events, plate reorganization, and changes in plate kinematics.

Trenches

Karig and Sharman (1975), Schweller and Kulm (1978), Thornburg and Kulm (1987), and Underwood and Moore (1995) summarized the

dynamic nature of sedimentation and tectonics in active trenches (Fig. 1.4a). The sediment wedge of a trench is in dynamic equilibrium when subduction rate and angle, sediment thickness on the

oceanic plate, rate of sedimentation, and distribution of sediment within the trench are constant. Thornburg and Kulm (1987) provided documentation of the dynamic interaction of longitudinally




Fig. 1.4. True-scale actualistic analog models for sedimentary basins in convergent settings. A remnant arc is shown on the left side of (A). Trench, trench-slope and forearc basins are labeled only in (A and B), but they are associated with all types of arc-trench systems. Intra-arc basins may be associated with any magmatic arc, but they are more common and more likely to be preserved in extensional and neutral settings (A, B, and C). Hinterland basins may form in compressional arc-trench systems (D, E), or in collisional systems (F and Figure 1.6A–B). Remnant ocean basins form between any colliding crustal margins; a compressional arc-trench system is shown converging with an intraplate margin in (E). Wedgetop basins may form in any compressional setting; a proforeland example is shown in (F). If neutral continental arc-trench systems (C) become extensional, then they may evolve into extensional oceanic systems (A). Symbols same as in Figure 1.3; slab-generated melts = red.

transported material (trench wedge with axial channel) and transversely fed material (trench fan). With increasing transverse supply of sediment to the trench, the axial channel of the trench wedge is forced seaward and the trench wedge widens. Contrasts in dynamic trench-fill processes help determine not only trench bathymetry and depositional systems, but also accretionary architecture (Thornburg and Kulm, 1987; Underwood and Moore, 1995). This dynamic model may be useful in reconstruction of sedimentary and tectonic processes in trenches, as expressed in ancient subduction complexes.

Scholl et al. (1980) developed conceptual models relating accretionary processes to subduction and sedimentary parameters that influence forearc and trench characteristics. Cloos and Shreve (1988a, 1988b) developed quantitative models for processes at greater depths in subduction zones, which affect the nature of deformation and metamorphism, and the overall character of forearcs. Reconstruction of sedimentary systems within the transient settings of ancient trenches is highly problematic because of difficulty of studying modern systems at such great water depths, contrast in scale of resolution between modern and ancient studies, and extreme structural deformation that occurs within subduction environments (Underwood and Moore, 1995). Nonetheless, advances in technology and continuing studies of modern and ancient systems are providing incremental improvements in our understanding of the sedimentary and tectonic systems (e.g., Maldonado et al., 1994; Mountney and Westbrook, 1996; Leverenz, 2000; Kopp and Kukowski, 2003).

Trench-slope basins

Moore and Karig (1976) developed a model for sedimentation in small ponded basins along inner trench walls (Fig. 1.4b). Deformation within and on subduction complexes results in irregular bathymetry; turbidites are ponded within resulting trench-slope basins. Average width, sediment thickness, and age of basins increase up slope due to progressive uplift of deformed material and widening of fault spacing during dewatering and deformation of offscraped sediment. In ancient subduction complexes, trench-slope basins are filled with relatively undeformed, locally derived turbidites surrounded by highly deformed accreted material of variable origin. Contacts

between trench-slope basins and accreted material are both depositional and tectonic. Moore and Karig's (1976) model was developed for Nias Island near Sumatra, an area of rapid accretion of thick sediments. Their model is less useful for sediment-starved forearc areas. Allen et al. (2008), and Hall and Smyth (2008) provided additional details concerning the Nias Island area and the Andaman Islands, including some alternative interpretations. Nonetheless, Moore and Karig's (1976) general principles governing the development of sedimentary basins on the lower trench slope are fundamental to reconstructing ancient subduction complexes.

Underwood and Moore (1995), Aalto and Miller (1999), Underwood et al. (2003), and Allen et al. (2008) discussed additional examples of both modern and ancient trench-slope basins, and their significance in paleotectonic reconstructions.

Forearc basins

Dickinson and Seely (1979) and Dickinson (1995) provided a classification of arc-trench systems, similar to Dewey's (1980), and outlined plate-tectonic controls governing subduction initiation and forearc development (Fig. 1.4b). Factors controlling forearc geometry include the (1) initial setting (2) sediment thickness on subducting plate (3) rate of sediment supply to trench (4) rate of sediment supply to forearc area (5) rate and orientation of subduction, and (6) time since initiation of subduction. Arc-trench gaps tend to widen through time (Dickinson, 1973) due to prograde accretion at trenches and retrograde migration of magmatic arcs following subduction initiation. Prograde accretion is especially rapid where thick sequences of sediment are accreted. The net result of widening of the arc-trench gap is the general tendency for forearc basins to enlarge through time (e.g., Great Valley forearc basin; Ingersoll, 1979, 1982; Dickinson, 1995).

Forearc basins include the following types (Dickinson and Seely, 1979; Dickinson, 1995): (1) intramassif (transitional to intra-arc) (2) accretionary (trench-slope) (3) residual (lying on oceanic or transitional crust trapped behind the trench when subduction initiated) (4) constructed (lying across the boundary of arc massif and subduction complex), and (5) composite (combination of above settings). Residual and constructed basins tend to evolve into composite basins; commonly,

this evolutionary trend is accompanied by filling and shallowing of forearc basins.

Stern and Bloomer (1992) discussed lithospheric extension along the front edge of the overriding plate at the time of initiation of the Mariana subduction zone (Oligocene). This type of extension to form new crust is only likely within 10–20 My of initiation of an intraoceanic subduction zone within old (strong) oceanic lithosphere, where slab rollback begins as soon as subduction initiates, and the weakest part of the overriding plate is near the edge. Soon after subduction begins, forearc areas are cooled by the cold subducting oceanic lithosphere; thus, mature intraoceanic forearcs tend to be underlain by cold and strong lithosphere, and resist crustal extension (i.e., Vink et al., 1984; Steckler and tenBrink, 1986; Dickinson, 1995). Normal faults are common in shallow levels of accretionary wedges (e.g., Platt, 1986; Underwood and Moore, 1995), but crustal rifting to form new crust has not been documented in any modern forearc, and is unlikely to have occurred in any ancient forearcs (Ingersoll, 2000). In contrast, arc axes of mature intraoceanic systems tend to be the weakest parts of overriding plates, and extension is accommodated by intra-arc and backarc spreading (Marsaglia, 1995).

Several recent studies of both modern and ancient forearc basins have verified the usefulness of the general models discussed by Dickinson (1995) (e.g., Einsele et al., 1994; Van der Werff, 1996; Mountney and Westbrook, 1997; Constenius et al., 2000; Trop, 2008).

Intra-arc basins

The origin of basins within magmatic arcs (Fig. 1.4a) is, in general, poorly understood, largely due to the paucity of studies that integrate volcanology, sedimentology and basin analysis (Ingersoll, 1988; Ingersoll and Busby, 1995). A further deterrent for many sedimentologists is the fact that arcs are characterized by high heat flow with steep geothermal gradients and intense magmatism, and are commonly subjected to crustal shortening at some time in their history; therefore, experience in “seeing through” the overprints of hydrothermal alteration, metamorphism and deformation is required. Fisher and Schmincke (1984), Cas and Wright (1987), Cas and Busby-Spera (1991), Fisher and Smith (1991), and Smith and Landis (1995) provided excellent summaries of knowledge prior to 1995.

Oceanic intra-arc basins

There are at least three major types of depocenters for volcanic and sedimentary accumulations within arcs (Ingersoll and Busby, 1995; Smith and Landis, 1995). Depocenters may occur in low regions between volcanoes and along their flanks, although these have high preservation potential only below sealevel (i.e., generally in oceanic arcs). Also, depocenters with high preservation potential may form when the axis of arc volcanism shifts to a new position on an oceanic arc platform, thus creating a low region between the active chain and the abandoned chain. Smith and Landis (1995) referred to both of these types of intra-arc basin as “volcano-bounded basins.” They also referred to “fault-bounded basins,” which are rapidly subsiding basins where tectonic structures, rather than constructional volcanic features, account for relief along the basin margins (Ingersoll and Busby, 1995).

Additional studies of sedimentation in oceanic intra-arc settings include those of Robertson and Degnan (1994), Fackler-Adams and Busby (1998), Sowerbutts and Underhill (1998), Sowerbutts (2000), Clift et al. (2005), and Busby et al. (2006).

Continental intra-arc basins

The most important mechanisms for accumulating and preserving thick stratigraphic successions in continental arcs appear to be, in descending scale (Busby-Spera, 1988b; Busby-Spera et al., 1990): (1) plate-margin-scale extension or transtension (2) extension on a more local scale during pluton or batholith emplacement, and (3) localized subsidence of calderas during large-volume ignimbrite eruptions. Plate-margin-scale extension or transtension produces belts of continental-arc sequences that are continuous or semi-continuous over hundreds to thousands of kilometers and record high rates of subsidence over tens of millions of years. The effects of extension in the roofs of plutons or batholiths (e.g., Tobisch et al., 1986) may be difficult to distinguish from plate-margin-scale extension, but the former should operate over shorter time scales (i.e., less than a few million years), and should not, by itself, produce a low-standing arc capable of trapping sediment derived from outside the arc. Continental calderas form small (10–60 km wide) but deep (1–4 km) depocenters for ignimbrite erupted during caldera collapse, as well as for volcanic and

sedimentary strata ponded within the caldera after collapse (e.g., Riggs and Busby-Spera, 1991; Lipman, 1992; Schermer and Busby, 1994). Funk et al. (2009) provided a detailed description of the Cenozoic tectonics of the intra-arc basins of Nicaragua and El Salvador.

Backarc basins

There are two types of backarc basins: (1) oceanic basins behind intraoceanic magmatic arcs, and (2) continental basins behind continental-margin arcs that lack foreland foldthrust belts (Ingersoll and Busby, 1995). Many backarc basins are extensional in origin, forming by rifting and seafloor spreading (Fig. 1.4a) (Marsaglia, 1995). These commonly originate through rifting of the arc, either along its axis (intra-arc) or immediately to the front or rear of its axis. The term “interarc basin” (Karig, 1970) has been widely superseded by the term “backarc basin,” but it can be used where rifting has occurred along or near an arc axis, thus eventually producing a remnant arc behind the backarc basin. The presence or preservation of a remnant arc is not a necessary condition for recognition of a backarc basin (Taylor and Karner, 1983).

Many backarc basins are nonextensional (Marsaglia, 1995), forming under neutral stress regimes (Fig. 1.4b). The most common type of nonextensional backarc basin consists of old ocean basins trapped during plate reorganization (e.g., the Bering Sea). Also, nonextensional backarc basins develop on continental crust (Fig. 1.4c) (e.g., Sunda shelf of Indonesia). Backarc shortening may occur in intraoceanic arc-trench systems involved in early stages of collision with buoyant crust (e.g., Greater Antilles and eastern Indonesia) (tenBrink et al., 2009); this shortening could represent early stages of subduction initiation during polarity reversal.

Oceanic backarc basins

Modern oceanic backarc basins may be distinguished from other ocean basins petrologically or by their positions behind active or inactive arc-trench systems (Taylor and Karner, 1983; Marsaglia, 1995). Such diagnostic features are commonly not preserved in ancient backarc basins, which commonly undergo metamorphic and structural modifications during emplacement in orogenic zones as ophiolites. The nature and

timing of deposition of sediment on top of ophiolite sections have proven more diagnostic for determining original plate-tectonic settings (e.g., Tanner and Rex, 1979; Hopson et al., 1981, 2008; Kimbrough, 1984; Busby-Spera, 1988a; Robertson, 1989).

The most detailed study of a backarc volcanoclastic apron and its substrate comes from Middle Jurassic rocks in Mexico (Busby-Spera, 1987, 1988a; Critelli et al., 2002). That study supported Karig and Moore’s (1975) assertion that oceanic backarc basins isolated from terrigenous sediment influx may show the following simple, uniform sedimentation patterns: (1) lateral and vertical differentiation of facies due to progradation of a thick volcanoclastic apron into a widening backarc basin; such an apron may extend for more than 100 km from a volcanic island and grow to a thickness of 5 km in 5 My (Lonsdale, 1975). This phase is followed by (2) blanketing of the apron with a thin sheet of mud and sand eroded from the arc after volcanism and spreading have ceased. This cycle reflects the temporal episodicity of seafloor spreading in oceanic backarc basins, which appear to form in 10–15 My or less (Taylor and Karner, 1983). As a result, extensional oceanic backarc basins generally have shorter life spans than intra-arc basins (Fig. 1.2). The shorter life span reflects temporal episodicity of extensional oceanic backarc basins, the most common type of backarc basin. In contrast, arcs may undergo episodic extension for many tens of millions of years, particularly in continental settings. Although backarc basins and their fill make an important contribution to orogenic belts, most ancient oceanic backarc basins have probably been subducted; the frontal-arc sides of backarc basins may be preferentially preserved in the geologic record (Busby-Spera, 1988a).

Marsaglia (1995) discussed modern and ancient backarc basins, both oceanic and continental, and both extensional and neutral settings, although her emphasis was on extensional backarcs of the western Pacific (also, see Klein, 1985). More recent publications addressed evolution of the complex extensional backarc basins of the Western Mediterranean (e.g., Maillard and Mauffret, 1999; Pascucci et al., 1999; Mattei et al., 2002; Rollet et al., 2002). Sibuet et al. (1998) synthesized the tectonic and magmatic evolution of the Okinawa Trough, and Critelli et al. (2002) analyzed the Jurassic backarc basin of Cedros Island, Baja California using sandstone petrology in conjunction with stratigraphy and sedimentology. Less

attention has been paid to nonextensional backarc basins (e.g., Aleutian Basin of the Bering Sea), formed by trapping of oceanic crust behind intraoceanic arcs following initiation of intraoceanic subduction zones (Ben-Avraham and Uyeda, 1983; Tamaki and Honza, 1991). Part of the Caribbean Sea, the West Philippine Basin, and perhaps part of the Okhotsk Sea north of the Kuril Basin also consist of oceanic crust trapped in backarc settings (Uyeda and Ben-Avraham, 1972; Scholl et al., 1975; Ben-Avraham and Uyeda, 1983; Marsaglia, 1995). These backarc basins generally have longer life spans and greater preservation potential than indicated in Figure 1.2, especially if they evolve into dormant ocean basins. Oceanic crust trapped in backarc settings is likely to be as complex as any other oceanic crust, with oceanic plateaus, continental fragments, and transform faults (Marsaglia, 1995).

Continental backarc basins

A modern continental backarc in a neutral stress regime is the Sunda Shelf of Indonesia (Hamilton, 1979; Ingersoll, 1988; Ingersoll and Busby, 1995). DeCelles and Giles (1996) utilized the Sunda Shelf as an example of the earliest stages of development of a retroforeland basin (see below), but Moss and McCarthy (1997) disputed this interpretation and suggested that there is no retroarc shortening behind the Indonesian magmatic arc (also see DeCelles and Giles, 1997). Moss and McCarthy (1997) interpreted part of the Sunda Shelf area as having a previous extensional history. In any case, an extensional backarc can evolve into a neutral backarc, which can evolve into a retroforeland basin. Variable stress regimes in backarc and retroarc settings are common.

A similar series of backarc-to-retroforeland basins developed during the Mesozoic in the western USA (Dickinson, 1981a, 1981b; Lawton, 1994; Ingersoll, 1997, 2008a). Following the Permian-Triassic Sonoma orogeny, a continental-margin magmatic arc developed following subduction initiation (Hamilton, 1969; Schweickert, 1976, 1978; Busby-Spera, 1988b). Shallow-marine and non-marine conditions prevailed in the dynamically neutral backarc area from mid-Triassic to Late Jurassic (Dickinson, 1981a, 1981b; Lawton, 1994; Ingersoll, 1997, 2008a), although crustal extension may have characterized some parts of the backarc (Wyld, 2000, 2002). The backarc evolved into a retroforeland as shortening initiated during the

Jurassic (Oldow, 1984; Oldow et al., 1989; Lawton, 1994; Wyld, 2002). Thus, relative timing of extension, neutrality, and shortening in the Sunda Shelf area is similar to the interpreted sequence of events in the Mesozoic backarc-retroforeland of the western USA.

Retroforeland basins

Compressional arc-trench systems commonly develop foreland basins behind arcs due to partial subduction of continental crust beneath arc orogens (Dickinson, 1974b; Dewey, 1980; Ingersoll, 1988; Ingersoll and Busby, 1995; DeCelles and Giles, 1996). “Foreland basin” is a pre-plate-tectonic term used to describe a basin between an orogenic belt and a craton (Allen et al., 1986). Dickinson (1974b) proposed that the term “retroarc” be used to describe foreland basins formed behind compressional arcs, in contrast to “peripheral” foreland basins formed on subducting plates during continental collisions. Thus, although “backarc” and “retroarc” are literally synonymous, the former is used for extensional and neutral arc-trench systems, whereas the latter is used for compressional arc-trench systems.

Willett et al. (1993), Johnson and Beaumont (1995), Beaumont et al. (1996), and Naylor and Sinclair (2008) modified Dickinson’s (1974b) original nomenclature for foreland basins by shortening “retroarc foreland” to “retroforeland” and changing “peripheral foreland” to “proforeland”. This nomenclature is adopted herein, with the clear stipulation that retroforeland basins form on the upper plates of convergent margins and proforeland basins form on the lower plates of convergent margins. Retroforelands tend to have longer histories than proforelands because the former commonly initiate during subduction of oceanic lithosphere (e.g., Andean retroforeland), whereas proforelands do not exist until buoyant continental crust enters subduction zones (inducing collision) (Dickinson, 1974b; Ingersoll, 1988; Cloos, 1993; Ingersoll and Busby, 1995). In order to clarify these distinctions, I propose that retroforelands be subdivided into retroarc forelands (formed behind continental-margin arcs, e.g., the Andes) and collisional retroforelands (formed on the overriding continental plate during continental collision, e.g., South Alpine foreland basin). The general term “retroforeland” may be used for any foreland on the upper plate of a convergent margin, whereas the more restricted terms would

be used to designate whether oceanic or continental crust was being subducted on the opposite side of the orogen. Approximately one third of active magmatic arcs have associated retroarc forelands, whereas arc activity commonly ceases as proforelands and collisional retroforelands develop.

Retroarc foreland basins

Jordan (1981) presented an analysis of the asymmetric Cretaceous retroarc foreland basin associated with the Idaho-Wyoming thrust belt. She used a two-dimensional elastic model to show how thrust loading and sedimentary loading resulted in broad flexure of the lithosphere (Fig. 1.4e). The location of maximum flexure migrated eastward as thrusting migrated eastward. The area of subsidence broadened due to erosional and depositional redistribution of part of the thrust load, and possibly enhanced by high eustatic sea level of the Late Cretaceous. Comparison of modeled basin and basement geometries with isopach maps provides tests of possible values of flexural rigidity of the lithosphere. The modern sub-Andean thrust belt and foreland basin have similar topography to that proposed for the Cretaceous of the Idaho-Wyoming system (Jordan, 1995). Topography is controlled by thrust-fault geometry and isostatic subsidence.

The models presented by Jordan (1981) and Beaumont (1981) are broadly applicable to other retroarc foreland basins (Jordan, 1995; DeCelles and Giles, 1996; Catuneanu, 2004). These and derivative models demonstrate that tectonic activity in foreland foldthrust belts is the primary cause of subsidence in associated foreland basins (Price, 1973). Sedimentary redistribution, autocyclic sedimentary processes, dynamic effects of asthenospheric circulation (e.g., Gurnis, 1993; Burgess et al., 1997), and eustatic sealevel changes are important modifying factors in terms of regressive-transgressive sequences, but compressional tectonics behind the arc-trench system is the driving force. The Cretaceous seaway of North America was largely the result of this compressional tectonic activity (combined with high eustatic sealevel) (Dickinson, 1976b, 1981a). Details concerning timing of thrusting and initial sedimentary response to thrusting within the Idaho-Wyoming thrust belt have been debated (e.g., Heller et al., 1986), but the essential role of compressional tectonics in creating retroarc foreland basins is clear (Price, 1973; DeCelles and Giles, 1996).

Jordan (1995) updated her analysis of the Cretaceous retroarc foreland of North America, synthesized the Neogene to Holocene retroarc foreland of South America, and discussed general models for retroarc foreland basins. DeCelles and Giles (1996) synthesized foreland-basin systems, including subdivision into four discrete depozones: wedgetop, foredeep, forebulge, and backbulge. All four depozones occur in the modern retroforeland east of the central Andes (Horton and DeCelles, 1997). DeCelles and Horton (2003) applied this model to interpretation of tertiary foreland strata of Bolivia, and concluded that approximately 1000 km of foreland crust have been thrust westward beneath the Andean orogenic belt. Fildani et al. (2003), Abascal (2005), Gomez et al. (2005), Hermoza et al. (2005), Horton (2005), and Uba et al. (2005) provided detailed analyses of diverse parts of the Andean retroforeland system.

DeCelles and Giles's (1996) subdivisions apply equally to all types of foreland-basin systems. Wedgetop basins are discussed separately below because their character is tied directly to fault dynamics, whereas the other three depozones are created by flexural loading of the overall thrust belt. Wedgetop and backbulge depozones have not been considered in most models for foreland evolution (DeCelles and Giles, 1996). Foreland models that include wedgetop depozones must utilize doubly tapered prisms in cross section, rather than the wedge that is commonly used as a simplification (DeCelles and Giles, 1996). Dorobek and Ross (1995) illustrated many types of models and case studies that improve our understanding of foreland basins.

Collisional retroforeland basins

The South Alpine collisional retroforeland basin developed synchronously with Alpine orogenesis as the European plate subducted beneath Adria (Bertotti et al., 1998; Carrapa, 2009) (Fig. 1.4f). Late Cretaceous evolution of this foreland began in a retroarc setting, but developed into a collisional retroforeland concurrent with Alpine orogenesis and development of the better known North Alpine (molasse) proforeland basin (Bertotti et al., 1998). Bertotti et al. (1998) suggested progressive weakening of the flexed Adria plate through time. Carrapa and Garcia-Castellanos (2005) demonstrated that the Tertiary Piedmont basin of the western Po Plain formed by Alpine retroforeland flexure of a visco-elastic plate during

Oligocene-Miocene time. Zattin et al. (2003) used provenance data from the Venetian basin to document the sequence of deformation in the eastern South Alpine collisional retroforeland. Apennine orogenesis has superposed the Po Valley proforeland basin on the older retroforeland basin, thus making this a hybrid foreland basin (Ingersoll and Busby, 1995; Miall, 1995).

The Triassic-Jurassic foreland sequences of the Ordos basin of central China represent deposition in a collisional retroforeland basin related to suturing of the North and South China blocks (Sitian et al., 1995; Ritts et al., 2009).

Broken retroforeland basins

Low-angle subduction beneath compressional arc-trench systems may result in basement-involved deformation within retroarc foreland basins (Fig. 1.4d) (Dickinson and Snyder, 1978; Jordan, 1995). The Rocky Mountain region of the western USA is the best-known ancient example of this style of deformation; similar modern provinces have been documented in the Andean foreland (e.g., Jordan et al., 1983a, 1983b; Jordan and Allmendinger, 1986; Jordan, 1995).

Chapin and Cather (1981), Dickinson et al. (1988, 1990), Cather and Chapin (1990), Dickinson (1990), Hansen (1990), and Lawton (2008) synthesized and discussed controls on latest Cretaceous through Eocene (Laramide) sedimentation and basin formation of the Colorado Plateau and Rocky Mountain area. They agreed that diverse types of uplifts and basins formed during this period, but they disagreed on paleodrainage networks, the relative importance of strike-slip deformation along the east side of the Colorado Plateau, and whether the Laramide orogeny occurred in two distinct stages or was a continuum of responses to a generally homogeneous strain field. Yin and Ingersoll (1997) and Ingersoll (2001) presented a model for Laramide crustal strain and basin evolution in northern New Mexico and southern Colorado, which is consistent with a generally homogeneous strain field. Hoy and Ridgway (1997) illustrated the complex structural, stratigraphic and sedimentologic relations that commonly developed along the margins of intraforeland uplifts in Wyoming. Cardozo and Jordan (2001), Davila and Astini (2003), Sobel and Strecker (2003), and Hilley and Strecker (2005) studied broken retroforeland basins and associated uplifts in Argentina.

Remnant ocean basins

Intense deformation occurs in suture belts during the attempted subduction of buoyant (nonsubductable) continental or magmatic-arc crust (e.g., Cloos, 1993). Suture belts can involve rifted continental margins and continental-margin magmatic arcs (terminal closing of an ocean basin) or various combinations of arcs and continental margins (Fig. 1.4e–f). Colliding continents tend to be irregular, and great variability of timing, structural deformation, sediment dispersal patterns and preservability occurs along strike (Dewey and Burke, 1974).

Graham et al. (1975) and Ingersoll et al. (1995, 2003) used Cenozoic development of the Himalayan-Bengal system as an analog for late Paleozoic development of the Appalachian-Ouachita system, and proposed a general model for sediment dispersal related to sequentially suturing orogenic belts. “Most sediment shed from orogenic highlands formed by continental collisions pours longitudinally through deltaic complexes into remnant ocean basins as turbidites that are subsequently deformed and incorporated into the orogenic belts as collision sutures lengthen” (Graham et al., 1975, 273). This model provides a general explanation for many synorogenic flysch and molasse deposits associated with suture belts, although many units called “flysch” and “molasse” have different tectonic settings (Ingersoll et al., 1995, 2003; Miall, 1995).

North American examples of arc-continent collisions, with variable volumes of remnant-ocean-basin flysch, include the Ordovician Taconic orogeny of the Appalachians (e.g., Rowley and Kidd, 1981; Stanley and Ratcliffe, 1985; Lash, 1988; Bradley, 1989; Bradley and Kidd, 1991) and the Devonian-Mississippian Antler orogeny of the Cordillera (e.g., Speed and Sleep, 1982; Dickinson et al., 1983). In both cases, it is difficult to clearly distinguish remnant ocean basins from incipient proforeland basins as the depositional sites for “flysch” (Ingersoll and Busby, 1995; Ingersoll et al., 1995; Miall, 1995).

Ingersoll et al. (1995, 2003) reviewed several remnant ocean basins, both modern and ancient, and demonstrated that submarine fans in remnant ocean basins represent the largest accumulations of sediment on Earth. The modern Bengal Fan is the largest sediment body and the Indus Fan is second largest; both are derived from the greatest uplifted area on Earth, the Tibetan Plateau and

Himalaya. The Triassic Songpan-Ganzi complex of northern Tibet and the Carboniferous-Permian Ouachita-Marathon flysch of Arkansas, Oklahoma, and Texas also were deposited in remnant ocean basins adjoining uplifted continental suture belts, and they are of comparable size to the Bengal and Indus fans, although deformation during suturing makes their reconstruction difficult (Ingersoll et al., 2003). No known or suggested mechanism can produce sediment masses of comparable volume. Many additional examples of remnant ocean basins, associated with both continent-continent and continent-intraoceanic-arc collisions, are discussed by Ingersoll et al. (2003).

Proforeland basins

As continental collision occurs between a rifted continental margin and the subduction zone of an arc-trench system, a tectonic load is placed on the rifted margin, first below sealevel, and later subaerially (Dickinson, 1974b; Ingersoll, 1988; Miall, 1995). A proforeland basin forms as the elastic lithosphere flexes under the encroaching dynamic load (Fig. 1.4f). Normal faulting in front of the dynamic load and uplift of a forebulge are initial responses to flexure as the dynamic load encroaches on the foreland (Bradley and Kidd, 1991; Miall, 1995; DeCelles and Giles, 1996).

Discrimination of ancient proforelands and collisional retroforelands (Fig. 1.4f) is difficult, but may be possible based on the following characteristics (Ingersoll, 1988; Ingersoll and Busby, 1995): (1) polarity of magmatic arc (2) presence of oceanic subduction complex associated with earliest phases of proforeland (3) greater water depths in proforeland (foredeep stage) (4) asymmetry of suture belt (closer to proforeland) (5) protracted development of retroforeland (long-term arc evolution) versus discrete development of proforeland (terminal ocean closure without precursor), and (6) possible volcanoclastic input to retroforeland, especially during early development, versus minimal volcanoclastic input to proforeland.

Stockmal et al. (1986) provided a dynamic 2D model for the development of proforeland basins, following finite times of rifting. They modified the model of Speed and Sleep (1982), and demonstrated the effects of rifted-margin age and topography on lithospheric flexure and basin development. The primary effect of age shows up as a higher flexural forebulge and thicker trench fill during earlier stages of attempted

subduction of an old (120 my) margin. Subsequent development is relatively insensitive to margin age. Foreland-basin subsidence is sensitive to overthrust load, with depths possibly exceeding 10 km. Crustal thickness may reach 70 km during the compressional phase (e.g., Himalayas). Tens of kilometers of uplift and erosion, of both the allochthon and the proximal foreland basin, are predicted during and after deformation. Most erosional detritus is deposited elsewhere due to uplift within the foreland; longitudinal transport into remnant ocean basins results (Graham et al., 1975; Ingersoll et al., 1995, 2003; Miall, 1995). Thick overthrusts with low topographic expression are to be expected where broad, attenuated rifted continental margins have been pulled into subduction zones (Stockmal et al., 1986).

Miall (1995) discussed all “collision-related foreland basins,” which include both proforeland and collisional retroforeland basins. Several studies of foreland basins were presented in Dorobek and Ross (1995) and Mascle et al. (1998). The rapidly expanding literature on proforeland basins includes study of the Pyrenean (e.g., Arenas et al., 2001; Jones et al., 2004), Alpine (e.g., Sinclair, 1997; Gupta and Allen, 2000; Allen et al., 2001; Pfiffner et al., 2002; Kempf and Pfiffner, 2004), Apennine (e.g., Bertotti et al., 2001; Lucente, 2004), Carpathian (e.g., Zoetemeijer et al., 1999; Tarapouca et al., 2004; Leever et al., 2006), Zagros (e.g., Alavi, 2004), Himalayan (e.g., Pivnik and Wells, 1996; DeCelles et al., 1998, 2001; Najman and Garzanti, 2000; Najman et al., 2004), Longmen Shan (e.g., Yong et al., 2003; Meng et al., 2005), West Taiwan (e.g., Chen et al., 2001; Lin and Watts, 2002), Papuan (e.g., Galewsky et al., 1996; Haddad and Watts, 1999), Appalachian (e.g., Thomas, 1995; Castle, 2001), and Proterozoic (e.g., Saylor, 2003) forelands.

Wedgetop basins

Ori and Friend (1984) defined “piggyback basins” as basins that form and fill while being carried on moving thrust sheets. DeCelles and Giles (1996) suggested “wedgetop” as a more general descriptive term, which includes both “piggyback” and “thrust-top” basins (Fig. 1.4f). Wedgetop basins are dynamic settings for sediment accumulation; most sediment is derived from associated foldthrust belts, with subordinate input from arc and basement terranes (Critelli and Le Pera, 1994; Trop and

Ridgway, 1997). The foldthrust belts can be in proforeland, retroforeland, or transpressional settings (Ingersoll and Busby, 1995). Wedgetop basins share characteristics with trench-slope basins. The submarine environment of southern Taiwan illustrates the transition from forearc/trench-slope/trench environments west of the Luzon Arc to orogenic-wedge/wedgetop/foredeep of the Taiwan collision zone (Chiang et al., 2004). This transition occurs where subduction of oceanic crust beneath the Luzon Arc evolves into attempted subduction of Asian continental crust to form the Taiwan suture zone. Both trench-slope and wedgetop basins have low preservation potential due to their development on growing thrust belts; therefore, they are generally found only in young orogenic systems (e.g., Burbank and Tahirkheli, 1985) (Fig. 1.2).

Jordan (1995), Miall (1995), Nilsen and Sylvester (1995), and Talling et al. (1995) provided additional insights regarding wedgetop basins and their relations with compressional processes. Foreland deposition can evolve into wedgetop deposition as frontal faults prograde into foreland basins (DeCelles and Giles, 1996; Pivnik and Khan, 1996); where rapid “sled run” advance of the thrust front occurs due to low basal friction, most foreland deposition may occur in diverse wedgetop basins (Evans and Elliott, 1999; Ford, 2004). Horton (1998) documented extensive Oligocene to Upper Miocene synorogenic wedgetop deposits in the Eastern Cordillera of southern Bolivia. He suggested that longterm preservation of these deposits (for up to 30 My) may reflect the semiarid climate of the region; their presence adds mass to the orogenic wedge, which, in turn, promotes propagation of the thrust front (Horton, 1998). Wedgetop basins form in both thin-skinned and thick-skinned (basement-involved) contractional settings (e.g., Casas-Sainz et al., 2000).

Hinterland basins

Horton (chapter 21, this volume) described two classes of hinterland basins: those formed in non-collisional retroarc orogens (e.g., Andes) and collisional orogens (e.g., Himalaya-Tibet). Because “hinterland” refers to the “internal” parts of orogens, opposite the direction of vergence of folds and faults, the term denotes a direction relative to a foldthrust belt. In a literal sense, all parts of an orogen behind a foldthrust belt constitute the hinterland, regardless of genetic origin; for example, a

retroforeland is part of the hinterland of a proforeland, and vice versa, in two-sided orogens (e.g., Alps and Pyrenees). Horton (chapter 21, this volume) and I define “hinterland basin” in the more restricted sense of basins within orogenic belts that do not fall into any other category described herein.

Hinterland basins record nonmarine sedimentation, usually at high elevations, that formed on thickened continental crust (Horton, chapter 21, this volume) (Fig. 1.4d). As a result, they have low preservation potential, and relatively short life spans (Fig. 1.2), although some basins have life spans of tens of millions of years (e.g., Altiplano; Horton et al., 2002). Extensional, contractional, and strike-slip processes can create the accommodation space for hinterland basins, with fault-induced crustal thinning, sedimentary and volcanic loading, and tectonic loading causing subsidence (Fig. 1.1). Horton (chapter 21, this volume) described two modes of hinterland-basin evolution: (1) basins that developed as new faults became active, and (2) basins that overprinted former foreland basins as the deformation front advanced.

Additional examples of hinterland basins are discussed by Burchfiel et al. (1992), Garzione et al. (2003), Alcicek (2007), DeCelles et al. (2007), Giovanni et al. (2010), and Saylor et al. (2010).

TRANSFORM SETTINGS

Strike-slip systems

The complexity and variability of sedimentary basins associated with strike-slip faults are almost as great as for all other types of basins (Ingersoll and Busby, 1995). Transform faults in oceanic lithosphere generally behave according to the plate-tectonic model, whereas strike-slip faults in continental lithosphere are extremely complex and difficult to fit into a model involving rigid plates.

Strike-slip faults within continental crust are likely to experience alternating periods of extension and compression as slip directions adjust along major crustal faults (Crowell, 1974a, 1974b; Reading, 1980). Thus, opening and closing of basins along strike-slip faults is analogous, at smaller spatial and time scales, to the opening and closing of ocean basins (the Wilson Cycle) (e.g., Wilson, 1966; Dewey and Burke, 1974). This process is illustrated beautifully by the Neogene to Holocene development of southern

California (e.g., Crowell, 1974a, 1974b; Schneider et al., 1996; Ingersoll and Rumelhart, 1999; Kellogg and Minor, 2005; Ingersoll, 2008b).

Basins related to strike-slip faults can be classified into end-member types, although most basins are hybrids. Transtensional (including pull-apart) basins form near releasing bends and transpressional basins form at constraining bends (Crowell, 1974b). Basins associated with crustal rotations about vertical axes within the rotating blocks (“transrotational”; Ingersoll, 1988) may experience any combination of extension, compression, and strike slip (Ingersoll and Busby, 1995).

Christie-Blick and Biddle (1985) and Nilsen and Sylvester (1995) reviewed structural and stratigraphic development of strike-slip basins, based, in large part, on the pioneering work of Crowell (1974a, 1974b). They illustrated structural complexity along strike-slip faults, and implications for associated basins. Primary controls on structural patterns are the (1) degree of convergence and divergence of adjacent blocks (2) magnitude of displacement (3) material properties of deformed rocks, and (4) preexisting structures (Christie-Blick and Biddle, 1985). Subsidence in sedimentary basins results from crustal attenuation, thermal subsidence during and following extension, flexural loading due to compression, and sedimentary loading. Thermal subsidence is faster, but less in total magnitude in narrow transtensional basins than in elongate orthogonal rifts due to lateral heat conduction in the former. Distinctive aspects of sedimentary basins associated with strike-slip faults include (Christie-Blick and Biddle, 1985) (1) mismatches across basin margins (2) longitudinal and lateral basin asymmetry (3) episodic rapid subsidence (4) abrupt lateral facies changes and local unconformities, and (5) marked contrasts in stratigraphy, facies geometry, and unconformities among different basins in the same region.

These characteristics of strike-slip systems have been documented by studies of both modern and ancient, and onshore and offshore fault systems (e.g., Barnes et al., 2001, 2005; Koukouvelas and Aydin, 2002; Hsiao et al., 2004; Okay et al., 2004; Seeber et al., 2004; Wakabayashi et al., 2004).

Transtensional basins

Transtensional basins (Fig. 1.5a) form at left-stepping sinistral fault junctures and at right-stepping dextral fault junctures (Crowell, 1974a, 1974b; Reading 1980; Christie-Blick and Biddle,

1985; Nilsen and Sylvester, 1995). Mann et al. (1983) proposed a model for such basins based on a comparative study of pull-apart basins at various stages of development. Pull-apart basins evolve through the following stages: (1) nucleation of extensional faulting at releasing bends of master faults; (2) formation of spindle-shaped basins defined and commonly bisected by oblique-slip faults; (3) further extension, producing “lazy-S” or “lazy-Z” basins; (4) development into rhomboclasts, commonly with two or more sub-circular deeps; and (5) continued extension, resulting in the formation of oceanic crust at short spreading centers offset by long transforms. Basaltic volcanism and intrusion may be important during stages 3 through 5 (e.g., Crowell, 1974b). Most pull-apart basins have low length-to-width ratios, due to their short histories in changing strike-slip regimes (Mann et al., 1983). Mann (1997) demonstrated how large transtensional basins commonly form in zones of tectonic escape. Long-lived transtensional plate margins may evolve into transtensional nascent ocean basins (e.g., Gulf of California) or intraplate transform continental margins (e.g., south coast of West Africa).

Physical analog modeling provides important insights concerning initiation and evolution of pull-apart basins (e.g., Dooley and McClay, 1997; Rahe et al., 1998). Integrated geophysical and geologic studies have been conducted on several young transtensional basins and fault zones in both sub-aerial and submarine environments, for example: Hope fault of New Zealand (Wood et al., 1994), Dead Sea transform (Katzman et al., 1995; Hurwitz et al., 2002; Lazar et al., 2006), and North Anatolian fault in Sea of Marmara (Okay et al., 1999; Rangin et al., 2004). Dorsey et al. (1995) discussed the effects of rapid fault-controlled subsidence on fan-delta sedimentation along the margin of the transtensional Gulf of California. Waldron (2004) demonstrated that the Middle Pennsylvanian Stellarton pull-apart basin of Nova Scotia had a complex history of multiple overprinted structures. The overall structural and stratigraphic development of this ancient transtensional feature is consistent with the models of Mann et al. (1983), Dooley and McClay (1997), and Rahe et al. (1998).

Transpressional basins

Transpressional basins (Fig. 1.5b) include two types: 1. severely deformed and overthrust margins along sharp restraining bends that result in flexural




Fig. 1.5. True-scale actualistic analog models for sedimentary basins in transform and miscellaneous settings. These cross sections are based on the Miocene to Holocene evolution of southern California (Ingersoll, 2008b); many other tectonic settings are common for transtensional and transpressional basins. Transrotational basins are less common; the three-plate interactions depicted in (C) are unique on Earth today. Successor basins may form following cessation of any tectonic activity; shown in (C) is the southern Basin and Range of the USA, where undeformed Neogene to Holocene strata overlie tilted fault blocks and strata of the older extensional regime. Symbols same as in Figure 1.3.

subsidence due to tectonic load (e.g., northern Los Angeles basin, southern California; Schneider et al., 1996); and 2. fault-wedge basins at gentle restraining bends that result in rapid uplift of one or two margins and rapid subsidence of a basin as one block moves past the restraining bend (e.g., Neogene Ridge basin, southern California) (Crowell, 1974b, 2003a, 2003b). A basin model for type 1 would involve flexural loading similar to the foreland models discussed above, although at smaller scale.

Ridge basin is one of the most elegantly exposed and carefully studied transpressional basins in the world, as summarized by Crowell and Link (1982) and Crowell (2003a). Crowell (2003b) presented a dynamic model for the evolution of Ridge basin (12–5 Ma), a narrow crustal sliver caught between the San Gabriel fault to the southwest, and northwest-trending faults that became active sequentially in a northeast direction on the northeast side of the basin. Ridge basin became inactive when motion was transferred completely to the modern San Andreas fault (Crowell and Link, 1982; Ingersoll and Rumelhart, 1999; Crowell, 2003b). As a result of movement on the San Gabriel fault, the southwest side of the basin was uplifted and the Violin Breccia was deposited

along the basin margin. The depressed floor of the basin moved past this uplifted margin, while concurrently receiving abundant sediment from the northeast. Older depocenters moved southeastward past the restraining bend, after receiving sediment in conveyor-belt fashion, with uplift and tilting following deposition. The result is a stratigraphic thickness of over 11 km in outcrop, although vertical thickness of the basin fill is approximately one third of this. Many extraordinarily thick coarse clastic units in ancient, narrow fault-bounded basins likely were deposited in similar settings. (See Ingersoll and Busby, 1995, for a discussion of May et al.'s [1993] rejected transtensional model for the development of Ridge basin.)

McClay and Bonora (2001) developed analog models for restraining stepovers. Several studies of young transpressional features have been completed in both submarine and subaerial settings, for example the Alpine fault, New Zealand (Norris and Cooper, 1995; Barnes et al., 2005); Kobe and northern Osaka basins, Japan (Itoh et al., 2000), Cibao basin, Hispaniola (Erikson et al., 1998); Maturin foreland basin, Venezuela (Jacome et al., 2003); and southern Falkland basin (Bry et al., 2004). Trop et al. (2004) documented a transpressional origin for the Oligocene Colorado Creek basin

along the Denali fault system of Alaska. Meng et al. (2005) demonstrated how the northwest Sichuan basin (south China) evolved from a proforeland basin into a transpressional basin during the Mesozoic.

Transrotational basins

Paleomagnetic data from southern California document extensive clockwise rotation of several crustal blocks (more than 90 degrees locally), beginning in the Miocene and continuing today (e.g., Luyendyk et al., 1980; Hornafius et al., 1986; Luyendyk, 1991). Luyendyk and Hornafius (1987) developed their geometric model in order to make testable predictions concerning amount and direction of slip on faults bounding rotated and non-rotated blocks, and areas of gaps (basins) and overlap (overthrusts) among blocks. Dickinson (1996) quantified the amount of cumulative slip along the San Andreas transform fault system that is contributed by transrotational tectonism in southern California. Recognition of this contribution helps resolve discrepancies between Pacific-North American plate motions, and demonstrated offset along and within the North American continental margin (Dickinson and Wernicke, 1997).

Nicholson et al. (1994) developed a model of microplate capture that explains how complex interactions among the North American, Pacific and Farallon plates, starting soon after 30 Ma (Atwater, 1970, 1989; Bohannon and Parsons, 1995) led to three distinct phases of transfer of sections of coastal southern California onto the Pacific plate. The first phase (18–12 Ma) resulted in rapid clockwise vertical-axis rotation, with accumulation of the Topanga Formation in complex supradetachment basins (Ingersoll and Rumelhart, 1999; Ingersoll, 2008b) (Fig. 1.5c). Crouch and Suppe (1993) proposed that large-magnitude, core-complex-style extension formed in the wake of the rotating western Transverse Ranges. The southern California borderland and Los Angeles basin are floored by the Catalina Schist, interpreted by Crouch and Suppe (1993) as a footwall metamorphic tectonite, tectonically denuded below a detachment.

A model that successfully explains the extraordinarily complex basins of the Los Angeles area will need to integrate the transrotational models of Luyendyk and Hornafius (1987) and Dickinson (1996), the detachment model of Crouch and

Suppe (1993), the microplate-capture model of Nicholson et al. (1994), and the detailed stratigraphic, sedimentologic and structural history of the Los Angeles and related basins (e.g., Wright, 1991; Ingersoll and Rumelhart, 1999; Ingersoll, 2008b).

MISCELLANEOUS AND HYBRID SETTINGS

Aulacogens

During continental rifting, three rifts commonly form at approximately 120 degrees, probably because this is a least-work configuration (Burke and Dewey, 1973). Regardless of whether initiating processes are “active” or “passive” (i.e., Sengor and Burke, 1978; Morgan and Baker, 1983), in the majority of cases, two rift arms proceed through the stages of continental separation, whereas sea-floor spreading fails to develop in the third arm, resulting in a fossil rift (Sengor, 1995). Hoffman et al. (1974) discussed resulting sedimentary basins, with emphasis on a Proterozoic example. They outlined five developmental stages of the Athapuscow aulacogen, which with slight modification, provide a model applicable to most aulacogens (linear sedimentary troughs at high angles to orogens) (Fig. 1.6a): the (1) rift stage (2) transitional stage (3) downwarping stage (4) reactivation stage, and (5) postorogenic stage.

Sengor (1995) demonstrated the diverse ways in which fossil rifts (precursors to aulacogens) form, including doming, rifting, and drifting (Hoffman et al., 1974), membrane stresses, rift-tip abandonment, and continental rotation. All of these processes can result in “a narrow, elongate and fairly straight depression trending into a craton commonly from a reentrant adjoining a major basin” (Shatsky, 1964, as quoted in Sengor, 1995, 78).

Rifts that evolve into ocean basins generally are overlain by nascent-ocean and shelf-slope-rise continental margins (Fig. 1.3c–d), whereas fossil rifts adjoining continental margins evolve into reentrants that capture major drainages of continental interiors; major deltas that form at these reentrants commonly construct continental embankments (e.g., Niger Delta) (Dickinson, 1974b; Ingersoll, 1988; Ingersoll and Busby, 1995) (Fig. 1.3f). Upon activation or collision of a continental margin, the rifted-margin sedimentary prisms are intensely deformed, especially at continental promontories (Dewey and Burke, 1974; Graham et al., 1975). As orogeny




Fig. 1.6. True-scale actualistic analog models for sedimentary basins in continental collisional settings, resulting in hybrid basins. Depiction of aulacogens and impactogens in these cross sections does not show key four-dimensional relations that dictate their histories (see text for discussion). Aulacogens commonly are associated with continental embankments deposited by deltas at the mouths of fossil rifts along intraplate continental margins; therefore, thicker precollisional strata are indicated in (A). Normal faults of the fossil rifts are commonly reactivated during the aulacogen stage (syncollisional). Impactogens do not have precollisional strata or structures to reactivate. The impactogen on the right side of (B) is similar to the proximal proforeland Rhine graben. The collisional broken foreland on the left might include a distal retroforeland impactogen such as the Baikal rift. Symbols same as in Figure 1.3.

proceeds, fossil rifts become aulacogens, which may experience compressional, extensional or translational deformation.

Sengor et al. (1978) and Sengor (1995) developed criteria for distinguishing fossil rifts formed during the opening of nearby oceans that are later closed (aulacogens) from rifts formed due to crustal collision (impactogens). Both types of rift valleys trend at high angles to orogenic belts; however, aulacogens have a rifting history coincident with initiation of a neighboring ocean basin prior to collision, whereas impactogens have no precollisional rift history. Tests for distinguishing them must come from the stratigraphic record because temporal correlation of initial rifting (or lack thereof) is the primary test for their geodynamic origin (Ingersoll and Busby, 1995). Aulacogens tend to form at reentrants along rifted continental margins (Dewey and Burke, 1974), whereas impactogens are more likely to form opposite coastal promontories, where deformation of colliding continents is more intense (Sengor, 1976, 1995). This criterion must be applied cautiously, however, due

to the difficulty of definitively reconstructing precollision geometry (e.g., Thomas, 1983, 1985).

Impactogens

Impactogens (Sengor et al., 1978; Sengor, 1995) resemble aulacogens (rifts at high angles to orogenic belts), but without preorogenic stages (Fig. 1.6b). They typically form during attempted subduction of continental crust (during collision, with either another continent or a magmatic arc). Two excellent examples, of contrasting style and tectonic setting, are the middle Cenozoic Rhine graben and the late Cenozoic Baikal rift. The Rhine graben formed as a transtensional impactogen proximal to the Alpine collision orogen (Sengor, 1976). It formed on the subducting plate (Europe), in a proforeland setting. The Baikal rift, which is still active, is also transtensional, but it is distal to the related Himalayan collision (Ingersoll and Busby, 1995). It is part of the collisional broken foreland of central Asia, which is the overriding plate (Fig. 1.6b). Thus, these are end-member types

of impactogens: the Rhine graben formed in a proximal proforeland, whereas the Baikal rift formed in a distal retroforeland. Sengor (1995) discussed these and other examples.

Collisional broken-foreland basins

The collision of continents of varying shapes and sizes usually results in extreme complexity in ancient orogenic belts and related sedimentary basins (e.g., Dewey and Burke, 1974; Graham et al., 1975, 1993; Molnar and Tapponnier, 1975; Sengor, 1976, 1995; Tapponnier et al., 1982). As Tapponnier et al. (1982) demonstrated through the use of plasticine models, the collision of India and Asia has resulted in major intracontinental strike-slip faults, with associated foreland, rift, transtensional, transpressional, and transrotational basins (e.g., Graham et al., 1993; Allen et al., 1999; Yin and Harrison, 2000; Howard et al., 2003). All of these types of basins may form in either proforeland (e.g., Rhine graben) or retroforeland (e.g., Baikal rift) collisional settings.

An excellent ancient example of collisional broken-foreland basins and uplifts is the Pennsylvanian-Permian Ancestral Rocky Mountain (ARM) orogenic belt (e.g., Kluth and Coney, 1981; Kluth, 1986; Dickinson and Lawton, 2003; Blakey, 2008). ARM deformation occurred concurrently with final suturing between Laurasia and Gondwana during the late Carboniferous into the Permian (Graham et al., 1975; Kluth and Coney, 1981; Kluth, 1986; Ingersoll et al., 1995, 2003; Dickinson and Lawton, 2003; Miall, 2008). Foreland and rift basins, commonly with transpressional and transtensional components, respectively, have been documented adjacent to basement-involved uplifts (e.g., Soreghan, 1994; Geslin, 1998; Hoy and Ridgway, 2002; Barbeau, 2003). Reactivation of basement features determined location and character of many ARM uplifts and basins, including reactivation of Proterozoic fossil rifts to form aulacogens (e.g., Sengor, 1995; Marshak et al., 2000; Dickinson and Lawton, 2003; Blakey, 2008; Miall, 2008).

Additional studies of broken forelands include late Proterozoic deformation of North America (e.g., Cannon, 1995), late Paleozoic Appalachian deformation (e.g., McBride and Nelson, 1999; Murphy et al., 1999; Root and Onasch, 1999), late Paleozoic deformation of Europe (e.g., Stollhofen and Stanistreet, 1994; Mattern, 2001; Vanbrabant et al., 2002), and Mesozoic deformation of central

Asia (e.g., Sobel, 1999; Vincent and Allen, 1999; Kao et al., 2001; Johnson, 2004; Ritts et al., 2009).

Halokinetic basins

Increased exploration of deep-marine continental margins (especially continental embankments such as the northern Gulf of Mexico) has demonstrated the importance of deformation of salt in producing ponded sedimentary basins (Fig. 1.3f) (Worrall and Snelson, 1989; Winker, 1996; Prather et al., 1998; Badalini et al., 2000; Beaubouef and Friedmann, 2000). Hudec et al. (2009) reviewed subsidence mechanisms for such “minibasins,” and suggested that they can be viewed as smaller-scale models of crustal basins. They suggested that subsidence in “minibasins” can be caused by (1) density contrasts (2) diapir shortening (3) extensional diapir fall (4) decay of salt topography (5) sedimentary topographic loading, and (6) subsalt deformation. They also discussed criteria for distinguishing these subsidence mechanisms.

Study of ancient settings, where salt has played important roles in determining kinematic response of weak sediment of contrasting densities to tectonic and gravitational forces (e.g., Giles and Lawton, 2002; Rowan et al., 2003), demonstrates the uniqueness of halokinetic structural development and formation of sedimentary basins. All basins directly related to halokinetic processes (the well-studied “mini-basins” of the Gulf of Mexico, as well as diverse other salt-related basins) are herein termed “halokinetic basins.”

Bolide basins

Discovery of an iridium anomaly at the Cretaceous-Paleogene boundary (i.e., Alvarez et al., 1980) raised awareness of the significance of the impact of extraterrestrial objects (bolides) in Earth history. Not only have large impacts resulted in major evolutionary changes (e.g., mass extinctions), but also they have produced widespread sedimentary deposits resulting from tsunamis, landslides, air fall, and other bolide-induced processes (e.g., Bourgeois et al., 1988; Alvarez et al., 1992; Smit et al., 1996; Bralower et al., 1998). The “smoking gun” to explain worldwide bolide-produced sediments at the Cretaceous-Paleogene boundary has been identified as the Chicxulub crater beneath the north coast of Yucatan, Mexico (Hildebrand et al., 1991; Pope et al., 1991). Significant sedimentary basins resulting from filling of impact craters

and related features are herein termed “bolide basins” (Fig. 1.3e).

Bolide basins are now recognized in many localities on Earth: Chicxulub, Chesapeake Bay (Shah et al., 2005; Gohn et al., 2006; Hayden et al., 2008), the North Sea (Stewart and Allen, 2002, 2005), and the Barents Sea (Tsikalas et al., 1998; Dypvik et al., 2004). Detailed stratigraphic and basin analyses have been conducted in some of these basins (e.g., Marin et al., 2001; Parnell et al., 2005; Hayden et al., 2008). Some of these basins are prolific hydrocarbon producers (e.g., Grajalas-Nishimura et al., 2000). An expanding literature concerning bolide basins demonstrates the importance of this type of sedimentary basin (e.g., Glickson and Haines, 2005; Evans et al., 2008). Stewart (2003) discussed criteria for the recognition of bolide basins.

Successor basins

The original definition of successor basins (King, 1966) as “deeply subsiding troughs with limited volcanism associated with rather narrow uplifts, and overlying deformed and intruded eugeosynclines” (Kay, 1951, 107; Eisbacher, 1974) needs modification; “deeply subsiding” and “eugeosynclines” should be replaced by “intermontane” and “terrane,” respectively (Ingersoll, 1988; Ingersoll and Busby, 1995). Within the context of plate tectonics, successor basins form primarily in intermontane settings on top of inactive foldthrust belts, suture belts, transform belts, and noncratonic fossil rifts. The presence of successor basins indicates the end of orogenic or taphrogenic activity; therefore, their ages constrain interpretations of timing of suturing, deformation, and rifting (Ingersoll and Busby, 1995). Thus, they have special significance in “terrane analysis”; they represent overlap assemblages which provide minimum ages for terrane accretion (e.g., Howell et al., 1985; Ricketts, 2008).

Little work has been published on actualistic models for such basins; Eisbacher (1974) summarized models based on work on ancient basins in the Canadian Cordillera. This dearth of work may reflect the diversity of successor basins and their tectonic settings. In a sense, all basins are successor basins because they form following some orogenic or taphrogenic event represented in the basement of the basin. In fact, one of Kay’s (1951) examples of eugeosynclines (successor basins) is the post-

Nevadan basin of central California, which is now interpreted as a forearc basin, overprinted in the Cenozoic by transform tectonics (Ingersoll, 1982; Ingersoll and Schweickert, 1986; Dickinson, 1995). Dickinson (1995) discussed examples of “sutural forearc basins,” remnants of which are found along suture zones; deposition that occurred following suturing would have been in successor basins (e.g., Ricketts, 2008). Modern use of the term “successor basin” should be restricted to post-orogenic and post-taphrogenic basins that do not fall into any other plate-tectonic framework (Ingersoll and Busby, 1995). For example, most of the southern Basin and Range Province has been tectonically inactive since the Miocene (Wernicke, 1992; Dickinson, 2006). Therefore, modern intermontane basins of this region may be considered successor basins (Ingersoll and Busby, 1995) (Fig. 1.5c).

DISCUSSION

Readers of this review might be overwhelmed by the complexity of tectonic processes controlling the evolution of sedimentary basins, and the resulting complexity of this catalog of basin types. The more we know about these processes and their consequences, the more complex become our models, and the more each basin seems unique (e.g., Dickinson, 1993). This outcome is both exhilarating and frustrating. Exhilaration results from new discoveries of both fact and insight; frustration results from the need to assimilate the overwhelming crush of new information. New models are developed each time insightful simplifications or generalizations are made. Integration of observation, modeling, and experiment is an iterative, self-adjusting process.

The ultimate goal of classifying and reviewing all types of sedimentary basins is the improvement of paleotectonic and paleogeographic reconstructions through the application of actualistic models for basin evolution. Related features, whose recognition aids paleotectonic reconstruction, include suture belts (e.g., Burke et al., 1977), magmatic arcs (e.g., Sengor et al., 1991), foldthrust belts (e.g., McClay, 1992), and metamorphic belts (e.g., Miyashiro, 1973). A skilled basin analyst needs to integrate these topics, as well as geochemistry, geophysics, petrology, paleoecology, and an array of other disciplines. In a complementary manner, workers in these other fields should draw on the insights provided by the sedimentary

record to constrain their paleotectonic reconstructions. I hope that this review encourages this process of interdisciplinary development and testing of models regarding Earth's evolution.

ACKNOWLEDGMENTS

I thank C.J. Busby, W.R. Dickinson, S.A. Graham, B.K. Horton, T.E. Jordan, K.M. Marsaglia, A.D. Miall, and A.M.C. Sengor for years of rewarding interactions regarding the subjects reviewed herein. I also thank all the authors referenced below, and I apologize if I have misrepresented any of their work or if I have missed important contributions. Attempting to stay up to date with all these topics is a risky business!

REFERENCES

- Aalto, K.R., and Miller, W., III (1999) Sedimentology of the Pliocene Upper Onzole Formation, an inner-trench slope succession in northwestern Ecuador. *Journal of South American Earth Sciences*, 12, 69–85.
- Abascal, L. del V. (2005) Combined thin-skinned and thick-skinned deformation in the central Andean foreland of northwestern Argentina. *Journal of South American Earth Sciences*, 19, 75–81.
- Alavi, M. (2004) Regional stratigraphy of the Zagros fold-thrust belt of Iran and its proforeland evolution. *American Journal of Science*, 304, 1–20.
- Alcicek, M.C. (2007) Tectonic development of an orogen-top rift recorded by its terrestrial sedimentation pattern: the Neogene Esen basin of southwestern Anatolia, Turkey. *Sedimentary Geology*, 200, 117–140.
- Allen, M.B., Vincent, S.J., and Wheeler, P.J. (1999) Late Cenozoic tectonics of the Kepingtage thrust zone: interactions of the Tien Shan and Tarim basin, northwest China. *Tectonics*, 18, 639–654.
- Allen, P.A., Homewood, P., and Williams, G.D. (1986) Foreland basins: an introduction. *International Association of Sedimentologists Special Publication*, 8, 3–12.
- Allen, P.A., Burgess, P.M., Galewsky, J., and Sinclair, H.D. (2001) Flexural-eustatic numerical model for drowning of the Eocene perialpine carbonate ramp and implications for Alpine geodynamics. *Geological Society of America Bulletin*, 113, 1052–1066.
- Allen, R., Carter, A., Najman, Y., Bandopadhyay, P.C., Chapman, H.J., Bickle, M.J., Garzanti, E., Vezzoli, G., Ando, S., Foster, G.L., and Gerring, C. (2008) New constraints on the sedimentation and uplift history of the Andaman-Nicobar accretionary prism, South Andaman Island. *Geological Society of America Special Paper*, 436, 223–255.
- Alvarez, L.W., Alvarez, W., Asaro, F., and Michel, H.V. (1980) Extraterrestrial cause for the Cretaceous-Tertiary extinction. *Science*, 208, 1095–1108.
- Alvarez, W., Smit, J., Lowrie, W., Asaro, F., Margolis, S.V., Claeys, P., Kastner, M., and Hildebrand, A.R. (1992) Proximal impact deposits at the Cretaceous-Tertiary boundary in the Gulf of Mexico: a restudy of DSDP Leg 77 Sites 536 and 540. *Geology*, 20, 697–700.
- Arenas, C., Millan, H., Pardo, G., and Pocovi, A. (2001) Ebro basin continental sedimentation associated with late compressional Pyrenean tectonics (north-eastern Iberia): controls on basin margin fans and fluvial systems. *Basin Research*, 13, 65–89.
- Atwater, T. (1970) Implications of plate tectonics for the Cenozoic tectonic evolution of western North America. *Geological Society of America Bulletin*, 81, 3513–3535.
- Atwater, T. (1989) Plate tectonic history of the northeast Pacific and western North America, in Winterer, E.L., Hussong, D.M., and Decker, R.W., eds., *The eastern Pacific Ocean and Hawaii. The geology of North America*, vol. N. Boulder, CO, Geological Society of America, 21–72.
- Atwater, T., and Stock, J. (1998) Pacific-North America plate tectonics of the Neogene southwestern United States: an update. *International Geology Review*, 40, 375–402.
- Audley-Charles, M.G., Curray, J.R., and Evans, G. (1977) Location of major deltas. *Geology*, 5, 341–344.
- Axen, G.J., and Fletcher, J.M. (1998) Late Miocene-Pleistocene extensional faulting, northern Gulf of California, Mexico and Salton Trough, California. *International Geology Review*, 40, 217–244.
- Badalini, G., Kneller, B., and Winker, C.D. (2000) Architecture and process in the late Pleistocene Brazos-Trinity turbidite system, Gulf of Mexico continental slope, in Weimer P., Slatt R.M., Coleman J., Rossen N.C., Nelson H., Bouma A.H., Styzen M.J., Lawrence D.T., eds., *Deep-water reservoirs of the world, 20th Annual Research Conference Proceedings. Society of Economic Paleontologist and Mineralogists, Gulf Coast Section, Houston, TX*, 16–34.
- Bally, A.W. (1989) Phanerozoic basins of North America, in Bally, A.W., and Palmer, A.R., eds., *The geology of North America: an overview. The Geology of North America*, A. Boulder, CO, Geological Society of America, 397–447.
- Barbeau, D.L. (2003) A flexural model for the Paradox basin: implications for the tectonics of the Ancestral Rocky Mountains. *Basin Research*, 15, 97–115.
- Barnes, P.M., Sutherland, R., Davy, B., and Delteil, J. (2001) Rapid creation and destruction of sedimentary basins on mature strike-slip faults: an example from the offshore Alpine fault, New Zealand. *Journal of Structural Geology*, 23, 1727–1739.
- Barnes, P.M., Sutherland, R., and Delteil, J. (2005) Strike-slip structure and sedimentary basins of the southern Alpine fault, Fiordland, New Zealand. *Geological Society of America Bulletin*, 117, 411–435.
- Beaubouef, R.T., and Friedmann, S.J. (2000) High resolution seismic/sequence stratigraphic framework for the evolution of Pleistocene intraslope basins, western Gulf of Mexico: Depositional models and reservoir analogs, in Weimer P., Slatt R.M., Coleman J., Rossen N.C., Nelson H., Bouma A.H., Styzen M.J., Lawrence D.T., eds., *Deep-water reservoirs of the world, 20th Annual Research Conference Proceedings. Society of Economic Paleontologist and Mineralogists, Gulf Coast Section, Houston, TX*, 40–60.
- Beaumont, C. (1981) Foreland basins. *Geophysical Journal of the Royal Astronomical Society*, 65, 291–329.

- Beaumont, C., Ellis, S., Hamilton, J., and Fullsack, P. (1996) Mechanical model for subduction-collision tectonics of Alpine-type compressional orogens. *Geology*, 24, 675–678.
- Beck, M. (1983) On the mechanism of tectonic transport in zones of oblique subduction. *Tectonophysics*, 93, 1–11.
- Ben-Avraham, Z., and Uyeda, S. (1983) Entrapment origin of marginal seas. *American Geophysical Union Geodynamics Series*, 11, 91–104.
- Berger, W.H. (1973) Cenozoic sedimentation in the eastern tropical Pacific. *Geological Society of America Bulletin*, 84, 1941–1954.
- Berger, W.H., and Winterer, E.L. (1974) Plate stratigraphy and the fluctuating carbonate line. *International Association of Sedimentologists Special Publication*, 1, 11–48.
- Berry, J.P., and Wilkinson, B.H. (1994) Paleoclimatic and tectonic control on the accumulation of North American cratonic sediment. *Geological Society of America Bulletin*, 106, 855–865.
- Bertotti, G., Picotti, V., and Cloetingh, S. (1998) Lithospheric weakening during “retroforeland” basin formation: tectonic evolution of the central South Alpine foredeep. *Tectonics*, 17, 131–142.
- Bertotti, G., Picotti, V., Chilovi, C., Fantoni, R., Merlini, S., and Mosconi, A. (2001) Neogene to Quaternary sedimentary basins in the south Adriatic (Central Mediterranean): foredeeps and lithospheric buckling. *Tectonics*, 20, 771–787.
- Blakey, R.C. (2008) Pennsylvanian-Jurassic sedimentary basins of the Colorado Plateau and southern Rocky Mountains, in Miall, A.D., ed., *The sedimentary basins of the United States and Canada. Sedimentary Basins of the World*, vol. 5. Amsterdam, Elsevier, 245–296.
- Bohannon, R.G. (1986a) How much divergence has occurred between Africa and Arabia as a result of the opening of the Red Sea? *Geology*, 14, 510–513.
- Bohannon, R.G. (1986b) Tectonic configuration of the western Arabian continental margin, southern Red Sea. *Tectonics*, 5, 477–499.
- Bohannon, R.G., and Parsons, T. (1995) Tectonic implications of post-30 Ma Pacific and North American relative plate motions. *Geological Society of America Bulletin*, 107, 937–959.
- Bond, G.C., Kominz, M.A., and Sheridan, R.E. (1995) Continental terraces and rises, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 149–178.
- Bostrom, R.C. (1971) Westward displacement of the lithosphere. *Nature*, 234, 536–538.
- Bosworth, W., Huchon, P., and McClay, K. (2005) The Red Sea and Gulf of Aden basins. *Journal of African Earth Sciences*, 43, 324–378.
- Bourgeois, J., Hansen, T.A., Wiberg, P.L., and Kauffman, E.G. (1988) A tsunami deposit at the Cretaceous-Tertiary boundary in Texas. *Science*, 241, 567–570.
- Bradley, D.C. (1989) Taconic plate kinematics as revealed by foredeep stratigraphy, Appalachian orogen. *Tectonics*, 8, 1037–1049.
- Bradley, D.C., and Kidd, W.S.F. (1991) Flexural extension of the upper continental crust in collisional foredeeps. *Geological Society of America Bulletin*, 103, 1416–1438.
- Bralower, T.J., Paull, C.K., and Leckie, R.M. (1998) The Cretaceous-Tertiary boundary in Busby cocktail: Chicxulub impact triggers margin collapse and extensive sediment gravity flows. *Geology*, 26, 331–334.
- Brunet, M-F., Korotaev, M.V., Ershov, A.V., and Nikishin, A. M. (2003) The South Caspian basin: a review of its evolution from subsidence modeling. *Sedimentary Geology*, 156, 119–148.
- Bry, M., White, N., Singh, S., England, R., and Trowell, C. (2004) Anatomy and formation of oblique continental collision: South Falkland basin. *Tectonics*, 23, TC4011, 20 p.
- Buffler, R.T., and Thomas, W.A. (1994) Crustal structure and evolution of the southeastern margin of North America and the Gulf of Mexico basin, in Speed, R.C., ed., *Phanerozoic evolution of North American continent-ocean transitions*. Geological Society of America, Decade of North American Geology Summary Volume to accompany the DNAG Continent-Ocean Transects Series, 219–264.
- Burbank, D.W., and Tahirkheli, R.A.K. (1985) The magnetostratigraphy, fission-track dating, and stratigraphic evolution of the Peshawar intermontane basin, northern Pakistan. *Geological Society of America Bulletin*, 96, 539–552.
- Burchfiel, B.C., Chen, Z., Hodges, K.V., Liu, Y., Royden, L.H., Deng, C., and Xu, J. (1992) The south Tibetan detachment system, Himalayan orogen: Extension contemporaneous with and parallel to shortening in a collisional mountain belt. *Geological Society of America Special Paper*, 269, 41 p.
- Burgess, P.M. (2008) Phanerozoic evolution of the sedimentary cover of the North American craton, in Miall, A.D., ed., *The sedimentary basins of the United States and Canada. Sedimentary basins of the world*, vol. 5. Amsterdam, Elsevier, 31–63.
- Burgess, P.M., and Gurnis, M. (1995) Mechanisms for the formation of cratonic stratigraphic sequences. *Earth and Planetary Science Letters*, 136, 647–663.
- Burgess, P.M., Gurnis, M., and Moresi, L. (1997) Formation of sequences in the cratonic interior of North America by interaction between mantle, eustatic, and stratigraphic processes. *Geological Society of America Bulletin*, 109, 1515–1535.
- Burke, K. (1972) Longshore drift, submarine canyons, and submarine fans in development of Niger delta. *American Association of Petroleum Geologists Bulletin*, 56, 1975–1983.
- Burke, K., and Dewey, J.F. (1973) Plume-generated triple junctions: key indicators in applying plate tectonics to old rocks. *Journal of Geology*, 81, 406–433.
- Burke, K., Dewey, J.F., and Kidd, W.S.F. (1977) World distribution of sutures: the sites of former oceans. *Tectonophysics*, 40, 69–99.
- Busby, C.J., Adams, B.F., Mattinson, J., and De Oreo, S. (2006) View of an intact oceanic arc, from surficial to mesozonal levels: Cretaceous Alisitos arc, Baja California, Mexico. *Journal of Volcanology and Geothermal Research*, 149, 1–46.
- Busby-Spera, C.J. (1987) Lithofacies of deep marine basalts emplaced on a Jurassic backarc apron, Baja California (Mexico). *Journal of Geology*, 95, 671–686.
- Busby-Spera, C.J. (1988a) Evolution of a Middle Jurassic back-arc basin, Cedros Island, Baja California: evidence from a marine volcanoclastic apron. *Geological Society of America Bulletin*, 100, 218–233.

- Busby-Spera, C.J. (1988b) Speculative tectonic model for the early Mesozoic arc of the southwest Cordilleran United States. *Geology*, 16, 1121–1125.
- Busby-Spera, C.J., Mattinson, J.M., Riggs, N.R., and Schermer, E.R. (1990) The Triassic-Jurassic magmatic arc in the Mojave-Sonoran Deserts and the Sierran-Klamath region: similarities and differences in paleogeographic evolution. *Geological Society of America Special Paper*, 255, 325–338.
- Cameron, T.D.J., Crosby, A., Balson, P.S., Jeffrey, D.H., Lott, G.K., Bulat, J., and Harrison, D.J. (1992) United Kingdom offshore regional report: the geology of the southern North Sea. HMSO for the British Geological Survey, London, 152 p.
- Cannon, W.F. (1995) Closing of the Midcontinent rift: a far-field effect of Grenvillian compression. *Geology*, 22, 155–158.
- Cardozo, N., and Jordan, T. (2001) Causes of spatially variable tectonic subsidence in the Miocene Bermejo foreland basin, Argentina. *Basin Research*, 13, 335–357.
- Carrapa, B. (2009) Tracing exhumation and orogenic wedge dynamics in the European Alps with detrital thermochronology. *Geology*, 37, 1127–1130.
- Carrapa, B., and Garcia-Castellanos, D. (2005) Western Alpine back-thrusting as subsidence mechanism in the Tertiary Piedmont basin (western Po Plain, NW Italy). *Tectonophysics*, 406, 197–212.
- Cas, R., and Busby-Spera, C., eds. (1991) Volcaniclastic sedimentation. *Sedimentary Geology*, 74, 362 p.
- Cas, R.A.F., and Wright, J.V. (1987) Volcanic successions modern and ancient. Boston, Allen and Unwin, 528 p.
- Casas-Sainz, A.M., Cortes-Garcia, A.L., and Maestro-Gonzalez, A. (2000) Intraplate deformation and basin formation during the Tertiary within the northern Iberian plate: origin and evolution of the Almazan basin. *Tectonics*, 19, 258–289.
- Castle, J.W. (2001) Appalachian basin stratigraphic response to convergent-margin structural evolution. *Basin Research*, 13, 397–418.
- Cather, S.M., and Chapin, C.E. (1990) Paleogeographic and paleotectonic setting of Laramide sedimentary basins in the central Rocky Mountain region: alternative interpretation. *Geological Society of America Bulletin*, 102, 256–260.
- Catuneanu, O. (2004) Retroarc foreland systems: evolution through time. *Journal of African Earth Sciences*, 38, 225–242.
- Chapin, C.E., and Cather, S.M. (1981) Eocene tectonics and sedimentation in the Colorado Plateau-Rocky Mountain area. *Arizona Geological Society Digest*, 14, 173–198.
- Chen, W.-S., Ridgway, K.D., Horng, C.-S., Chen, Y.-G., Shea, K.-S., and Yeh, M.-G. (2001) Stratigraphic architecture, magnetostratigraphy, and incised-valley systems of the Pliocene-Pleistocene collisional marine foreland basin of Taiwan. *Geological Society of America Bulletin*, 113, 1249–1271.
- Chiang, C.-S., Yu, H.-S., and Chou, Y.-W. (2004) Characteristics of the wedge-top depozone of the southern Taiwan foreland basin system. *Basin Research*, 16, 65–78.
- Christie-Blick, N., and Biddle, K.T. (1985) Deformation and basin formation along strike-slip faults. *Society of Economic Paleontologists and Mineralogists Special Publication*, 37, 1–34.
- Clague, D.A. (1981) Linear island and seamount chains, aseismic ridges and intraplate volcanism: results from DSDP. *Society of Economic Paleontologists and Mineralogists Special Publication*, 32, 7–22.
- Clift, P.D., Draut, A.E., Kelemen, P.B., Blusztajn, J., and Greene, A. (2005) Stratigraphic and geochemical evolution of an oceanic arc upper crustal section: the Jurassic Talkeetna Volcanic Formation, south-central Alaska. *Geological Society of America Bulletin*, 117, 902–925.
- Cloetingh, S. (1988) Intraplate stress: a new element in basin analysis, in Kleinspehn, K.L., and Paola, C., eds., *New perspectives in basin analysis*. New York, Springer-Verlag, 205–230.
- Cloos, M. (1993) Lithospheric buoyancy and collisional orogenesis: subduction of oceanic plateaus, continental margins, island arcs, spreading ridges, and seamounts. *Geological Society of America Bulletin*, 105, 715–737.
- Cloos, M., and Shreve, R.L. (1988a) Subduction-channel model of prism accretion, melange formation, sediment subduction, and subduction erosion at convergent plate margins: 1. background and description. *Pure and Applied Geophysics*, 128, 455–500.
- Cloos, M., and Shreve, R.L. (1988b) Subduction-channel model of prism accretion, melange formation, sediment subduction, and subduction erosion at convergent plate margins: 2. implications and discussion. *Pure and Applied Geophysics*, 128, 501–545.
- Cochran, J.R. (1983) A model for development of Red Sea. *American Association of Petroleum Geologists Bulletin*, 67, 41–69.
- Cogne, J.P., Humler, E., and Courtillot, V. (2006) Mean age of oceanic lithosphere drives eustatic sea-level change since Pangea breakup. *Earth and Planetary Science Letters*, 245, 115–122.
- Coleman, R.G. (1993) *Geologic evolution of the Red Sea*. New York, Oxford University Press, 186 p.
- Constenius, K.N., Johnson, R.A., Dickinson, W.R., and Williams, T.A. (2000) Tectonic evolution of the Jurassic-Cretaceous Great Valley forearc, California: implications for the Franciscan thrust-wedge hypothesis. *Geological Society of America Bulletin*, 112, 1703–1723.
- Cook, H.E. (1975) North American stratigraphic principles as applied to deep-sea sediments. *American Association of Petroleum Geologists Bulletin*, 59, 817–837.
- Cooper, M.A., and Williams, G.D., eds. (1989) *Inversion tectonics*. Geological Society of London Special Publication, 44, 375 p.
- Critelli, S., and Le Pera, E. (1994) Detrital modes and provenance of Miocene sandstones and modern sands of the southern Apennines thrust-top basins (Italy). *Journal of Sedimentary Research*, A64, 824–835.
- Critelli, S., Marsaglia, K.M., and Busby, C.J. (2002) Tectonic history of a Jurassic backarc-basin sequence (the Gran Canon Formation, Cedros Island, Mexico) based on compositional modes of tuffaceous deposits. *Geological Society of America Bulletin*, 114, 515–527.
- Crouch, J.K., and Suppe, J. (1993) Late Cenozoic tectonic evolution of the Los Angeles basin and inner California borderland: a model for core complex-like crustal extension. *Geological Society of America Bulletin*, 105, 1415–1434.

- Crowell, J.C. (1974a) Sedimentation along the San Andreas fault, California. Society of Economic Paleontologists and Mineralogists Special Publication, 19, 292–303.
- Crowell, J.C. (1974b) Origin of late Cenozoic basins in southern California. Society of Economic Paleontologists and Mineralogists Special Publication, 22, 190–204.
- Crowell, J.C., ed. (2003a) Evolution of Ridge basin, southern California: an interplay of sedimentation and tectonics. Geological Society of America Special Paper, 367, 247 p.
- Crowell, J.C. (2003b) Tectonics of Ridge basin, southern California, in Crowell, J.C., ed., Evolution of Ridge basin, southern California: an interplay of sedimentation and tectonics. Geological Society of America Special Paper, 367, 157–203.
- Crowell, J.C. and Link, M.H., eds. (1982) Geologic history of Ridge basin southern California. Book 22. Pacific Section, Society of Economic Paleontologists and Mineralogists, Los Angeles, 304 p.
- Cruciani, C., Carminati, E., and Doglioni, C. (2005) Slab dip v. lithosphere age: no direct function. *Earth and Planetary Science Letters*, 238, 298–310.
- Davila, F.M., and Astini, R.A. (2003) Early Middle Miocene broken foreland development in the southern Central Andes: evidence for extension prior to regional shortening. *Basin Research*, 15, 379–396.
- DeCelles, P.G., and Giles, K.A. (1996) Foreland basin systems. *Basin Research*, 8, 105–123.
- DeCelles, P.G., and Giles, K.A. (1997) Foreland basin systems: Reply. *Basin Research*, 9, 171–176.
- DeCelles, P.G., and Horton, B.K. (2003) Early to middle Tertiary foreland basin development and the history of Andean crustal shortening in Bolivia. *Geological Society of America Bulletin*, 115, 58–77.
- DeCelles, P.G., Gehrels, G.E., Quade, J., and Ojha, T.P. (1998) Eocene-early Miocene foreland basin development and the history of Himalayan thrusting, western and central Nepal. *Tectonics*, 17, 741–765.
- DeCelles, P.G., Robinson, D.M., Quade, J., Ojha, T.P., Garzione, C.N., Copeland, P., and Upreti, B.N. (2001) Stratigraphy, structure, and tectonic evolution of the Himalayan fold-thrust belt in western Nepal. *Tectonics*, 20, 487–509.
- DeCelles, P.G., Quade, J., Kapp, P., Fan, M.J., Dettman, D.L., and Ding, L. (2007) High and dry in central Tibet during the Late Oligocene. *Earth and Planetary Science Letters*, 253, 389–401.
- DeRito, R.F., Cozzarelli, F.A. and Hodge, D.S. (1983) Mechanism of subsidence of ancient cratonic rift basins. *Tectonophysics*, 94, 141–168.
- Dewey, J.F. (1980) Episodicity, sequence and style at convergent plate boundaries. *Geological Association of Canada Special Paper*, 20, 553–573.
- Dewey, J.F., and Burke, K. (1974) Hot spots and continental break-up: implications for collisional orogeny. *Geology*, 2, 57–60.
- Dickinson, W.R. (1973) Widths of modern arc-trench gaps proportional to past duration of igneous activity in associated magmatic arcs. *Journal of Geophysical Research*, 78, 3376–3389.
- Dickinson, W.R. (1974a) Sedimentation within and beside ancient and modern magmatic arcs. Society of Economic Paleontologists and Mineralogists Special Publication, 19, 230–239.
- Dickinson, W.R. (1974b) Plate tectonics and sedimentation. Society of Economic Paleontologists and Mineralogists Special Publication, 22, 1–27.
- Dickinson, W.R. (1976a) Plate tectonic evolution of sedimentary basins. American Association of Petroleum Geologists Continuing Education Course Notes Series 1, 62 p.
- Dickinson, W.R. (1976b) Sedimentary basins developed during evolution of Mesozoic-Cenozoic arc-trench system in western North America. *Canadian Journal of Earth Sciences*, 13, 1268–1287.
- Dickinson, W.R. (1978) Plate tectonic evolution of north Pacific rim. *Journal of the Physics of the Earth*, 26, Supplement, S1–S19.
- Dickinson, W.R. (1981a) Plate tectonic evolution of the southern Cordillera. *Arizona Geological Society Digest*, 14, 113–135.
- Dickinson, W.R. (1981b) Plate tectonics and the continental margin of California, in Ernst, W.G., ed., *The geotectonic development of California*. Rubey, vol. 1. Englewood Cliffs, NJ, Prentice Hall, 1–28.
- Dickinson, W.R. (1990) Paleogeographic and paleotectonic setting of Laramide sedimentary basins in the central Rocky Mountain region: reply. *Geological Society of America Bulletin*, 102, 281–282.
- Dickinson, W.R. (1993) Basin geodynamics. *Basin Research*, 5, 195–196.
- Dickinson, W.R. (1995) Forearc basins, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 221–261.
- Dickinson, W.R. (1996) Kinematics of transrotational tectonism in the California Transverse Ranges and its contribution to cumulative slip along the San Andrea transform fault system. *Geological Society of America Special Paper*, 305, 46 p.
- Dickinson, W.R. (2004) Impacts of eustasy and hydro-isostasy on the evolution and landforms of Pacific atolls. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 213, 251–269.
- Dickinson, W.R. (2006) Geotectonic evolution of the Great Basin. *Geosphere*, 7, 353–368.
- Dickinson, W.R. (2008) Tectonic lessons from the configuration and internal anatomy of the Circum-Pacific orogenic belt, in Spencer, J.E., and Titley, S.R., eds., *Ores and orogenesis: Circum-Pacific tectonics, geologic evolution, and ore deposits*. Arizona Geological Society Digest 22, 5–18.
- Dickinson, W.R., and Lawton, T.F. (2001) Carboniferous to Cretaceous assembly and fragmentation of Mexico. *Geological Society of America Bulletin*, 113, 1142–1160.
- Dickinson, W.R., and Lawton, T.F. (2003) Sequential intercontinental suturing as the ultimate control for Pennsylvanian Ancestral Rocky Mountains deformation. *Geology*, 31, 609–612.
- Dickinson, W.R., and Seely, D.R. (1979) Structure and stratigraphy of forearc regions. *American Association of Petroleum Geologists Bulletin*, 63, 2–31.
- Dickinson, W.R., and Snyder, W.S. (1978) Plate tectonics of the Laramide orogeny. *Geological Society of America Memoir*, 151, 355–366.

- Dickinson, W.R., and Wernicke, B.P. (1997) Reconciliation of San Andreas slip discrepancy by a combination of interior Basin and Range extension and transrotation near the coast. *Geology*, 25, 663–665.
- Dickinson, W.R., Harbaugh, D.W., Saller, A.H., Heller, P.L., and Snyder, W.S. (1983) Detrital modes of upper Paleozoic sandstones derived from Antler orogen in Nevada: implications for nature of Antler orogeny. *American Journal of Science*, 283, 481–509.
- Dickinson, W.R., Klute, M.A., Hayes, M.J., Janecke, S.U., Lundin, E.R., McKittrick, M.A., and Olivares, M.D. (1988) Paleogeographic and paleotectonic setting of Laramide sedimentary basins in the central Rocky Mountain region. *Geological Society of America Bulletin*, 100, 1023–1039.
- Dickinson, W.R., Klute, M.A., Hayes, M.J., Janecke, S.U., Lundin, E.R., McKittrick, M.A., and Olivares, M.D. (1990) Paleogeographic and paleotectonic setting of Laramide sedimentary basins in the central Rocky Mountain region: reply. *Geological Society of America Bulletin*, 102, 259–260.
- Doglioni, C. (1994) Foredeeps versus subduction zones. *Geology*, 22, 271–274.
- Doglioni, C., Harabaglia, P., Merlini, S., Mongelli, F., Peccerillo, A., and Piromallo, C. (1999) Orogens and slabs vs. their direction of subduction. *Earth-Science Reviews*, 45, 167–208.
- Dooley, T., and McClay, K. (1997) Analog modeling of pull-apart basins. *American Association of Petroleum Geologists Bulletin*, 81, 1804–1826.
- Dorobek, S.L. (2008) Carbonate-platform facies in volcanic-arc settings: characteristics and controls on deposition and stratigraphic development. *Geological Society of America Special Paper*, 436, 55–90.
- Dorobek, S.L., and Ross, G.M., eds. (1995) Stratigraphic evolution of foreland basins. *SEPM (Society for Sedimentary Geology) Special Publication*, 52, 310 p.
- Dorsey, R.J., Umhoefer, P.J., and Renne, P.R. (1995) Rapid subsidence and stacked Gilbert-type fan deltas, Pliocene Loreto basin, Baja California Sur, Mexico. *Sedimentary Geology*, 98, 181–204.
- Dypvik, H., Sandbakken, P.T., Postma, G., and Mork, A. (2004) Early post-impact sedimentation around the central high of the Mjolnir impact crater (Barents Sea, Late Jurassic). *Sedimentary Geology*, 168, 227–247.
- Einsele, G. (1985) Basaltic sill-sediment complexes in young spreading centers: genesis and significance. *Geology*, 13, 249–252.
- Einsele, G., Liu, B., Durr, S., Frisch, W., Liu, G., Luterbacher, H.P., Ratschbacher, L., Ricken, W., Wendt, J., Wetzel, A., Yu, G., and Zheng, H. (1994) The Xigaze forearc basin: evolution and facies architecture (Cretaceous, Tibet). *Sedimentary Geology*, 90, 1–32.
- Eisbacher, G.H. (1974) Evolution of successor basins in the Canadian Cordillera. *Society of Economic Paleontologists and Mineralogists Special Publication*, 19, 274–291.
- Erikson, J.P., Pindall, J.L., Karner, G.D., Sonder, L.J., Fuller, E., and Dent, L. (1998) Neogene sedimentation and tectonics in the Cibao basin and northern Hispaniola: an example of basin evolution near a strike-slip-dominated plate boundary. *Journal of Geology*, 106, 473–494.
- Evans, K.R., Horton, J.W., Jr., King, D.T., Jr., and Morrow, J.R., eds. (2008) The sedimentary record of meteorite impacts. *Geological Society of America Special Paper*, 437, 213 p.
- Evans, M.J., and Elliott, T. (1999) Evolution of a thrust-sheet-top basin: the Tertiary Barreme basin, Alpes-de-Haute-Provence, France. *Geological Society of America Bulletin*, 111, 1617–1643.
- Fackler-Adams, B., and Busby, C.J. (1998) Structural and stratigraphic evolution of extensional oceanic arcs. *Geology*, 26, 735–738.
- Faulds, J.E., and Varga, R.J. (1998) The role of accommodation zones and transfer zones in the regional segmentation of extended terranes. *Geological Society of America Special Paper*, 323, 1–45.
- Fildani, A., Cope, T.D., Graham, S.A., and Wooden, J.L. (2003) Initiation of the Magallanes foreland basin: timing of the southernmost Patagonian Andes orogeny revised by detrital zircon provenance analysis. *Geology*, 31, 1081–1084.
- Fisher, R.V., and Schmincke, H.-U. (1984) *Pyroclastic rocks*. New York, Springer-Verlag, 472 p.
- Fisher, R.V., and Smith, G.A., eds. (1991) *Sedimentation in volcanic settings*. *SEPM (Society for Sedimentary Geology) Special Publication*, 45, 257 p.
- Ford, M. (2004) Depositional wedge tops: interaction between low basal friction external orogenic wedges and flexural foreland basins. *Basin Research*, 16, 361–375.
- Friedmann, S.J., and Burbank, D.W. (1995) Rift basins and supradetachment basins: intracontinental extensional end-members. *Basin Research*, 7, 109–127.
- Funk, J., Mann, P., McIntosh, K., and Stephens, J. (2009) Cenozoic tectonics of the Nicaraguan depression, Nicaragua, and Median Trough, El Salvador, based on seismic-reflection profiling and remote-sensing data. *Geological Society of America Bulletin*, 121, 1491–1521.
- Galewsky, J., Silver, E.A., Gallup, C.D., Edwards, R.L., and Potts, D.C. (1996) Foredeep tectonics and carbonate platform dynamics in the Huon Gulf, Papua New Guinea. *Geology*, 24, 819–822.
- Galloway, W.E., Ganey-Curry, P.E., Li, X., and Buffler, R.T. (2000) Cenozoic depositional history of the Gulf of Mexico basin. *American Association of Petroleum Geologists Bulletin*, 84, 1743–1774.
- Garzione, C.N., DeCelles, P.G., Hodkinson, D.G., Ojha, T.P., and Upreti, B.N. (2003) East-west extension and Miocene environmental change in the southern Tibetan plateau: Thakkhola graben, central Nepal. *Geological Society of America Bulletin*, 115, 3–20.
- Gawthorpe, R.L., and Leeder, M.R. (2000) Tectono-sedimentary evolution of active extensional basins. *Basin Research*, 12, 195–218.
- Geslin, J.K. (1998) Distal Ancestral Rocky Mountains tectonism: evolution of the Pennsylvanian-Permian Oquirrh-Wood River basin, southern Idaho. *Geological Society of America Bulletin*, 110, 644–663.
- Giles, K.A., and Lawton, T.F. (2002) Halokinetic sequence stratigraphy adjacent to the El Papalote diapir, north-eastern Mexico. *American Association of Petroleum Geologists Bulletin*, 86, 823–840.
- Giovanni, M.K., Horton, B.K., Garzione, C.N., McNulty, B., and Grove, M. (2010) Extensional basin evolution in the Cordillera Blanca, Peru: stratigraphic and isotopic

- records of detachment faulting and orogenic collapse in the Andean hinterland. *Tectonics*, 29, TC6007, 21 p.
- Glickson, A.Y., and Haines, P.W., eds. (2005) Thematic issue: Shoemaker memorial issue on the Australian impact record: 1997–2005 update. *Australian Journal of Earth Sciences*, 52, 475–798.
- Gohn, G.S., Koeberl, C., Miller, K.G., Reimold, W.U., and the Scientific Staff of the Chesapeake Bay Impact Structure Drilling Project (2006) Chesapeake Bay Impact Structure Deep Drilling Project completes coring. *Scientific Drilling* (3), 34–39.
- Gomez, E., Jordan, T.E., Allmendinger, R.W., and Cardozo, N. (2005) Development of the Colombian foreland-basin system as a consequence of diachronous exhumation of the northern Andes. *Geological Society of America Bulletin*, 117, 1272–1292.
- Gould, S.J. (1989) *Wonderful life: the Burgess Shale and the nature of history*. New York, W. W. Norton, 347 p.
- Graham, S.A., Dickinson, W.R., and Ingersoll, R.V. (1975) Himalayan-Bengal model for flysch dispersal in Appalachian-Ouachita system. *Geological Society of America Bulletin*, 86, 273–286.
- Graham, S.A., Hendrix, M.S., Wang, L.B., and Carroll, A.R. (1993) Collisional successor basins of western China: impact of tectonic inheritance on sand composition. *Geological Society of America Bulletin*, 105, 323–344.
- Grajales-Nishimura, J.M., Cedillo-Pardo, E., Rosales-Dominguez, C., Moran-Zenteno, D.J., Alvarez, W., Claeys, P., Ruis-Morales, J., Garcia-Hernandez, J., Padilla-Avila, P., and Sanchez-Rios, A. (2000) Chicxulub impact: the origin of reservoir and seal facies in the southeastern Mexican oil fields. *Geology*, 28, 307–310.
- Gupta, S., and Allen, P.A. (2000) Implications of foreland paleotopography for stratigraphic development in the Eocene distal Alpine foreland basin. *Geological Society of America Bulletin*, 112, 515–530.
- Gurnis, M. (1993) Depressed continental hypsometry behind oceanic trenches: a clue to subduction controls on sea-level change. *Geology*, 21, 29–32.
- Haddad, D., and Watts, A.B. (1999) Subsidence history, gravity anomalies, and flexure of the northeast Australian margin in Papua New Guinea. *Tectonics*, 18, 827–842.
- Hall, R., and Smyth, H.R. (2008) Cenozoic arc processes in Indonesia: identification of the key influences on the stratigraphic record in active volcanic arcs. *Geological Society of America Special Paper*, 436, 27–54.
- Hamilton, W. (1969) Mesozoic California and the underflow of Pacific mantle. *Geological Society of America Bulletin*, 80, 2409–2430.
- Hamilton, W. (1979) *Tectonics of the Indonesian region*. United States Geological Survey Professional Paper 1078, 345 p.
- Hansen, W.R. (1990) Paleogeographic and paleotectonic setting of Laramide sedimentary basins in the central Rocky Mountain region: alternative interpretation. *Geological Society of America Bulletin*, 102, 280–282.
- Hayden, T., Kominz, M., Powars, D.S., Edwards, L.E., Miller, K.G., Browning, J.V., and Kulpecz, A.A. (2008) Impact effects and regional tectonic insights: Backstripping the Chesapeake Bay impact structure. *Geology*, 36, 327–330.
- Heckel, P.H. (1984) Changing concepts of Midcontinent Pennsylvanian cyclothems, North America, in *Congress International de Stratigraphie et de Geologie du Carbonifere*, 9th. *Compte Rendu*, 3. Carbondale, Southern Illinois University Press, 535–538.
- Heezen, B.C., MacGregor, I.D., Foreman, H.P., Forristal, G., Hekel, H., Hesse, R., Hoskins, R.H., Jones, E.J.W., Kaneps, A., Krashennnikov, V.A., Okada, H., and Ruef, M.H. (1973) Diachronous deposits: a kinematic interpretation of the post Jurassic sedimentary sequence on the Pacific Plate. *Nature*, 241, 25–32.
- Heller, P.L., and Angevine, C.L. (1985) Sea-level cycles during the growth of Atlantic-type oceans. *Earth and Planetary Science Letters*, 75, 417–426.
- Heller, P.L., Bowdler, S.S., Chambers, H.P., Coogan, J.C., Hagen, E.S., Shuster, M.W., Winslow, N.S., and Lawton, T.F. (1986) Time of initial thrusting in the Sevier orogenic belt, Idaho-Wyoming and Utah. *Geology*, 14, 388–391.
- Hermoza, W., Brusset, S., Baby, P., Gil, W., Roddaz, M., Guerrero, N., and Bolanos, M. (2005) The Huallaga foreland basin evolution: thrust propagation in a deltaic environment, northern Peruvian Andes. *Journal of South American Earth Sciences*, 19, 21–34.
- Hildebrand, A.R., Penfield, G.T., Kring, D.A., Pilkington, M., Camargo Z., A., Jacobsen, S.B., and Boynton, W.V. (1991) Chicxulub crater: a possible Cretaceous/Tertiary boundary impact crater on the Yucatan Peninsula, Mexico. *Geology*, 19, 867–871.
- Hilley, G.E., and Strecker, M.R. (2005) Processes of oscillatory basin filling and excavation in a tectonically active orogen: Quebrada del Toro basin, NW Argentina. *Geological Society of America Bulletin*, 117, 887–901.
- Hoffman, P.F., Dewey, J.F., and Burke, K. (1974) Aulacogens and their genetic relation to geosynclines with a Proterozoic example from Great Slave Lake, Canada. *Society of Economic Paleontologists and Mineralogists Special Publication*, 19, 38–55.
- Hopson, C.A., Mattinson, J.M., and Pessagno, E.A., Jr. (1981) Coast Range ophiolite, western California, in Ernst, W.G., ed., *The geotectonic development of California* (Rubey, vol. 1). Englewood Cliffs, NJ, Prentice Hall, 418–510.
- Hopson, C. A., Mattinson, J.M., Pessagno, E.A., and Luyendyk, B.P. (2008) California Coast Range ophiolite: composite Middle and Late Jurassic oceanic lithosphere. *Geological Society of America Special Paper*, 438, 1–101.
- Hornafius, J.S., Luyendyk, B.P., Terres, R.R., and Kamerling, M.J. (1986) Timing and extent of Neogene tectonic rotation in the western Transverse Ranges, California. *Geological Society of America Bulletin*, 97, 1476–1487.
- Horton, B.K. (1998) Sediment accumulation on top of the Andean orogenic wedge: Oligocene to late Miocene basins of the Eastern Cordillera, southern Bolivia. *Geological Society of America Bulletin*, 110, 1174–1192.
- Horton, B.K. (2005) Revised deformation history of the central Andes: inferences from Cenozoic foredeep and intermontane basins of the Eastern Cordillera, Bolivia. *Tectonics*, 24, TC3011, 18 p.
- Horton, B.K., and DeCelles, P.G. (1997) The modern foreland basin system adjacent to the Central Andes. *Geology*, 25, 895–898.

- Horton, B.K., Hampton, B.A., LaReau, B.N., and Baldellón, E. (2002) Tertiary provenance history of the northern and central Altiplano (central Andes, Bolivia): A detrital record of plateau-margin tectonics. *Journal of Sedimentary Research*, 72, 711–726.
- Howard, J.P., Cunningham, W.D., Davies, S.J., Dijkstra, A. H., and Badarch, G. (2003) The stratigraphic and structural evolution of the Dzereg basin, western Mongolia: clastic sedimentation, transpressional faulting and basin destruction in an intraplate, intracontinental setting. *Basin Research*, 15, 45–72.
- Howell, D.G., Jones, D.L., and Schermer, E.R. (1985) Tectonostratigraphic terranes of the circum-Pacific region. *Circum-Pacific Council for Energy and Mineral Resources Earth Sciences Series*, 1, 3–30.
- Howell, P.D., and van der Pluijm, B.A. (1999) Structural sequences and styles of subsidence in the Michigan basin. *Geological Society of America Bulletin*, 111, 974–991.
- Hoy, R.G., and Ridgway, K.D. (1997) Structural and sedimentological development of footwall growth synclines along an intraforeland uplift, east-central Bighorn Mountains, Wyoming. *Geological Society of America Bulletin*, 109, 915–935.
- Hoy, R.G., and Ridgway, K.D. (2002) Syndepositional thrust-related deformation and sedimentation in an Ancestral Rocky Mountains basin, central Colorado trough, Colorado, USA. *Geological Society of America Bulletin*, 114, 804–828.
- Hsiao, L-Y., Graham, S.A., and Tilander, N. (2004) Seismic reflection imaging of a major strike-slip fault zone in a rift system: Paleogene structure and evolution of the Tan-Lu fault system, Liaodong Bay, Bohai, offshore China. *American Association of Petroleum Geologists Bulletin*, 88, 71–97.
- Hudec, M.R., Jackson, M.P.A., and Schultz-Ela, D.D. (2009) The paradox of minibasin subsidence into salt: Clues to the evolution of crustal basins. *Geological Society of America Bulletin*, 121, 201–221.
- Hurwitz, S., Garfunkel, Z., Ben-Gai, Y., Reznikov, M., Rotstein, Y., and Gvirtzman, H. (2002) The tectonic framework of a complex pull-apart basin: seismic reflection observations in the Sea of Galilee, Dead Sea transform. *Tectonophysics*, 359, 289–306.
- Ingersoll, R.V. (1979) Evolution of the Late Cretaceous forearc basin, northern and central California. *Geological Society of America Bulletin*, 90, part 1, 813–826.
- Ingersoll, R.V. (1982) Initiation and evolution of the Great Valley forearc basin of northern and central California, U.S.A. *Geological Society of London Special Publication*, 10, 459–467.
- Ingersoll, R.V. (1988) Tectonics of sedimentary basins. *Geological Society of America Bulletin*, 100, 1704–1719.
- Ingersoll, R.V. (1997) Phanerozoic tectonic evolution of central California and environs. *International Geology Review*, 39, 957–972.
- Ingersoll, R.V. (2000) Models for origin and emplacement of Jurassic ophiolites of northern California, in Dilek, Y., Moores, E.M., Elthon, D., and Nicolas, A., eds., *Ophiolites and oceanic crust: new insights from field studies and the Ocean Drilling Program*. *Geological Society of America Special Paper*, 349, 395–402.
- Ingersoll, R.V. (2001) Structural and stratigraphic evolution of the Rio Grande rift, northern New Mexico and southern Colorado. *International Geology Review*, 43, 867–891.
- Ingersoll, R.V. (2008a) Subduction-related sedimentary basins of the U.S.A. Cordillera, in Miall, A.D., ed., *The sedimentary basins of the United States and Canada*. *Sedimentary Basins of the World*, vol. 5. Amsterdam, Elsevier, 395–428.
- Ingersoll, R.V. (2008b) Reconstructing southern California, in Spencer, J.E., and Titley, S.R., eds., *Ores and orogenesis: Circum-Pacific tectonics, geologic evolution, and ore deposits*. *Arizona Geological Society Digest* 22, 409–417.
- Ingersoll, R.V., and Busby, C.J. (1995) Tectonics of sedimentary basins, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 1–51.
- Ingersoll, R.V., and Rumelhart, P.E. (1999) Three-stage evolution of the Los Angeles basin, southern California. *Geology*, 27, 593–596 (Correction, 27, 864).
- Ingersoll, R.V., and Schweickert, R.A. (1986) A plate-tectonic model for Late Jurassic ophiolite genesis, Nevada orogeny and forearc initiation, northern California. *Tectonics*, 5, 901–912.
- Ingersoll, R.V., Graham, S.A., and Dickinson, W.R. (1995) Remnant ocean basins, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 363–391.
- Ingersoll, R.V., Dickinson, W.R., and Graham, S.A. (2003) Remnant-ocean submarine fans: largest sedimentary systems on Earth, in Chan, M.A., and Archer, A.W., eds., *Extreme depositional environments: mega end members in geologic time*. *Geological Society of America Special Paper*, 370, 191–208.
- Itoh, Y., Takemura, K., Ishiyama, T., Tanaka, Y., and Iwaki, H. (2000) Basin formation at a contractional bend of a large transcurrent fault: Plio-Pleistocene subsidence of the Kobe and northern Osaka basins, Japan. *Tectonophysics*, 321, 327–341.
- Jacome, M.I., Kusznir, N., Audemard, F., and Flint, S. (2003) Formation of the Maturin foreland basin, eastern Venezuela: thrust sheet loading or subduction dynamic topography. *Tectonics*, 22, TC1046, 17 p.
- Jarrard, R.D. (1986) Relations among subduction parameters. *Reviews of Geophysics*, 24, 217–284.
- Jenkyns, H.C., and Wilson, P.A. (1999) Stratigraphy, paleoceanography, and evolution of Cretaceous Pacific Guyots: relics from a greenhouse Earth. *American Journal of Science*, 299, 341–392.
- Johnson, C.L. (2004) Polyphase evolution of the East Gobi basin: sedimentary and structural records of Mesozoic-Cenozoic intraplate deformation in Mongolia. *Basin Research*, 16, 79–99.
- Johnson, D.D., and Beaumont, C. (1995) Preliminary results from a planform kinematic model of orogen evolution, surface processes and the development of clastic foreland basin stratigraphy, in Dorobek, S.L., and Ross, G.M., eds., *Stratigraphic evolution of foreland basins*. *SEPM (Society for Sedimentary Geology) Special Publication*, 52, 3–24.
- Jones, M.A., Heller, P.L., Roca, E., Garces, M., and Cabrera, L. (2004) Time lag of syntectonic sedimentation across an

- alluvial basin: theory and example from the Ebro basin, Spain. *Basin Research*, 16, 467–488.
- Jordan, T.E. (1981) Thrust loads and foreland basin evolution, Cretaceous, western United States. *American Association of Petroleum Geologists Bulletin*, 65, 2506–2520.
- Jordan, T.E. (1995) Retroarc foreland and related basins, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 331–362.
- Jordan, T.E., and Allmendinger, R.W. (1986) The Sierras Pampeanas of Argentina: a modern analogue of Rocky Mountain foreland deformation. *American Journal of Science*, 286, 737–764.
- Jordan, T.E., Isacks, B.L., Ramos, V.A., and Allmendinger, R.W. (1983a) Mountain building in the central Andes. *Episodes*, 1983, 20–26.
- Jordan, T.E., Isacks, B.L., Allmendinger, R.W., Brewer, J.A., Ramos, V.A., and Ando, C.J. (1983b) Andean tectonics related to geometry of subducted Nazca plate. *Geological Society of America Bulletin*, 94, 341–361.
- Kanamori, H. (1986) Rupture process of subduction-zone earthquakes. *Annual Reviews of Earth and Planetary Science*, 14, 293–322.
- Kao, H., Gao, R., Rau, R.-J., Shi, D., Chen, R.-Y., Guan, Y., and Wu, F.T. (2001) Seismic image of the Tarim basin and its collision with Tibet. *Geology*, 29, 575–578.
- Karig, D.E. (1970) Kermadec arc-New Zealand tectonic confluence. *New Zealand Journal of Geology and Geophysics*, 13, 21–29.
- Karig, D.E., and Moore, G.F. (1975) Tectonically controlled sedimentation in marginal basins. *Earth and Planetary Science Letters*, 26, 233–238.
- Karig, D.E., and Sharman, G.F., III (1975) Subduction and accretion in trenches. *Geological Society of America Bulletin*, 86, 377–389.
- Katzman, R., tenBrink, U.S., and Lin, J. (1995) Three-dimensional modeling of pull-apart basins: implications for the tectonics of the Dead Sea basin. *Journal of Geophysical Research*, 100, 6295–6312.
- Kay, M. (1951) North American geosynclines. *Geological Society of America Memoir* 48, 143 p.
- Kellogg, K.S., and Minor, S.A. (2005) Pliocene transpressional modification of depositional basins by convergent thrusting adjacent to the “Big Bend” of the San Andreas fault: an example from Lockwood Valley, southern California. *Tectonics*, 24, TC1004, 12 p.
- Kempf, O., and Pfiffner, O.A. (2004) Early Tertiary evolution of the north Alpine foreland basin of the Swiss Alps and adjoining areas. *Basin Research*, 16, 549–567.
- Kimbrough, D.L. (1984) Paleogeographic significance of the Middle Jurassic Gran Canon Formation, Cedros Island, Baja California, in Frizzell, V.A., Jr., ed., *Geology of the Baja California Peninsula*. Pacific Section, Society of Economic Paleontologists and Mineralogists, Los Angeles, 107–118.
- King, P.B. (1966) The North American Cordillera. *Canadian Institute of Mining and Metallurgy Special*, 8, 1–25.
- Kinsman, D.J.J. (1975) Rift valley basins and sedimentary history of trailing continental margins, in Fischer, A.G. and Judson, S., eds., *Petroleum and global tectonics*. Princeton, NJ, Princeton University Press, 83–126.
- Klein, G.deV. (1985) The control of depositional depth, tectonic uplift, and lockwoodism on sedimentation processes in the back-arc basins of the western Pacific Ocean. *Journal of Geology*, 93, 1–25.
- Klein, G.deV. (1992) Climatic and tectonic sea-level gauge for midcontinent Pennsylvanian cyclothems. *Geology*, 20, 363–366.
- Klein, G.D. (1995) Intracratonic basins, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 459–478.
- Klein, G.deV., and Kupperman, J.B. (1992) Pennsylvanian cyclothems: methods of distinguishing tectonically-induced changes in sea level from climatically-induced changes. *Geological Society of America Bulletin*, 104, 166–175.
- Kluth, C.F. (1986) Plate tectonics of the Ancestral Rocky Mountains. *American Association of Petroleum Geologists Memoir* 42, 353–369.
- Kluth, C.F., and Coney, P.J. (1981) Plate tectonics of the ancestral Rocky Mountains. *Geology*, 9, 10–15.
- Kopp, H., and Kukowski, N. (2003) Backstop geometry and accretionary mechanics of the Sunda margin. *Tectonics*, 22, TC1072, 16 p.
- Koukouvelas, I.K., and Aydin, A. (2002) Fault structure and related basins of the North Aegean Sea and its surroundings. *Tectonics*, 21, TC1046, 17 p.
- Lash, G.G. (1988) Along-strike variations in foreland basin evolution: possible evidence for continental collision along an irregular margin. *Basin Research*, 1, 71–83.
- Lawton, T.F. (1994) Tectonic setting of Mesozoic sedimentary basins, Rocky Mountain region, United States, in Caputo, M.V., Peterson, J.A., and Franczyk, K.J., eds., *Mesozoic systems of the Rocky Mountain region, USA*. Rocky Mountain Section, Society for Sedimentary Geology, Denver, 1–25.
- Lawton, T.F. (2008) Laramide sedimentary basins, in Miall, A.D., ed., *The sedimentary basins of the United States and Canada*. *Sedimentary Basins of the World*, vol. 5. Amsterdam, Elsevier, 429–450.
- Lazar, M., Ben-Avraham, Z., and Schattner, U. (2006) Formation of sequential basins along a strike-slip fault: geophysical observations from the Dead Sea basin. *Tectonophysics*, 421, 53–69.
- Leeder, M.R. (1995) Continental rifts and proto-oceanic rift troughs, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 119–148.
- Leeder, M.R., and Gawthorpe, R.L. (1987) Sedimentary models for extensional tilt-block/half-graben basins. *Geological Society of London Special Publication*, 28, 139–152.
- Leever, K.A., Bertotti, G., Zoetemeijer, R., Matenco, L., and Cloetingh, S.A.P.L. (2006) The effects of lateral variation in lithospheric strength on foredeep evolution: implications for the East Carpathian foredeep. *Tectonophysics*, 421, 251–267.
- Leverenz, A. (2000) Trench-sedimentation versus accreted submarine fan: an approach to regional-scale facies analysis in a Mesozoic accretionary complex: “Torlesse” terrane, northeastern North Island, New Zealand. *Sedimentary Geology*, 132, 125–160.
- Lin, A.T., and Watts, A.B. (2002) Origin of the west Taiwan basin by orogenic loading and flexure of a rifted continental margin. *Journal of Geophysical Research*, 107, n. B9, 2185.

- Lipman, P.W. (1992) Magmatism in the Cordilleran United States; progress and problems, in Burchfiel, B.C., Lipman, P.W., and Zoback, M.L., eds., *The Cordilleran orogen: conterminous U.S. Geology of North America*, G3. Boulder, CO, Geological Society of America, 481–514.
- Lonsdale, P. (1975) Sedimentation and tectonic modification of the Samoan archipelagic apron. *American Association of Petroleum Geologists Bulletin*, 59, 780–798.
- Lonsdale, P. (1991) Structural patterns of the Pacific floor offshore of peninsular California. *American Association of Petroleum Geologists Memoir* 47, 87–125.
- Lucente, C.C. (2004) Topography and palaeogeographic evolution of a middle Miocene foredeep basin plain (northern Apennines, Italy). *Sedimentary Geology*, 170, 107–134.
- Luyendyk, B.P. (1991) A model for Neogene crustal rotations, transtension, and transpression in southern California. *Geological Society of America Bulletin*, 103, 1528–1536.
- Luyendyk, B.P., and Hornafius, J.S. (1987) Neogene crustal rotations, fault slip, and basin development in southern California, in Ingersoll, R.V., and Ernst, W.G., eds., *Cenozoic basin development of coastal California* (Rubey, vol. 6). Englewood Cliffs, NJ, Prentice Hall, 259–283.
- Luyendyk, B.P., Kamerling, M.J., and Terres, R. (1980) Geometric model for Neogene crustal rotations in southern California. *Geological Society of America Bulletin*, 91, Part I, 211–217.
- Mack, G.H., Seager, W.R., and Leeder, M.R. (2003) Synclinal-horst basins: examples from the southern Rio Grande rift and southern transition zone of southwestern New Mexico, USA. *Basin Research*, 15, 365–377.
- Maillard, A., and Mauffret, A. (1999) Crustal structure and riftogenesis of the Valencia Trough (north-western Mediterranean Sea). *Basin Research*, 11, 357–379.
- Maldonado, A., Larter, R.D., and Aldaya, F. (1994) Forearc tectonic evolution of the South Shetland margin, Antarctic Peninsula. *Tectonics*, 13, 1345–1370.
- Mann, P. (1997) Model for the formation of large, transtensional basins in zones of tectonic escape. *Geology*, 25, 211–214.
- Mann, P., Hempton, M.R., Bradley, D.C., and Burke, K. (1983) Development of pull-apart basins. *Journal of Geology*, 91, 529–554.
- Marin, L.E., Sharpton, V.L., Fucugauchi, J.U., Smit, J., Sikora, P., Carney, C., Rebolledo-Vieyra, M. (2001) Stratigraphy at ground zero: a contemporary evaluation of well data in the Chicxulub impact basin. *International Geology Review*, 43, 1145–1149.
- Marsaglia, K.M. (1995) Interarc and backarc basins, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 299–329.
- Marshak, S., Karlstrom, K., and Timmons, J.M. (2000) Inversion of Proterozoic extensional faults: an explanation for the pattern of Laramide and Ancestral Rockies intracratonic deformation, United States. *Geology*, 28, 735–738.
- Masclé, A., Puigdefabregas, C., Luterbacher, H.P., and Fernandez, M., eds. (1998) *Cenozoic foreland basins of western Europe*. Geological Society of London Special Publication, 134, 427 p.
- Mattei, M., Cipollari, P., Cosentino, D., Argentieri, A., Rossetti, F., Speranza, F., and Di Bella, L. (2002) The Miocene tectono-sedimentary evolution of the southern Tyrrhenian Sea: stratigraphy, structural and palaeomagnetic data from the on-shore Amantea basin (Calabrian arc, Italy). *Basin Research*, 14, 147–168.
- Mattern, F. (2001) Permo-Silesian movements between Baltica and western Europe: tectonics and ‘basin families’. *Terra Nova*, 13, 368–375.
- May, S.R., Ehman, K.D., Gray, G.G., and Crowell, J.C. (1993) A new angle on the tectonic evolution of the Ridge basin, a “strike-slip” basin in southern California. *Geological Society of America Bulletin*, 105, 1357–1372.
- McBride, J.H., and Nelson, W.J. (1999) Style and origin of mid-Carboniferous deformation in the Illinois basin, USA: Ancestral Rockies deformation? *Tectonophysics*, 305, 249–273.
- McClay, K.R., ed. (1992) *Thrust tectonics*. London, Chapman and Hall, 447 p.
- McClay, K., and Bonora, M. (2001) Analog models of restraining stepovers in strike-slip fault systems. *American Association of Petroleum Geologists Bulletin*, 85, 233–260.
- Meng, Q.-R., Wange, E., and Hu, J.-M. (2005) Mesozoic sedimentary evolution of the northwest Sichuan basin: implication for continued clockwise rotation of the South China block. *Geological Society of America Bulletin*, 117, 396–410.
- Miall, A.D. (1995) Collision-related foreland basins, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 393–424.
- Miall, A.D. (2008) The southern midcontinent, Permian basin, and Ouachitas, in Miall, A.D., ed., *The sedimentary basins of the United States and Canada* *Sedimentary Basins of the World*, vol. 5. Amsterdam, Elsevier, 297–327.
- Miyashiro, A. (1973) *Metamorphism and metamorphic belts*. New York, John Wiley, 492 p.
- Molnar, and Atwater, T. (1978) Interarc spreading and Cordilleran tectonics as alternates related to age of subducted oceanic lithosphere. *Earth and Planetary Science Letters*, 41, 330–340.
- Molnar, P., and Tapponnier, P. (1975) Cenozoic tectonics of Asia: effects of a continental collision. *Science*, 189, 419–426.
- Moore, G.F., and Karig, D.E. (1976) Development of sedimentary basins on the lower trench slope. *Geology*, 4, 693–697.
- Moore, G.W. (1973) Westward tidal lag as the driving force of plate tectonics. *Geology*, 1, 99–100.
- Morgan, P., and Baker, B.H. (1983) Introduction: processes of continental rifting. *Tectonophysics*, 94, 1–10.
- Moss, S.J., and McCarthy, A.J. (1997) Foreland basin systems: discussion. *Basin Research*, 9, 171–172.
- Mountney, N.P., and Westbrook, G.K. (1996) Modelling sedimentation in ocean trenches: the Nankai Trough from 1 Ma to the present. *Basin Research*, 8, 85–101.
- Mountney, N.P., and Westbrook, G.K. (1997) Quantitative analysis of Miocene to Recent forearc basin evolution along the Colombian convergent margin. *Basin Research*, 9, 177–196.
- Murphy, J.B., Keppie, J.D., and Nance, R.D. (1999) Fault reactivation within Avalonia: plate margin to continental interior deformation. *Tectonophysics*, 305, 183–204.

- Najman, Y., and Garzanti, E. (2000) Reconstructing early Himalayan tectonic evolution and paleogeography from Tertiary foreland basin sedimentary rocks, northern India. *Geological Society of America Bulletin*, 112, 435–449.
- Najman, Y., Johnson, K., White, N., and Oliver, G. (2004) Evolution of the Himalayan foreland basin, NW India. *Basin Research*, 16, 1–24.
- Naylor, M., and Sinclair, H.D. (2008) Pro- vs. retro-foreland basins. *Basin Research*, 20, 285–303.
- Nicolas, A. (1985) Novel type of crust produced during continental rifting. *Nature*, 315, 112–115.
- Nicholson, C., Sorlien, C.C., Atwater, T., Crowell, J.C., and Luyendyk, B.P. (1994) Microplate capture, rotation of the western Transverse Ranges, and initiation of the San Andreas transform as a low-angle fault system. *Geology*, 22, 491–495.
- Nilsen, T.H., and Sylvester, A.G. (1995) Strike-slip basins, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 425–457.
- Norris, R.J., and Cooper, A.F. (1995) Origin of small-scale segmentation and transpressional thrusting along the Alpine fault, New Zealand. *Geological Society of America Bulletin*, 107, 231–240.
- Okay, A.I., Sengor, A.M.C., and Gorur, N. (1994) Kinematic history of the opening of the Black Sea and its effect on the surrounding regions. *Geology*, 22, 267–270.
- Okay, A.I., Demirbag, E., Kurt, H., Okay, N., and Kuscü, I. (1999) An active, deep marine strike-slip basin along the north Anatolian fault in Turkey. *Tectonics*, 18, 129–147.
- Okay, A.I., Tuysuz, O., and Kaya, S. (2004) From transpression to transtension: changes in morphology and structure around a bend on the North Anatolian fault in the Marmara region. *Tectonophysics*, 391, 259–282.
- Oldow, J.S. (1984) Evolution of a late Mesozoic back-arc fold and thrust belt in northwestern Great Basin, U.S.A. *Tectonophysics*, 102, 245–274.
- Oldow, J.S., Bally, A.W., Ave Lallement, H.G., and Leeman, W.P. (1989) Phanerozoic evolution of the North American Cordillera; United States and Canada, in Bally, A.W., and Palmer, A.R., eds., *The geology of North America; an overview [The Geology of North America Volume A]*. Boulder, CO, Geological Society of America, 139–232.
- Ori, G.G., and Friend, P.F. (1984) Sedimentary basins formed and carried piggyback on active thrust sheets. *Geology*, 12, 475–478.
- Parnell, J., Osinski, G.R., Lee, P., Green, P.F., and Baron, M.J. (2005) Thermal alteration of organic matter in an impact crater and the duration of postimpact heating. *Geology*, 33, 373–376.
- Parsons, B., and Sclater, J.G. (1977) An analysis of the variation of ocean floor bathymetry and heat flow with age. *Journal of Geophysical Research*, 82, 803–827.
- Pascucci, V., Merlin, S., and Martini, I.P. (1999) Seismic stratigraphy of the Miocene-Pleistocene sedimentary basins of the northern Tyrrhenian Sea and western Tuscany (Italy). *Basin Research*, 11, 337–356.
- Pfiffner, O.A., Schlunegger, F., and Buitner, S.J.H. (2002) The Swiss Alps and their peripheral foreland basin: stratigraphic response to deep crustal processes. *Tectonics*, 21, TC1054, 16 p.
- Pindell, J.L. (1985) Alleghenian reconstruction and subsequent evolution of the Gulf of Mexico, Bahamas, and proto-Caribbean. *Tectonics*, 4, 1–39.
- Pindell, J., and Dewey, J.F. (1982) Permo-Triassic reconstruction of western Pangea and the evolution of the Gulf of Mexico/Caribbean region. *Tectonics*, 1, 179–211.
- Pivnik, D.A., and Khan, M.J. (1996) Transition from foreland- to piggyback-basin deposition, Plio-Pleistocene upper Siwalik Group, Shinghar Range, NW Pakistan. *Sedimentology*, 43, 631–646.
- Pivnik, D.A., and Wells, N.A. (1996) The transition from Tethys to the Himalaya as recorded in northwest Pakistan. *Geological Society of America Bulletin*, 108, 1295–1313.
- Platt, J.P. (1986) Dynamics of orogenic wedges and the uplift of high-pressure metamorphic rocks. *Geological Society of America Bulletin*, 97, 1037–1053.
- Pope, K.O., Ocampo, A.C., and Duller, C.E. (1991) Mexican site for K/T impact crater. *Nature*, 351, 105.
- Prather, B.E., Booth, J.R., Steffens, G.S., and Craig, P.A. (1998) Classification, lithologic calibration, and stratigraphic succession of seismic facies of intraslope basins, deep-water Gulf of Mexico. *American Association of Petroleum Geologists Bulletin*, 82, 701–728.
- Price, R.A. (1973) Large-scale gravitational flow of supracrustal rocks, southern Canadian Rockies, in DeJong, K.A., and Scholten, R., eds., *Gravity and tectonics*. New York, John Wiley and Sons, 491–502.
- Purser, B.H., and Bosence, D.W.J., eds. (1998) *Sedimentation and tectonics in rift basins: Red Sea: Gulf of Aden*. New York, Chapman and Hall, 663 p.
- Quinlan, G.M. (1987) Models of subsidence mechanisms in intracratonic basins, and their applicability to North American examples. *Canadian Society of Petroleum Geologists Memoir* 12, 463–481.
- Rahe, B., Ferrill, D.A., and Morris, A.P. (1998) Physical analog modeling of pull-apart basin evolution. *Tectonophysics*, 285, 21–40.
- Rangin, C., LePichon, X., Demirbag, E., and Imren, C. (2004) Strain localization in the Sea of Marmara: propagation of the north Anatolian fault in a now inactive pull-apart. *Tectonics*, 23, TC2014, 18 p.
- Reading, H.G. (1980) Characteristics and recognition of strike-slip fault systems. *International Association of Sedimentologists Special Publication*, 4, 7–26.
- Ricketts, B.D. (2008) Cordilleran sedimentary basins of western Canada record 180 million years of terrane accretion, in Miall, A.D., ed., *The sedimentary basins of the United States and Canada Sedimentary Basins of the World*, vol. 5. Amsterdam, Elsevier, 363–394.
- Riggs, N.R., and Busby-Spera, C.J. (1991) Facies analysis of an ancient, dismembered, large caldera complex and implications for intra-arc subsidence: Middle Jurassic strata of Cobre Ridge, southern Arizona, USA. *Sedimentary Geology*, 74, 39–68.
- Ritts, B.D., Weislogel, A., Graham, S.A., and Darby, B.J. (2009) Mesozoic tectonics and sedimentation of the giant polyphase nonmarine intraplate Ordos basin, western North China block. *International Geology Review*, 51, 95–115.
- Robertson, A.H.F. (1989) Palaeoceanography and tectonic setting of the Jurassic Coast Range ophiolite, central

- California: evidence from the extrusive rocks and the volcanoclastic sediment cover. *Marine and Petroleum Geology*, 6, 194–220.
- Robertson, A., and Deggan, P. (1994) The Dras arc complex: lithofacies and reconstruction of a Late Cretaceous oceanic volcanic arc in the Indus suture zone, Ladakh Himalaya. *Sedimentary Geology*, 92, 117–145.
- Rollet, N., Deverchere, J., Beslier, M.-O., Guennoc, P., Rehault, J.P., Sosson, M., and Truffert, C. (2002) Back arc extension, tectonic inheritance, and volcanism in the Ligurian Sea, Western Mediterranean. *Tectonics*, 21 (3), 23 p.
- Root, S., and Onasch, C.M. (1999) Structure and tectonic evolution of the transitional region between the central Appalachian foreland and interior cratonic basins. *Tectonophysics*, 305, 205–223.
- Rosendahl, B.R. (1987) Architecture of continental rifts with special reference to East Africa. *Annual Review of Earth and Planetary Sciences*, 15, 445–503.
- Rowan, M.G., Lawton, T.F., Giles, K.A., and Ratliff, R.A. (2003) Near-salt deformation in La Popa basin, Mexico, and the northern Gulf of Mexico: a general model for passive diapirism. *American Association of Petroleum Geologists Bulletin*, 87, 733–756.
- Rowley, D.B., and Kidd, W.S.F. (1981) Stratigraphic relationships and detrital composition of the medial Ordovician flysch of western New England: implications for the tectonic evolution of the Taconic orogeny. *Journal of Geology*, 89, 199–218.
- Salvador, A., ed. (1991) The Gulf of Mexico basin [The Geology of North America, J]. Boulder, CO, Geological Society of America, 568 p.
- Saylor, B.Z. (2003) Sequence stratigraphy and carbonate-siliciclastic mixing in a terminal Proterozoic foreland basin, Urusis Formation, Nama Group, Namibia. *Journal of Sedimentary Research*, 73, 264–279.
- Saylor, J., DeCelles, P., Gehrels, G., Murphy, M., Zhang, R., and Kapp, P. (2010) Basin formation in the high Himalaya by arc-parallel extension and tectonic damming: Zhada basin, southwestern Tibet. *Tectonics*, 29, TC1004, 24 p.
- Schellart, W.P. (2007) The potential influence of subduction zone polarity on overriding plate deformation, trench migration and slab dip angle. *Tectonophysics*, 445, 363–372.
- Schellart, W.P. (2008) Overriding plate shortening and extension above subduction zones. a parametric study to explain formation of the Andes Mountains. *Geological Society of America Bulletin*, 120, 1441–1454.
- Schermer, E.R., and Busby, C. (1994) Jurassic magmatism in the central Mojave Desert: implications for arc paleogeography and preservation of continental volcanic sequences. *Geological Society of America Bulletin*, 106, 767–790.
- Schneider, C.L., Hummon, C., Yeats, R.S., and Huftile, G.L. (1996) Structural evolution of the northern Los Angeles basin, California, based on growth strata. *Tectonics*, 15, 341–355.
- Scholl, D.W., Buffington, E.C., and Marlow, M.S. (1975) Plate tectonics and the structural evolution of the Aleutian-Bering Sea region. *Geological Society of America Special Paper*, 151, 1–31.
- Scholl, D.W., Von Huene, R., Vallier, T.L. and Howell, D.G. (1980) Sedimentary masses and concepts about tectonic processes at underthrust ocean margins. *Geology*, 8, 564–568.
- Schweickert, R.A. (1976) Shallow-level plutonic complexes in the eastern Sierra Nevada, California and their tectonic implications. *Geological Society of America Special Paper*, 176, 58 p.
- Schweickert, R.A. (1978) Triassic and Jurassic paleogeography of the Sierra Nevada and adjacent regions, California and western Nevada, in Howell, D.G., and McDougall, K.A., eds., *Mesozoic paleogeography of the western United States*. Pacific Section, Society of Economic Paleontologists and Mineralogists Pacific Coast Paleogeography Symposium 2, 361–384.
- Schweller, W.J. and Kulm, L.D. (1978) Depositional patterns and channelized sedimentation in active eastern Pacific trenches, in Stanley, D.J. and Kelling, G., eds., *Sedimentation in submarine canyons, fans, and trenches*. Stroudsburg, PA, Dowden, Hutchinson and Ross, Inc., 311–324.
- Sclater, J.G., Anderson, R.N., and Bell, M.L. (1971) Elevation of ridges and evolution of the central eastern Pacific. *Journal of Geophysical Research*, 76, 7888–7915.
- Seeber, L., Emre, O., Cormier, M.-H., Sorlien, C.C., McHugh, C.M.G., Polonia, A., Ozer, N., and Cagatay, N. (2004) Uplift and subsidence from oblique slip: the Ganos-Marmara bend of the North Anatolian transform, western Turkey. *Tectonophysics*, 391, 239–258.
- Sengor, A.M.C. (1976) Collision of irregular continental margins: implications for foreland deformation of Alpine-type orogens. *Geology*, 4, 779–782.
- Sengor, A.M.C. (1995) Sedimentation and tectonics of fossil rifts, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 53–117.
- Sengor, A.M.C. and Burke, K. (1978) Relative timing of rifting and volcanism on Earth and its tectonic implications. *Geophysical Research Letters*, 5, 419–421.
- Sengor, A.M.C., Burke, K., and Dewey, J.F. (1978) Rifts at high angles to orogenic belts: tests for their origin and the upper Rhine graben as an example. *American Journal of Science*, 278, 24–40.
- Sengor, A.M.C., Cin, A., Rowley, D.B., and Nie, S. (1991) Magmatic evolution of the Tethysides: a guide to reconstruction of collage history. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 87, 411–440.
- Sengor, A.M.C., Graham, S.A., and Biddle, K.T. (1996) Is the Tarim basin underlain by a Neoproterozoic oceanic plateau? *Geological Society of America Abstracts with Programs*, 28 (7), A–67.
- Shah, A.K., Brozena, J., Vogt, P., Daniels, D., and Plescia, J. (2005) New surveys of the Chesapeake Bay impact structure suggest melt pockets and target-structure effect. *Geology*, 33, 417–420.
- Shatsky, N.S. (1964) O progibach Donetskogo tipa, in Akademik N.S. Shatsky, *Izbrannie Trudi*, 2. “Nauka,” Moscow, 544–552.
- Sibuet, J.-C., Deffontaines, B., Hsu, S.-K., Thareau, N., Le Formal, J.-P., Liu, C.-S., and ACT party (1998) Okinawa trough backarc basin: early tectonic and magmatic evolution. *Journal of Geophysical Research*, 103, 30245–30267.
- Sinclair, H.D. (1997) Tectonostratigraphic model for underfilled peripheral foreland basins: an Alpine perspective. *Geological Society of America Bulletin*, 109, 324–346.

- Sitian, L., Shigong, Y., and Jerzykiewicz, T. (1995) Upper Triassic-Jurassic foreland sequences of the Ordos basin in China. *SEPM (Society for Sedimentary Geology) Special Publication*, 52, 233–241.
- Sloss, L.L. (1988) Tectonic evolution of the craton in Phanerozoic time, in Sloss, L.L., ed., *Sedimentary cover: North American craton: U.S. The Geology of North America*, D-2. Boulder, CO, Geological Society of America, 25–51.
- Smit, J., Roep, T.B., Alvarez, W., Montanari, A., Claeys, P., Grajales-Nishimura, J.M., and Bermúdez-Santana, J. (1996) Coarse-grained, clastic sandstone complex at the K/T boundary around the Gulf of Mexico: Deposition by tsunami waves induced by the Chicxulub impact? *Geological Society of America Special Paper*, 307, 151–186.
- Smith, G.A., and Landis, C.A. (1995) Intra-arc basins, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 263–298.
- Sobel, E.R. (1999) Basin analysis of the Jurassic-Lower Cretaceous southwest Tarim basin, northwest China. *Geological Society of America Bulletin*, 111, 709–724.
- Sobel, E.R., and Strecker, M.R. (2003) Uplift, exhumation and precipitation: tectonic and climatic control of late Cenozoic landscape evolution in the northern Sierras Pampeanas, Argentina. *Basin Research*, 15, 431–451.
- Soreghan, G.S. (1994) Stratigraphic responses to geologic processes: Late Pennsylvanian eustasy and tectonics in the Pedregosa and Orogrande basins, Ancestral Rocky Mountains. *Geological Society of America Bulletin*, 106, 1195–1211.
- Sowerbutts, A. (2000) Sedimentation and volcanism linked to multiphase rifting in an Oligo-Miocene intra-arc basin, Anglona, Sardinia. *Geological Magazine*, 137, 395–418.
- Sowerbutts, A.A., and Underhill, J.R. (1998) Sedimentary response to intra-arc extension: controls on Oligo-Miocene deposition, Sarcidano sub-basin, Sardinia. *Journal of the Geological Society of London*, 155, 491–508.
- Speed, R.C., and Sleep, N.H. (1982) Antler orogeny and foreland basin: a model. *Geological Society of America Bulletin*, 93, 815–828.
- Stanley, R.S., and Ratcliffe, N.M. (1985) Tectonic synthesis of the Taconian orogeny in western New England. *Geological Society of America Bulletin*, 96, 1227–1250.
- Steckler, M.S., and tenBrink, U.S. (1986) Lithospheric strength variations as a control on new plate boundaries: examples from the northern Red Sea region. *Earth and Planetary Science Letters*, 79, 120–132.
- Stein, C.A. and Stein, S. (1992) A model for the global variation in oceanic depth and heat flow with lithospheric age. *Nature*, 359, 123–129.
- Stern, R.J., and Bloomer, S.H. (1992) Subduction zone infancy: examples from the Eocene Izu-Bonin-Mariana and Jurassic California arcs. *Geological Society of America Bulletin*, 104, 1621–1636.
- Stewart, S.A. (2003) How will we recognize buried impact craters in terrestrial sedimentary basins? *Geology*, 31, 929–932.
- Stewart, S.A., and Allen, P.J. (2002) A 20-km-diameter multi-ringed impact structure in the North Sea. *Nature*, 418, 520–523. (Also see Discussion and Reply, 428.)
- Stewart, S.A., and Allen, P.J. (2005) 3D seismic reflection mapping of the Silverpit multi-ringed crater, North Sea. *Geological Society of America Bulletin*, 117, 354–368.
- Stockmal, G.S., Beaumont, C., and Boutilier, R. (1986) Geodynamic models of convergent margin tectonics: transition from rifted margin to overthrust belt and consequences for foreland-basin development. *American Association of Petroleum Geologists Bulletin*, 70, 181–190.
- Stollhofen, H., and Stanistreet, I.G. (1994) Interaction between bimodal volcanism, fluvial sedimentation and basin development in the Permo-Carboniferous Saar-Nahe basin (south-west Germany). *Basin Research*, 6, 245–267.
- Talling, P.J., Lawton, T.F., Burbank, D.W., and Hobbs, R.S. (1995) Evolution of latest Cretaceous-Eocene nonmarine deposystems in the Axhandle piggyback basin of central Utah. *Geological Society of America Bulletin*, 107, 297–315.
- Tamaki, K., and Honza, E. (1991) Global tectonics and formation of marginal basins: role of the western Pacific. *Episodes*, 14, 224–230.
- Tanner, P.W.G., and Rex, D.C. (1979) Timing of events on a Lower Cretaceous island-arc marginal basin system on South Georgia. *Geological Magazine*, 116, 167–179.
- Tapponnier, P., Peltzer, G., LeDain, A.Y., Armijo, R., and Cobbold, P. (1982) Propagating extrusion tectonics in Asia: new insights from simple experiments with plasticine. *Geology*, 10, 611–616.
- Tarapoonca, M., Garcia-Castellanos, D., Bertotti, G., Matenco, L., Cloetingh, S.A.P.L., and Dinu, C. (2004) Role of the 3-D distributions of load and lithospheric strength in orogenic arcs: polystage subsidence in the Carpathians foredeep. *Earth and Planetary Science Letters*, 221, 163–180.
- Taylor, B. (2006) The single largest oceanic plateau: Ontong Java-Manihiki-Hikurangi. *Earth and Planetary Science Letters*, 241, 372–380.
- Taylor, B., and Karner, G.D. (1983) On the evolution of marginal basins. *Reviews of Geophysics and Space Physics*, 21, 1727–1741.
- tenBrink, U.S., Marshak, S., and Bruna, J.-L. G. (2009) Bivergent thrust wedges surrounding oceanic island arcs: insight from observations and sandbox models of north-eastern Caribbean plate. *Geological Society of America Bulletin*, 121, 1522–1536.
- Thomas, W.A. (1983) Continental margins, orogenic belts, and intracratonic structures. *Geology*, 11, 270–272.
- Thomas, W.A. (1985) The Appalachian-Ouachita connection: Paleozoic orogenic belt at the southern margin of North America. *Annual Review of Earth and Planetary Sciences*, 13, 175–199.
- Thomas, W.A. (1991) The Appalachian-Ouachita rifted margin of southeastern North America. *Geological Society of America Bulletin*, 103, 415–431.
- Thomas, W.A. (1995) Diachronous thrust loading and fault partitioning of the Black Warrior foreland basin within the Alabama recess of the late Paleozoic Appalachian: Ouachita thrust belt. *SEPM (Society for Sedimentary Geology) Special Publication*, 52, 111–126.
- Thornburg, T.M., and Kulm, L.D. (1987) Sedimentation in the Chile Trench: depositional morphologies, lithofacies, and stratigraphy. *Geological Society of America Bulletin*, 98, 33–52. (Also see Discussion and Reply, 99, 598–600.)
- Tobisch, O.T., Saleeby, J.B., and Fiske, R.S. (1986) Structural history of continental volcanic arc rocks, eastern

- Sierra Nevada, California: a case for extensional tectonics. *Tectonics*, 5, 65–94.
- Trop, J.M. (2008) Latest Cretaceous forearc basin development along an accretionary convergent margin: south-central Alaska. *Geological Society of America Bulletin*, 120, 207–224.
- Trop, J.M., and Ridgway, K.D. (1997) Petrofacies and provenance of a Late Cretaceous suture zone thrust-top basin, Cantwell basin, central Alaska Range. *Journal of Sedimentary Research*, 67, 469–485.
- Trop, J.M., Ridgway, K.D., and Sweet, A.R. (2004) Stratigraphy, palynology, and provenance of the Colorado Creek basin, Alaska, USA: Oligocene transpressional tectonics along the central Denali fault system. *Canadian Journal of Earth Sciences*, 41, 457–480.
- Tsikalas, F., Gudlaugsson, S.T., and Faleide, J.I. (1998) Collapse, infilling, and postimpact deformation of the Mjølner impact structure, Barents Sea. *Geological Society of America Bulletin*, 110, 537–552.
- Turner, J.P., Rosendahl, B.R., and Wilson, P.G. (2003) Structure and evolution of an obliquely sheared continental margin: Rio Muni, West Africa. *Tectonophysics*, 374, 41–55.
- Uba, C.E., Heubeck, C., and Hulka, C. (2005) Facies analysis and basin architecture of the Neogene subandean synorogenic wedge, southern Bolivia. *Sedimentary Geology*, 180, 91–123.
- Underwood, M.B., and Moore, G.F. (1995) Trenches and trench-slope basins, in Busby, C.J., and Ingersoll, R.V., eds., *Tectonics of sedimentary basins*. Oxford, Blackwell Science, 179–219.
- Underwood, M.B., Moore, G.F., Taira, A., Klaus, A., Wilson, M.E.J., Fergusson, C.L., Hirano, S., Steurer, J., and Leg 190 Shipboard Scientific Party (2003) Sedimentary and tectonic evolution of a trench-slope basin in the Nankai subduction zone of southwest Japan. *Journal of Sedimentary Research*, 73, 589–602.
- Uyeda, S., and Ben-Avraham, Z. (1972) Origin and development of the Philippine Sea. *Nature*, 240, 176–178.
- Uyeda, S., and Kanamori, H. (1979) Back-arc opening and the mode of subduction. *Journal of Geophysical Research*, 84, 1049–1061.
- Vanbrabant, Y., Braun, J. and Jongmans, D. (2002) Models of passive margin inversion: implications for the Rhenohercynian fold-and-thrust belt, Belgium and Germany. *Earth and Planetary Science Letters*, 202, 15–29.
- van der Pluijm, B.A., Craddock, J.P., Graham, B.R., and Harris, J.H. (1997) Paleostress in cratonic North America: implications for deformation of continental interiors. *Science*, 277, 794–796.
- Van der Werff, W. (1996) Variation in forearc basin development along the Sunda Arc, Indonesia. *Journal of Southeast Asian Earth Sciences*, 14, 331–349.
- Vincent, S.J., and Allen, M.B. (1999) Evolution of the Minle and Chaoshui basins, China: implications for Mesozoic strike-slip basin formation in central Asia. *Geological Society of America Bulletin*, 111, 725–742.
- Vincent, S.J., Allen, M.B., Ismail-Zadeh, A.D., Flecker, R., Foland, K.A., and Simmons, M.D. (2005) Insights from the Talysh of Azerbaijan into the Paleogene evolution of the South Caspian region. *Geological Society of America Bulletin*, 117, 1513–1533.
- Vink, G.E., Morgan, W.J., and Zhao, W.-L. (1984) Preferential rifting of continents: a source of displaced terranes. *Journal of Geophysical Research*, 89, 10072–10076.
- Wakabayashi, J., Hengesh, J.V., and Sawyer, T.L. (2004) Four-dimensional transform fault processes: progressive evolution of step-overs and bends. *Tectonophysics*, 392, 279–301.
- Waldron, J.W.F. (2004) Anatomy and evolution of a pull-apart basin, Stellarton, Nova Scotia. *Geological Society of America Bulletin*, 116, 109–127.
- Wernicke, B.P. (1992) Cenozoic extensional tectonics of the U.S. Cordillera, in Burchfiel, B.C., Lipman, P.W., and Zoback, M.L., eds., *The Cordilleran orogen: conterminous U.S. The Geology of North America*, G-3. Boulder, CO, Geological Society of America, 553–581.
- Willett, S., Beaumont, C., and Fullsack, P. (1993) Mechanical model for the tectonics of doubly vergent compressional orogens. *Geology*, 21, 371–374.
- Wilson, J.T. (1966) Did the Atlantic close and then re-open? *Nature*, 211, 676–681.
- Wilson, P.G., Turner, J.P., and Westbrook, G.K. (2003) Structural architecture of the ocean-continent boundary at an oblique transform margin through deep-imaging seismic interpretation and gravity modeling: Equatorial Guinea, West Africa. *Tectonophysics*, 374, 19–40.
- Winker, C.D. (1996) High-resolution seismic stratigraphy of a late Pleistocene submarine fan ponded by salt-withdrawal minibasins on the Gulf of Mexico continental slope. *Proceedings of 1996 Offshore Technology Conference*, Houston, 619–662.
- Winterer, E.L. (1973) Sedimentary facies and plate tectonics of equatorial Pacific. *American Association of Petroleum Geologists Bulletin*, 57, 265–282.
- Withjack, M.O., Schlische, R.W., and Olsen, P.E. (1998) Diachronous rifting, drifting, and inversion on the passive margin of central eastern North America: an analog for other passive margins. *American Association of Petroleum Geologists Bulletin*, 82, 817–835.
- Wood, R.A., Pettinga, J.R., Bannister, S., Lamarche, G., and McMorran, T.J. (1994) Structure of the Hammer strike-slip basin, Hope fault, New Zealand. *Geological Society of America Bulletin*, 106, 1459–1473.
- Woodcock, N.H. (2004) Life span and fate of basins. *Geology*, 32, 685–688.
- Worrall, D.M., and Snelson, S. (1989) Evolution of the northern Gulf of Mexico, with emphasis on Cenozoic growth faulting and the role of salt, in Bally, A.W., and Palmer, A.R., eds., *The geology of North America: an overview*. The Geology of North America, A. Boulder, CO, Geological Society of America, 91–138.
- Wright, T. (1991) Structural geology and tectonic evolution of the Los Angeles basin, California. *American Association of Petroleum Geologists Memoir*, 52, 35–134.
- Wyld, S.J. (2000) Triassic evolution of the arc and backarc of northwestern Nevada, and evidence for extensional tectonism. *Geological Society of America Special Paper*, 347, 185–207.
- Wyld, S.J. (2002) Structural evolution of a Mesozoic backarc fold-and-thrust belt in the U.S. Cordillera: new evidence from northern Nevada. *Geological Society of America Bulletin*, 114, 1452–1468.

- Yin, A., and Harrison, T.M. (2000) Geologic evolution of the Himalayan-Tibetan orogen. *Annual Reviews of Earth and Planetary Sciences*, 28, 211–280.
- Yin, A., and Ingersoll, R.V. (1997) A model for evolution of Laramide axial basins in the southern Rocky Mountains, U.S.A. *International Geology Review*, 39, 1113–1123.
- Yong, L., Allen, P.A., Densmore, A.L., and Qiang, X. (2003) Evolution of the Longmen Shan foreland basin (western Sichuan, China) during the Late Triassic Indosinian orogeny. *Basin Research*, 15, 117–138.
- Zattin, M., Stefani, C., and Martin, S. (2003) Detrital fission-track analysis and sedimentary petrofacies as keys of Alpine exhumation: the example of the Venetian foreland (European southern Alps, Italy). *Journal of Sedimentary Research*, 73, 1051–1061.
- Zoetemeijer, R., Tomek, C., and Cloetingh, S. (1999) Flexural expression of European continental lithosphere under the western outer Carpathians. *Tectonics*, 18, 843–861.

