

# 1

## Global Approaches to Understanding Protein Kinase Functions

*Jennifer L. Gorman and James R. Woodgett*

### 1.1 A Brief History of the Structure of the Human Kinome

Initial guesstimates of the complexity of the human kinome suggested that our genomes would code for around 1000 protein kinases [1]. Subsequent sequencing of the human genome identified 518 protein kinases [2] (subsequently revised to add some atypical kinases and to delete pseudokinases), which were classified into subgroups based on sequence similarities of their catalytic domain. These subclasses are the AGC kinases, CMGC kinases, CAMK (calcium/calmodulin-dependent kinase), STE kinases, tyrosine kinases, casein kinases, tyrosine kinase-like protein kinases as well as the receptor guanylyl cyclase (RGC) kinases, and other/atypical protein kinases [2]. The degree of similarity of their kinase domains that enabled such classifications was also thought to preclude meaningful specificity of small molecule inhibitors, especially as most kinase inhibitors tend to interfere with ATP binding as a mechanism of action. The family, as a whole, was initially deemed undruggable. It is testament to the capabilities of synthetic chemists that kinase inhibitors with remarkable selectivity, potency, and specificity have become available. Add to this library the collection of gene knockouts and RNA interference (RNAi) reagents and global analysis of the 500+ gene family become feasible.

The remarkable scope of the kinase inhibitor libraries reflects the high level pharmaceutical interest in this class of proteins due to repeated examples of the dysregulation of these enzymes in human diseases. Here, we initially describe the subclasses of protein kinases, provide examples of their roles in disease, and discuss methods for their characterization and functional assignments.

#### 1.1.1 AGC Kinases

This family of serine/threonine kinases [3] is named after three representative families PKA, PKG, and PKC [4] and contains 60 members with a propensity for phosphorylating Ser/Thr residues preceded by basic amino acids [5]. The master

regulator of the AGC kinases is phosphoinositide-dependent kinase-1 (PDK1), which lies directly upstream of at least 23 other AGC kinase family members including Akt/PKB, protein kinase C (PKC), and serum- and glucocorticoid-induced protein kinase (SGK) and phosphorylates these protein kinases within their catalytic domain at the so-called “T-loop” [3]. Within the cell, PDK1 is constitutively active due to its intrinsic ability to autophosphorylate within its own active site and also contains a C-terminal pleckstrin homology (PH) domain that has high affinity for specific phosphoinositide lipids [3]. Given its wide range of substrates, PDK1 has very diverse cellular functions in regulating metabolism, cell proliferation, and cell survival [6]. PDK1 is inhibited by staurosporine (among other targets) which, given the proclivity of PDK1 to regulate many other kinases, explains why this drug was thought to be relatively nonselective, albeit a potent inducer of cellular apoptosis [7].

PDK1 was discovered by Alessi and colleagues while searching for a protein kinase activity that could phosphorylate and activate Akt/PKB. Akt/PKB is activated by agonists of phosphatidylinositol 3'kinase (PI3K) and, like PDK1, has a PH domain. Stimuli that induce PI3K activity result in generation of 3'-phosphorylate phosphoinositides that act to relocate Akt/PKB and PDK1 to colocalize at the plasma membrane, resulting in Akt/PKB phosphorylation and activation [3]. Full activation also requires phosphorylation of residues in the C-terminal domain of Akt/PKB, known as the hydrophobic motif. The kinase responsible for this modification is mTORC2, the rapamycin-insensitive complex of mammalian target of rapamycin (mTOR), Rictor, and other proteins. Isoform-specific knockouts of the three known isoforms (Akt1, 2, 3) have revealed nonredundant actions of each [8], as Akt1 knockout mice display increased rates of apoptosis [9], while Akt2 knockouts display a phenotype similar to type 2 diabetes with impaired glucose utilization [10]. Akt3 is primarily expressed in neuronal tissue, with Akt3 knockout mice displaying abnormalities in brain development [11]. Akt activation promotes cell proliferation via phosphorylation of substrates involved in regulating apoptosis and the cell cycle. For example, one group of targets, members of the FOXO family, induces transcription of genes that promote apoptosis and cell cycle arrest. Akt phosphorylation promotes exclusion of FOXO from the nucleus, turning down expression of these pro-apoptotic genes [12].

The PKC family consists of at least 12 isoforms, which are divided in four subcategories: conventional ( $\alpha$ ,  $\beta$ , and  $\gamma$ ); novel ( $\delta$ ,  $\epsilon$ ,  $\eta$ , and  $\theta$ ); atypical ( $\zeta$  and  $\iota$ ); as well as PKN- and PKC-related PKN1, PKN2, and PKN3 [3]. While PKC activation is regulated via phosphorylation at a C-terminal hydrophobic motif by mTORC2 and within the “T-loop” by PDK1, PKC activation can also be regulated through lipid second messengers. These vary based on the isoform, with conventional PKCs regulated by both diacylglycerol (DAG) and calcium, novel PKCs regulated solely by DAG, with neither required for atypical PKC activation [3, 13, 14]. PKC activation has two primary consequences; activation of the mitogen-activated protein kinase–extracellular signal regulated kinase (MAPK/ERK) pathway and activation of the PI3K/Akt pathway [15], leading to critical roles in cell survival, growth, and motility.

The SGKs are serine/threonine kinases that were initially found to act, as the name implies, in response to serum, glucocorticoids, cytokines, as well as other growth factors, through phosphorylation by both PDK1 and mTORC2 [13]. SGKs also have additional roles in the regulation of both ion channel and enzyme activity, including the ubiquitin ligase NEDD4, and with the transcription factors beta-catenin and p53. Through its downstream targets, the SGKs play a role in regulation of transport, hormone release, neuroexcitability, inflammation, cell proliferation, and apoptosis [16]. Of note, SGK3 harbors a PX domain that, while related to PH domains, has a distinct binding affinity for 3'-phosphoinositides found in the endosomal compartment [17].

### 1.1.2

#### The CaMK Family

The CaMKs also are basophilic amino acid directed with the majority of kinases in this group initially activated by calcium/calmodulin binding [5], followed by subsequent autophosphorylation. Binding of  $\text{Ca}^{2+}$ /calmodulin induces a conformational change that exposes the activation loop site for phosphorylation by other CaMK kinases [18]. As CaMKs are activated through  $\text{Ca}^{2+}$ , their ultimate functions lie in processes intimately associated with intracellular calcium levels, including excitation–contraction coupling in muscle [19], neuronal development, and synaptic plasticity [18].

Two other well-characterized members of this structurally related family are Myosin Light Chain Kinase (MLCK) and AMP-activated Protein Kinase (AMPK). MLCK is activated following release of calcium from sarcomeres in skeletal muscle [20]. Once activated, MLCK phosphorylates myosin, ensuring cross-bridge formation and subsequent muscle contraction [20]. Activation of AMPK occurs through binding of AMP, resulting in allosteric activation, and phosphorylation on Thr172 by the upstream kinase LKB1 [21]. AMPK acts to promote metabolic pathways that generate ATP: increasing glucose uptake, inactivating acetyl-coA carboxylase (ACC) to inhibit fatty acid synthesis, and enhancing fatty acid oxidation, as well as stimulating mitochondrial biogenesis through either phosphorylation of PGC1- $\alpha$  or activation of SIRT1, leading to inhibition of both protein and rRNA syntheses [22].

### 1.1.3

#### CMGC Family Kinases

The CMGC subfamily of protein kinases is composed of four groups; the cyclin-dependent kinases (Cdks), the mitogen-activated protein kinases (MAPKs), glycogen synthase kinase-3 (GSK-3), and Cdk-like kinases [23]. Cdks are a family of proline-directed serine/threonine protein kinases comprising 21 genes that encode a Cdk, with 5 additional genes encoding Cdk-related kinases. While nomenclature differs, approximately 20 different Cdk have been identified [24]. Cdk activation is typically achieved through binding of cyclin subunits, as in the

classic example of Cdk2 activation via cyclin A association. Other Cdks, such as Cdk4, require additional partners for full activation, as cyclin D binding alone does not confer the active conformation [25–27]. Cdk4 and Cdk6 play roles in promoting entry into the cell cycle and lead to inhibition of the retinoblastoma (Rb) protein. After DNA duplication in S phase, Cdk1 becomes activated promoting chromosome condensation and nuclear envelope breakdown [27]. Cdk5 is an atypical Cdk highly expressed in neural tissues. Its function is less associated with cell cycle regulation [28, 29] and is activated by the non-Cyclin proteins p35 and p39 [28]. Cdk5 functions in cytoskeletal organization, membrane trafficking, and neuronal differentiation [28].

MAPKs are also proline-directed and regulate proliferation, differentiation, apoptosis, survival, inflammation, and innate immunity [30]. The signaling cascades that lead to their activation consist of a hierarchy of kinases, each phosphorylating and activating the next kinase in the pathway. Following a stimulus, the activation of Extracellular signal-Regulated Kinase (ERK) cascade is triggered by the mitogen-activated protein kinase kinase kinases (MAP3Ks), which includes Raf [31, 32]. The MAP3K in turn activates the mitogen-activated protein kinase kinases (MAP2Ks), MEK1/2, leading to dual tyrosine and threonine phosphorylation of ERK [32–34]. Two other pathways that primarily respond to cellular stress stimuli are similarly structured. In these cases, the p38 MAPK and Stress-Activated Protein Kinase/Jun N-terminal Kinase (SAPK/JNK) proteins are activated by the MAP2Ks MKK3 and 6 or MKK4 and 7, respectively. These MKKs are respectively activated by MAP3Ks such as ASK1, MEKK, and MLK3 (mixed-lineage kinase) [32–34]. Each MAPK tends to comprise several isoforms with varying tissue distribution and function, although these typically overlap.

The ERK1/2 MAPK cascade is triggered by extracellular stimuli including growth factors and hormones and regulates cell proliferation and differentiation through control of certain transcription factors including myc, Ets, CREB, Jun, and Fos [32]. ERK5 is usually described separately from the rest of the ERK1/2 pathway as its activation pathway (MAPKKK and MAPKK) differs from that of ERK, p38MAPK, and SAPK/JNK. ERK5 activation is triggered by both cellular osmotic stress and mitogens [35, 36], and downstream substrates include SGK and the transcription factors c-myc, MEF2, and c-fos [37]. p38 MAPK has four isoforms termed  $\alpha$  to  $\delta$ . The  $\alpha$  and  $\beta$  isoforms can be triggered in response to cellular stress, as well as immune responses and inflammation [37]. Downstream targets of p38 MAPK include the MAPKAPKs MNK1/2 and MSK1/2, heat shock proteins, and the transcription factors ATF2 and MEF2C [37]. Similar to p38 MAPK, SAPK/JNK is triggered in response to cellular stresses. Although the three known genes of SAPK/JNK are expressed in most tissues, SAPK $\beta$ /JNK3 expression is primarily found in neuronal cells [30, 38]. Following activation by MKK4/7, SAPK/JNK transmits the signal to a variety of mediators including the MAPKAPK3 and the transcription factors ATF2, c-Jun, and Elk1 [39].

Two genes encoding GSK-3 exist in mammals,  $\alpha$  and  $\beta$ , which share high sequence homology of up to 98% in their catalytic domains [40]. GSK-3 is active

in the basal state and serves to regulate activation of numerous cell signaling pathways including the Wnt, Notch, Hedgehog, and growth factor signaling pathways [42]. Mitogenic pathways activated through stimulation of PI3K induce phosphorylation and inhibition of GSK-3 at serines 21 and 9 in the  $\alpha$  and  $\beta$  isoforms, respectively. This mechanism is irrelevant to regulation of GSK-3 by other signals such as Wnt due to insulation of a small (<10%) of the protein kinase within specific complexes [41]. By this means, the same protein kinase can act downstream of several systems yet be controlled in a highly specific manner. A common theme to many of the substrates of GSK-3 is that their phosphorylation sites fall within “phospho-degrons” that when phosphorylated lead to targeting for ubiquitination and degradation by the 26S proteasome [42]. Other substrates are suppressed in their function (such as glycogen synthase, the rate-limiting enzyme of glycogen synthesis) [42]. GSK-3 might therefore be thought of as a constitutive cellular brake that acts to dampen functions, and its negative effects are selectively switched off by specific signals, permitting induction of a variety of cellular processes. Genetic studies have revealed that despite high homology, GSK-3 $\alpha$  and  $\beta$  have nonredundant functions. GSK-3 $\alpha$  knockout mice are viable, while mice lacking GSK-3 $\beta$  either die late in embryogenesis due to significant liver apoptosis or just after birth due to heart-patterning defects [43–45].

#### 1.1.4

##### **STE Family Kinases**

The STE (for “sterile”) family of kinases contains a variety of atypical kinases including WNK (With No Lysine) and the MAPK cascade kinases [5]. WNK kinases differ from all other kinases as they lack a lysine in  $\beta$  sheet 3, which is instead located in  $\beta$  sheet 2 [46]. These kinases are involved in a variety of signaling pathways where, for example, WNK1 acts as an upstream MAP4K for ERK5. The WNKs themselves can be regulated by Akt/PKB- and SGK-mediated phosphorylation [47]. Recognized functions of WNK include cell proliferation, differentiation, exocytosis, and critical roles in chloride, sodium, and potassium transport [47].

The STE family also includes the MAPK cascade kinases (MEK1/2, MKK3/4/6/7), which transmit the extracellular stimulus signal to the MAPK, leading to alterations in cell proliferation and survival. The STE20 subfamily consists of MST, PAKA, and PAKB and serve as MAP4Ks, which transmit signals to the STE11 (MAP3K) family of which ASK1 and MEKK2 are two examples [48].

#### 1.1.5

##### **Tyrosine Kinases**

This family consists of two distinct classes: membrane-bound receptor tyrosine kinases (RTKs) and non-RTKs. RTKs are transmembrane proteins harboring an extracellular domain that contains high-affinity binding sites for ligands and a

C-terminal, intracellular domain which includes a protein kinase domain with specificity for tyrosine. Examples include many growth factor receptors including those for platelet-derived growth factor, epidermal growth factor, colony stimulating factor 1, insulin, insulin-like growth factor, heregulins, vascular endothelial growth factor, angiopoietins, nerve growth factor, and so on [49]. This family is most targeted pharmaceutically due to the identified roles of these enzymes in many forms of cancer (such as breast, lung, stomach) as well as angiogenesis (VEGF). Therapeutic agents targeting RTKs have flourished in the past decade and are either monoclonal antibodies to the extracellular domain (e.g., Herceptin) or small molecule inhibitors to the intracellular tyrosine kinase domain (e.g., Tarceva, Iressa) [50]. The substrates of tyrosine kinases tend to be primarily their dimeric partners, which are phosphorylated at multiple tyrosines lying within a C-terminal tail domain. The phosphorylated residues, within local amino acid sequence contexts, recruit proteins harboring Src Homology 2 (SH2) or PhosphoTyrosine Binding (PTB) domains that include the regulatory subunits of PI3K, certain protein tyrosine phosphatases, phospholipase  $C\gamma$ , and adaptor proteins such as Grb2, SHC, Gab1, and the insulin receptor substrate (IRS) proteins, which, in turn, recruit other signaling proteins [51].

In comparison, non-RTKs are typically intracellular proteins and can also associate with membranes to serve as catalytic domain donors to RTK that lack such a domain [52]. This family includes the three Src-like kinases (Src, Fyn, and Yes) as well as Lck, which lies downstream of the T-cell receptor and c-Abl, the infamous kinase that as a result of the 9:22 translocation associated with the Philadelphia chromosome in chronic myeloid leukemia (CML) is oncogenically activated (Bcr-Abl) [53, 54]. This latter kinase is also the target for the poster-child of molecularly targeted anticancer agents, Gleevec/imatinib, which has had remarkable impact in the clinical treatment of CML. Indeed, Gleevec is so effective in blocking the activity of Bcr-Abl that patients in whom the disease returns virtually exhibit mutations in the tyrosine kinase that interfere with Gleevec binding. This has given rise to second-line tyrosine kinase inhibitors designed to target common resistance mutations such as nilotinib [55]. In spite of their name, many of the non-RTKs act in close association with RTKs.

#### 1.1.6

##### Casein Kinases

There are two primary subclasses of casein kinases. Casein kinase 1 (CK1) consists of eight genes in mice, which function as monomeric enzymes:  $\alpha$ ,  $\gamma 1$ ,  $\gamma 2$ ,  $\gamma 3$ ,  $\delta$ ,  $\epsilon 1$ ,  $\epsilon 2$ , and  $\epsilon 3$ . CK1 is basally active and serves as a priming kinase for other protein kinases such as GSK-3 (in the case of  $\beta$ -catenin and LRP5/6) and an upstream regulator of Cdk5 [56–58]. These are under-appreciated kinases that touch upon many cellular processes through selective association with macromolecular complexes. Like its distally related cousins, the casein kinase II subfamily is implicated in a wide panoply of cellular events, largely via basal activity although this kinase has also been implicated in animal tumors. Notably, neither subclass is a

physiological casein kinase. That role is owned by Fam20C, a protein without obvious structural similarity to protein kinases [59].

#### 1.1.7

##### **Tyrosine Kinase-Like Family**

This is a diverse group containing both serine and tyrosine protein kinases that includes MLK, LISK, IRAK, Raf, RIPK, and STRK [2]. Despite its name, this family is strictly comprised of ser/thr specific kinases that act as intermediaries in a number of signaling cascades. Their structure suggests that they may have been evolutionary stepping stones toward tyrosine kinases (the yeast genomes are effectively barren of “true” tyrosine kinases, although Wee1 targets Cdc2 at Tyrosine 15 [60]).

#### 1.1.8

##### **RGC Kinases**

The RGCs is an atypical family with a domain sequence similar to tyrosine kinases that possess enzymatic capabilities [2]. Members of this family are activated by hormones, peptides, and low calcium-induced guanylyl cyclase-activating proteins [61]. They possess an active guanylate cyclase domain, which generates cGMP, and a catalytically inactive kinase domain [62]. RGC have a wide array of substrates and have roles in hypertension, cardiac hypertrophy, and vision [63–65].

#### 1.1.9

##### **Atypical/Other Protein Kinases**

This subgroup includes kinases that lack primary sequence similarity to the canonical protein kinase profile, but have experimentally demonstrated protein kinase activity. The subgroups include Alpha Kinases, ABC1, pyruvate dehydrogenase kinases, RIO, Bromo Domain kinases (BRDs), TATA binding factor associated factors (TAFs), BCR, FASTK, G11, Transcriptional Intermediary Factor 1 family (TIF1), H11, Fam20C, and PIKK, a member of the PI3K-related kinases [2]. Of this diverse group, one PIKK in particular has been well studied, FRAP, better known as mTOR.

mTOR is an atypical serine/threonine kinase [13] involved in the regulation of cell growth, proliferation, and metabolism and is associated with two distinct complexes termed mTORC1 and mTORC2. In mTORC1, mTOR interacts with Raptor, Deptor, and GβL to regulate protein translation and metabolism through phosphorylation of downstream targets including 4E-BP1, S6K, SREBP [13]. In the mTORC2 complex, mTOR interacts with Rictor, GβL, Sin1, Deptor, and PRR5 and is activated in response to extracellular cues. It remains to be elucidated via what molecular mechanisms these cues are “sensed,” though mTORC2 carries out its role via phosphorylation of substrate kinases in the AGC family [13, 66].

## 1.2

### Why Study Protein Kinases – Their Roles in Disease

After the sequencing of the human kinome was completed, the kinase chromosomal map was compared to disease loci, revealing 164 kinases that map to sites frequently observed in tumors [2, 67], with 80 kinases mapping to other (non-cancer) disease loci [2]. The remarkably high proportion of kinases associated with such loci highlights the important role of protein kinases in disease initiation and progression. Below we exemplify the role of key protein kinases in the pathophysiology of just two of the chronic diseases that affect tens of millions of people worldwide: Alzheimer's disease (AD) and cancer.

#### 1.2.1

##### Neurodegenerative Disease

AD is a progressive neurodegenerative disorder characterized by both the loss of acetylcholine and neuronal overstimulation, leading to neuronal decay and impairments in both cognition and behavior [68]. The pathological hallmarks of AD progression include protein deposits found inside (neurofibrillary tangles – NFTs) and outside (amyloid beta – A $\beta$  plaques) of the neuron [68]. In healthy individuals, the processing of amyloid precursor protein (APP) is regulated by  $\alpha$ -secretase, which cleaves a 695 amino acid fragment, which is further processed by a second enzyme,  $\gamma$ -secretase. This is described as the nonamyloidogenic pathway, as it does not produce the toxic A $\beta$  peptides. The production of amyloid plaques in Alzheimer's patients is regulated by the amyloidogenic pathway in which A $\beta$  peptides are generated through increased APP cleavage by  $\beta$ -secretase. The cleavage of APP by  $\beta$ -secretase releases a longer fragment that is subsequently cleaved by  $\gamma$ -secretase. APP cleavage initiated by  $\beta$ -secretase results in the formation of peptides of varying length including A $\beta$ 40 and A $\beta$ 42. These two peptides differ in their effects on neighboring neurons as A $\beta$ 42 peptides have neurotoxic properties, at least *in vitro* [69]. Recent studies have suggested that it is the relative ratio of these two peptides, and not their absolute quantities, which is key in the induction of neurotoxicity [70, 71].

The microtubule-associated protein Tau is important for the assembly and stabilization of neurofilaments, which play key roles in axonal growth and transport [72]. In healthy individuals, Tau phosphorylation typically occurs to target the protein to various compartments and is essential for its mobility within the neuron [72]. In AD, hyperphosphorylation of Tau leads to its dissociation from the microtubule and the formation of an insoluble complex of paired helical fragments. These fragments then aggregate together inside the neuron, leading to the formation of NFT [73, 74], resulting in decreased transport of essential components within the neuron and the eventual loss of the neuron.

Protein kinases have well-documented roles in the progression of AD and in the two pathological processes that are characteristic of the disease. The CMGC group of protein kinases have been of particular focus due to the significant

consequences of alterations in GSK-3, neuronal-enriched Cdk5, and MAPKs in the brain. Increased GSK-3 activity has been documented in postmortem analysis of the brain of suspected Alzheimer's patients through increased protein expression and phosphorylation at tyrosine sites linked to increased activity [75–77]. Studies have connected GSK-3 expression to both amyloid- and tau-associated pathologies in AD [78–80]. Treatment of rat hippocampal neurons in culture with A $\beta$  peptides increases GSK-3 activity [81], and GSK-3 $\beta$  was specifically shown to regulate processing of APP by  $\beta$ -secretase through NF- $\kappa$ B-induced BACE1 promoter activity and gene expression both in culture and in mouse models [82]. Tyrosine-phosphorylated GSK-3 also has been shown to colocalize with NFT [77], and GSK-3 acts as a physiological tau protein kinase, playing a significant role in the hyperphosphorylated state of tau observed in brain slices from AD-afflicted individuals [83].

The roles of GSK-3 in neurobehavior and neuronal health have been assessed in mice with altered GSK-3 expression in various brain regions. Transgenic overexpression of GSK-3 $\beta$  in neurons of the cortex and hippocampus resulted in tau hyperphosphorylation, neuronal death, and reduced spatial learning and memory, as assessed by the Morris water maze test [78, 79, 84]. The tau pathology was linked to the effects on spatial learning and memory, as performance on the Morris water maze test improved through tau deletion in these mice [85]. A conditional mouse model was later used to demonstrate the effect of both increased GSK-3 activity on forebrain neurons and the consequence of lowering activity back to normal levels. Increased GSK-3 activity leads to hyperphosphorylation of tau, neuronal loss, and deficits in spatial learning. Following reduction of GSK-3 activity, phosphorylation of tau decreased, neuronal loss was diminished, and deficits in spatial learning were prevented [86]. Also, inhibition of GSK-3 activity, either through pharmacological or genetic methods, in mouse models of AD had neuroprotective effects, with decreased A $\beta$  production and reduced hyperphosphorylation of tau observed [87]. Treatment of rats with lithium, a relatively nonspecific inhibitor of GSK-3 [88, 89], before and after infusion of A $\beta$  fibrils was shown to prevent spatial memory deficits, as well as tau hyperphosphorylation [90, 91]. In the TgCRND8 AD mouse model, lithium treatment also rescued deficits in spatial learning and memory [92, 93]; however, studies on the effects of lithium on AD and cognition have not all shown benefit. Chronic lithium treatment begun after amyloid plaque development in TgCRND8 mice showed no improvement in cognition, even though the average size of the A $\beta$  plaques was reduced [93]. Also, treatment of early stage AD patients with lithium showed no improvement in cognition [94]. Currently, a series of small molecule inhibitors are being developed and tested to target GSK-3, and recent studies have shown cognitive improvements and reduced tau phosphorylation and neuron degeneration in mouse models of AD [92, 95, 96].

Cdk5 is a unique Cdk, as it is highly expressed in postmitotic neurons [97], and in the adult brain it regulates neuronal survival, synaptic plasticity, and in the formation of learning and memory [97]. In AD, truncation of the Cdk5-regulating protein p35 occurs in response to A $\beta$ -induced activation of calcium-dependent proteases, the calpains, chopping down p35 to p25 [97]. As the p25 form of the

activating subunit is resistant to ubiquitin-mediated proteolysis, it extends the activation time of Cdk5, leading to hyperphosphorylation of its substrates, including tau [68, 74, 98]. The phosphorylation of tau by Cdk5 also serves to prime it for further phosphorylation by GSK-3, heightening the level of tau hyperphosphorylation [99]. Mice with regulatable expression of p25 displayed brain atrophy, synaptic dysfunction, and NFT formation [100]. Knockdown of Cdk5 expression in a mouse model of AD reduced tau phosphorylation as well as the accumulation of NFT and rescued hippocampal neuronal loss [101]. The development of Cdk5-specific inhibitors has been hampered by the high sequence homology between the various Cdk isoforms. A recently described Cdk5-specific inhibitor was shown to decrease A $\beta$  production in both young and old APP mutant mice, but in young mice it also relieved inhibition of GSK-3, leading to increased hyperphosphorylation of tau [102].

All three MAPK family members have documented roles in AD pathophysiology, in part, through phosphorylation of tau at various sites [103, 104]. Both the ERK and SAPK/JNK pathways play a role in A $\beta$ -induced neurotoxicity as exposure of hippocampal neuronal slices to A $\beta$  peptides induced cell death via ERK-induced caspase-3 expression [105] and exposure of cortical neurons to A $\beta$  peptides activated SAPK/JNK leading to Fas-dependent neuronal death [106]. Infusion of a MEK inhibitor into mice displaying A $\beta$ - and tau-induced pathology and behavioral deficits reversed the decline in memory displayed by these animals [107]. In response to oxidative stress, SAPK/JNK activation also has been shown to increase phosphorylation of APP, which promotes generation of A $\beta$  peptides [108]. The A $\beta$ 42 peptide has specifically been shown to induce SAPK/JNK-mediated BACE1 expression, further leading to increased A $\beta$  peptide generation [109]. This effect of SAPK/JNK on A $\beta$  generation has been confirmed *in vivo*, where SAPK/JNK inhibition decreased A $\beta$  peptide generation and plaque deposition in a mouse model of AD [110]. SAPK/JNK activation also has been demonstrated in the synapse prior to the establishment of cognitive impairments and treatment of TgCRND8 mice, a model of AD, with a jun N-terminal kinase (JNK) inhibitor prevented both the loss of postsynaptic proteins and reduction in excitatory synapse size, preventing neural spine degeneration [30, 111].

Activation of the p38 MAPK pathway in AD has also been reported to play a significant role in pathophysiology of the disease [112–115]. Patients in the early stages of AD show evidence of p38 MAPK pathway activation in both their neurons and glia, which has been linked to glial proinflammatory cytokine production and synaptic dysfunction [112–114, 116]. p38 MAPK may also contribute to tau phosphorylation, leading to its dissociation from microtubules and aggregation into NFT [83, 117]. The role of p38 MAPK in AD-associated neuroinflammation was highlighted in recent studies, where p38 MAPK inhibitor treatment decreased both A $\beta$  plaque accumulation and A $\beta$ -induced production of inflammatory regulators IL-6 and TNF- $\alpha$  [30, 118]. A novel p38alpha MAPK inhibitor, MW150, has been tested for efficacy on memory-related deficits associated with AD in two mouse models of AD: one overexpressing APP, leading to rapid onset of symptoms, and one utilizing endogenous promoters with no overexpression and slow

progression. In both models, treatment with the p38 $\alpha$  MAPK inhibitor halted AD-associated deficits in memory, as assessed by the radial arm water maze test and the contextual fear conditioning test [119].

Several members of the AGC kinase family also have described functions in the progression of AD. Under normal physiological conditions, mTOR plays a role in tau protein synthesis and phosphorylation via activation of p70S6K [120]. In contrast, it has recently been reported that A $\beta$  accumulation inhibits the mTOR pathway and that upregulation of mTOR signaling prevents A $\beta$ -induced deficits in long-term potentiation [121, 122]. In addition, treatment of rats with rapamycin, a potent and selective mTOR inhibitor, was shown to enhance the neurotoxicity of A $\beta$  peptides [123]. Like mTOR, PKC isoforms appear to have contrasting roles with other kinases in AD. Activation of PKC isoforms stimulates APP cleavage by  $\alpha$ -secretase, resulting in decreased A $\beta$  peptide production, both in culture and in mouse models of AD [124–126]. Also, A $\beta$  has been shown by various researchers to inhibit PKC activation via multiple mechanisms, including direct binding via a putative PKC pseudosubstrate domain [127, 128]. Recent studies have shown that human PKC $\epsilon$  overexpression reduces A $\beta$  expression and increases expression of endothelin-converting enzyme type 1, which plays a role in A $\beta$  clearance [129, 130].

Overall, these studies have revealed the integral role for several protein kinases in the progression of AD, and future work needs to focus both on further increasing our understanding of the mechanisms of kinase action in the progression of this disease, as well as developing novel therapies to target these kinases to halt disease progression. A substantial challenge in the field is that onset of symptoms usually occurs long after neuronal loss has occurred due to the remarkable synaptic plasticity and neuronal rewiring. As a consequence, effective therapies are likely going to need to be used prior to presentation of symptoms, raising the bar for safety of any prophylactic intervention.

### 1.2.2

#### Hallmarks of Cancer

Cancer is the result of a complex series of events that lead to uncontrolled growth and spread of tumor cells. Hanahan and Weinberg described the six hallmarks of cancer, which include sustaining a proliferative signal, evading growth suppressors, activation of invasion and metastasis, resisting cell death, inducing angiogenesis, and enabling replicative immortality [131]. Protein kinases have documented roles in each of these cancer hallmarks, and therefore extensive study is ongoing to assess their role in cancer progression and to ascertain whether their inhibition would be a useful therapeutic approach.

The initiation of disruptions to normal growth restrictions within a cell typically occurs via mutation of critical signaling components. Mutations in protein kinase signaling pathways account for a high proportion of the total amount of driver mutations found in human cancers. Point mutations in the PIK3CA gene (hotspots include E542K, E545K, and H1047R), coding for the p110 $\alpha$  subunit of

PI3K, are among the most frequent in human cancer and all result in enhanced catalytic activity of PI3K [132, 133]. Aside from gene mutation, activation of PI3K, and its downstream target Akt/PKB, are observed in breast, ovarian, pancreatic, esophageal, and thyroid cancer due to gene amplification [132]. In mouse models, expression of a single PIK3CA point mutation targeted to the mammary gland was sufficient for mammary tumor growth [134–137]. In contrast, PIK3CA point mutation expression in mouse models of colorectal, ovarian, endometrial, and lung cancer required a separate cooperating mutation in another pathway in order for tumor growth to occur [138–141]. Constitutively active Akt/PKB isoforms do not induce mammary gland tumorigenesis when expressed via the MMTV promoter [142, 143], indicating that overexpression of Akt/PKB alone is insufficient for mammary tumorigenesis. Following this result, the role of the Akt/PKB isoforms was examined in the context of mouse models of breast cancer driven by PyMT or ErbB-2 [8]. Constitutively active Akt1/PKB $\alpha$  was shown to accelerate tumor growth in the ErbB-2 mouse model of breast cancer, with tumors detectable at earlier time points but with less metastasis observed. In contrast, constitutively active Akt2/PKB $\beta$  had no effect on tumor latency but metastasis to the lung was increased [144, 145]. Akt1/PKB $\alpha$  and Akt2/PKB $\beta$  knockouts crossed to the PyMT and ErbB-2 mouse mammary tumorigenesis models also revealed opposing effects with loss of Akt1/PKB $\alpha$  resulting in delayed tumor onset, while loss of Akt2/PKB $\beta$  accelerated tumor growth [8, 146].

Due to its high mutation and amplification rate in human cancer, PI3K represents an important node for drug discovery research. Currently, numerous pan-PI3K and dual PI3K/mTOR inhibitors are at varying stages of clinical trials for treatment of breast cancer, nonsmall cell lung cancer, and renal cell carcinoma [147]. To date only one PI3K inhibitor has received regulatory approval. Idelalisib, a specific PI3K $\delta$  inhibitor, was approved in mid-2014 for combination treatment of patients with B cell malignancies [148, 149].

mTOR lies downstream of both the Ras/MAPK and PI3K pathways, where mutations in oncogenes or tumor suppressors are found in a high percentage of human cancers, leading to constitutive activation of mTORC1 and resultant tumor growth [150, 151]. The mTORC1 inhibitors rapamycin and rapalogs Temsirolimus and Everolimus are approved for the treatment of renal cell carcinoma, hormone refractory breast cancer, and pancreatic neuroendocrine tumors [152–154]; however, effects in patients have been modest as while these inhibitors affect cell proliferation, they are not toxic to tumor cells [151]. Also, the mTORC1 plays a role in feedback loops to repress signaling through RTKs, and inhibition of the complex allows this signaling system to remain unchecked [151, 155].

Activating mutations in PKC isoforms in cancer are rare [15], while PKC isoforms show altered expression in various cancers including PKC $\epsilon$  in stomach, lung, and breast cancer, PKC $\theta$  in GIST, and PKC $\eta$  in nonsmall cell lung cancer [14]. Using selective isoform inhibitors in mouse models of breast cancer, it has been shown that augmenting PKC $\alpha$  expression abrogates metastasis without an effect on primary tumor growth [156]. The role of PKC $\beta$  in tumor progression is

somewhat controversial as inhibitor studies have shown growth suppression in mouse xenograft models, whereas overexpression in mouse mammary cell lines has revealed suppression of both tumor growth and metastasis [15, 157, 158]. The novel PKC $\epsilon$  isoform is considered oncogenic as early studies revealed that overexpression of PKC $\epsilon$  in NIH 3T3 cells injected into mice induced tumor growth, while untransfected and PKC $\delta$  overexpressing cells failed to grow [159, 160]. Knockdown of PKC $\epsilon$  in MDA-MB-231 breast cancer cells has been shown to significantly reduce both primary mammary tumor growth and metastasis to the lung [161]. PKC $\zeta$  overexpression in cancer has been linked to the amplification of its gene *PRKCI*, which is located on chromosome 3q26, one of the most commonly amplified regions in human cancer [162], with copy number gains established in lung squamous cell carcinoma and serous epithelial ovarian cancer [162]. PKC $\zeta$  knockdown in esophageal cancer xenografts caused decreases in tumor growth and metastasis to the lung [163].

The CMGC group of protein kinases also have well-documented roles in cancer progression and serve as drug discovery targets for new therapeutics. Melanoma, thyroid cancer, and colon cancer all show evidence of alterations in the MAPK pathway, with this activation, in many cases, the result of activating point mutations in RAS family members (HRAS, KRAS, and NRAS) [164, 165]. Trametinib is the first MEK inhibitor approved for cancer treatment [164], and recent studies have shown that combination with B-Raf inhibitors results in enhanced antitumor activity in patients with melanoma [166]. The p38 $\alpha$  MAPK isoform has also been shown in preclinical models to play a role in resistance to various chemotherapeutic drugs for the treatment of colon cancer, as well as to kinase inhibitor treatment of hepatocellular carcinoma [30, 167, 168]. Overactivation of SAPK/JNK has been observed in hepatocellular carcinoma and mouse models of this disease in which JNK1 has been inactivated show decreases in cell proliferation and tumor formation [169, 170]. JNK2 is also implicated in tumor progression, as ablation of this kinase blocks chemical-induced skin carcinogenesis in the mouse [171].

The transcriptional regulator Snail plays an important role in epithelial–mesenchymal transition (EMT) of various solid tumors including gastric, liver, and colorectal cancer [172–174]. This protein is rapidly turned over by ubiquitin-mediated degradation, triggered by phosphorylation by GSK-3. If GSK-3 activity is inhibited, Snail protein accumulates and represses transcription of the epithelial cell adhesion protein, E-Cadherin (as well as promoting mesenchymal gene expression). GSK-3 overexpression has been observed in colon, liver, ovarian, and pancreatic tumors [175], while inhibition of GSK-3 $\beta$  activity promoted development of hepatocellular carcinoma [176], as well as stabilization of  $\beta$ -catenin and mammary gland hyperplasia [177]. Knockdown of GSK-3 was shown to inhibit pancreatic cancer growth and angiogenesis [175, 178] associated with decreased Bcl-2 and VEGF expression. Studies have also highlighted GSK-3 as a tumor suppressor as the overexpression of a constitutively active form of GSK-3 lead to increased chemosensitivity, cell cycle arrest, and reduced tumorigenicity of breast cancers [175]. GSK-3 $\beta$  has also been identified in several “sleeping beauty” mediated screens for tumor suppressors [179, 180].

As observed in these studies, as well as many others we do not have space to include protein kinases play a complex role in the initiation and progression of cancer. Using our increased understanding of the role of kinases in these processes, coupled with remarkable advances in medicinal chemistry that is increasing selectivity and potency of kinase inhibitions, should allow for improved and precise therapeutics to help conquer kinase-inhibitor drug resistance typically observed in drugs currently approved for therapeutic use.

### 1.3

#### Methodology for Assessment of Protein Kinase Functions

While AD and the many forms of cancer clearly impose huge burdens on patients and healthcare, there are many other conditions for which modulation of protein kinase activities offer promising potential including diabetes, arthritis/autoimmunity, viral infections, and psychiatric disorders. That said, there are significant hurdles in targeting these cellular regulators. It is critical to remember that signaling pathways play many distinct roles in distinct tissues and organs in response to a wide array of stimuli. Protein kinases are the cells “mail delivery system” and most drugs are indiscriminate in terms of where they act. Hence, therapeutic options for targeting members of this class of signaling proteins must take into account the possible impact on other areas of the body where these signaling molecules are essential for providing normal cellular homeostasis. A first step in delineating physiological functions of these pleiotropic enzymes is measurement of their normal functions and how these are perturbed – a task assisted by new technologies that are relatively unbiased.

#### 1.3.1

##### Mass Spectrometry

Protein mass spectrometry (MS) monitors the mass-to-charge ratio of ionized peptides followed by further fragmentation of the peptides to obtain the mass-to-charge ratio of the components producing a spectrum that identifies unequivocally not only the peptide but also any posttranslational modifications including phosphorylation [181]. As analysis of MS data accumulated, it became clear that previously identified phosphorylation sites represented a tiny portion of those actually present when assessed in an unbiased manner. Databases such as PhosphoELM (<http://phospho.elm.eu.org>) provide searchable interfaces for probing all phosphorylation, as well as other modifications, identified to date, by protein. This avalanche of new data uncovered a major gap in understanding and raised important questions. Firstly, because a phosphorylation site exists, does that mean it is physiologically relevant? Secondly, what are the kinases and phosphatases acting on these sites? Several approaches have been developed to try to answer these questions. One, termed NetworkKIN (part of a suite of kinase bioinformatics tools from the University of Copenhagen: <http://kinomexplorer.info>), uses regular

expression analysis of PubMed abstracts along with databases such as PhosphoELM and known primary sequence preferences to predict the kinases responsible for modification of any given phosphorylated residue. This type of information can play a vital role in assessing the likely importance of a site and provides insight into experimental approaches for validation. Other approaches have mapped the changes observed in response to treatment of cells with protein kinase inhibitors. MS analysis of the impact of the broad-spectrum kinase inhibitor CTx-0294885 revealed 235 protein kinases that were targetable with this inhibitor in the human breast cancer cell line MDA-MB-231 [182]. Of note, this screen revealed the interaction of this broad-spectrum inhibitor with all Akt/PKB isoforms as well as previously unreported phosphosites on HIPK2, a CMGC protein kinase and MELK, a CaMK [182].

Affinity-purification mass spectrometry (AP-MS) combines the specificity of antibody-based affinity purification with the sensitivity of MS. A typical work flow involves immobilizing a ligand, usually a well-characterized antibody, on a solid support such as an agarose or magnetic bead, allowing for capture of a target protein or protein complex from the soluble phase [183, 184]. Advantages of AP-MS include scalability, as it can be performed on as many proteins as can be epitope-tagged and/or antibodies that are available for immunoprecipitation, the fact that it can be performed on proteins in their native cellular environment and that it can monitor dynamic changes in protein–protein interactions [181]. Following AP-MS, it is possible to integrate the information on disease loci with protein interaction data to gain a new understanding of how kinases may control disease pathways. In a study by Varjosalo *et al.*, AP-MS data was used to identify 91 disease-associated proteins that physically interacted with CMGC kinases, with cancer being the most common disease associated with the CMGC interaction proteome. As an example, CDK6 was found to interact with cyclin D2, cyclin D3, and the CDK inhibitor p18, which are linked to various tumor types including non-Hodgkin lymphomas and glioma. These results served to confirm previously obtained results as well as showcase new potential mediators as CLK2 was shown to form interactions with SETD2, a methyltransferase that is linked to renal carcinoma [23, 185].

Another study used AP-MS to identify interacting partners with three critical proteins in colorectal cancer, namely adenomatous polyposis coli (APC), a tumor suppressor protein that is mutated in over 80% of colon cancers, MRE11A, a component of the MRE11A-RAD50-NBN complex, involved in the response to DNA damage and DNMT1, a DNA methyltransferase involved in the regulation of DNA methylation. AP-MS confirmed previously identified binding partners, validating the system, and also combined subcellular fractionation to localize protein–protein interactions to specific cell compartments [184]. In particular, although levels of APC and Flightless-1 homolog (FLII) were higher in the cytosol compared with the nucleus, MS revealed that their interaction only occurred in the nucleus, which is in line with previously published reports of FLII associated with  $\beta$ -catenin in the nucleus [186, 187]. Although this study did not examine protein kinases in particular, the technique could be applied to kinase interactions

that are critical within a particular disease model and the subcellular compartment in which they take place – an important element of cellular compartmentalization and signaling specificity.

Liquid chromatography tandem mass spectrometry (LC-MS/MS) is a further MS technique useful in the study of protein kinases in disease as it allows protein identification within mixed solutions. Recent studies have employed this technique to search for novel liquid biopsy markers in AD. Analysis of plasma from Tau transgenic mice and control mice revealed that changes in protein levels could be detected in the plasma, even during early stages of disease progression. This study confirmed three potential biomarkers in Histone H2B type 1-F/J/L, Clusterin, and Transthyretin, which were increased in the plasma of tau mice. Adenosine kinase was also decreased in the plasma of Tau transgenic mice during the early stages of disease progression, which was linked to the neurotoxicity observed in AD, as decreased adenosine kinase activity can lead to adenosine accumulation within the cell [188].

A recent innovation variation of traditional MS is MIB/MS, in which beads containing multiple kinase inhibitors are used to capture any kinases capable of interacting with the inhibitor, with the identity of the captured kinases determined through normal MS workflow. This technique can provide a snapshot of the signaling occurring in a particular tumor or tissue and can also be used to assess changes in kinase activity following drug treatment, for example, revealing potential drug resistance within a given tumor [189]. In response to MEK inhibition, the kinase activation profile is altered within 24 h, leading to the reactivation of the MEK/MAPK pathway within a few days [190]. This technique has also been used to compare kinase activation profiles between leukemia cells lines that were sensitive or resistant to imatinib treatment. MIB/MS revealed 35 significant kinases that are altered in imatinib-resistant cells including increased activity of MEK and ERK, while activity of Kit and SAPK/JNK were decreased [191]. These data allowed for relevant drug combinations to be tested in human cancer cell lines, revealing that cotargeting with inhibitors of MEK or IKK $\alpha$ , resulted in decreased cell viability and increased apoptosis – offering potential new treatments following resistance to the original therapy [191].

### 1.3.2

#### Fluorescence Resonance Energy Transfer

Fluorescence resonance energy transfer (FRET) is a technique for assessing direct molecular interactions between two molecules in their natural state (i.e., within cells). FRET consists of energy transfer from an excited donor fluorophore to a proximal acceptor fluorophore due to overlapping emittance and absorbance spectra. The energy transfer alters the amount and wavelength of the emitted energy, which can be quantified. This technique allows determination of how specific molecules interact as well as how this interaction may be altered during drug treatment [192]. When measured by fluorescence lifetime imaging microscopy (FLIM), FRET can be used to track kinase activity in fixed cells as well

as tissue samples. Analysis of mean fluorescence lifetime images revealed that while there is a small degree of PKC $\alpha$  activation in unstimulated cells, treatment with TPA, a PKC activator, rapidly increased PKC $\alpha$  phosphorylation/activation, validating the technique in the assessment of kinase activity. This activation could then be tracked to differing regions of the cell with activated PKC relocating to the perinuclear compartment [193]. FRET has also been used to ascertain novel binding partners of mTOR and exactly which proteins directly interacted within the mTOR complex. Use of GFP-mTOR and DsRED-Rheb, a GTP-binding protein, revealed direct interaction between these two proteins, which had been postulated via immunoprecipitation experiments. Through the use of these biosensors, it was also determined that Raptor and mTOR directly interact but Raptor and Rheb do not [194].

FRET can also be performed on formalin-fixed breast tumor sections, and this analysis revealed that only about half of the examined tumors showed increased PKC $\alpha$  activity, while the rest showed increases in expression with no effect on activity [193]. Studies on primary breast cancer tissue assays also have examined the utility of FRET to assess activity of a given enzyme in relation to prognosis, as current methods rely on staining intensity measurements by independent observers. FRET sensors monitoring Akt/PKB phosphorylation at Thr308 revealed heterogeneous Akt/PKB activity in breast tumors although higher phospho-Thr308 levels were correlated with worse disease-free survival [195].

This technique can also be used to identify novel binding partners of a particular kinase under a certain set of conditions or confirm interaction results obtained via high-throughput screens. Previous results had revealed that the AGC kinase ROCK1 activated LIMK2, and this interaction was hypothesized to be critical in membrane blebbing-mediated cell motility. FRET-FLIM allowed the interaction between ROCK and LIMK to be assessed in cells with spread morphology and those displaying membrane blebs. No FRET was detected between GFP-ROCK1 and mRFP-LIMK2 in blebbing cells, while the interaction was confirmed in cells with a spread morphology [196]. Also, time-resolved-fluorescence resonance energy transfer (TR-FRET) was used to assess receptor heterodimerization in response to drug treatment. TR-FRET combines the standard FRET technique with time-resolved (TR) measurement of fluorescence, involving the delayed measurement of fluorescence following excitation, in order to reduce the inclusion of short-lived, nonspecific emissions, to only capture long-lived fluorescence engaged in a FRET process [197]. TR-FRET examining EGFR-HER2 heterodimer formation in response to combination therapy drug treatment revealed that treatment with two monoclonal antibodies, cetuximab (EGFR inhibitor), and trastuzumab (HER2 inhibitor) resulted in significantly reduced pancreatic xenograft tumor growth, in part, due to significantly reduced heterodimer formation [198].

FRET is also being used in investigations of the mechanisms of drug resistance and the effectiveness of second-generation therapies. Mizutani *et al.* tested two second-generation therapies on imatinib-resistant cell lines. The biosensor used enabled the evaluation of BCR-ABL kinase activity in living cells and utilized

the downstream substrate of BCR–ABL, CrkL. In this study, FRET efficiency increased up to 40% in the presence of the BCR–ABL fusion, which could be decreased with imatinib treatment. In cell lines expressing the BCR–ABL fusion, these new inhibitors showed faster kinetics compared with imatinib; however, in imatinib-resistant cell lines, some sensitivity was observed by decreased FRET efficiencies with one of the drugs, while no change was detected in FRET in the other cell line tested with both second-generation inhibitors [199].

FRET techniques allow for the study of protein kinase activity in a variety of disease-related contexts. A resource for further examples on the use of FRET/fluorescence biosensors in probing kinase function and drug discovery can be found in a recent compilation [200].

### 1.3.3

#### **Assessment of Kinase Functions *in vitro*: Genetic and Chemical**

Knockdown of kinase expression in cell lines has been instrumental in establishing the complex roles of protein kinases in cellular function in both the normal and disease states. Establishing kinase knockdown in cells has relied on two main technologies: RNAi and chemical inhibitors of kinase function. RNAi relies on endogenous cellular processes, which following introduction of double-stranded (ds) RNA, posttranscriptionally silence gene expression [201]. It requires the introduction of target-specific dsRNA (short hairpin RNA (shRNA)) into the cell of interest through viral or nonviral methods. Once in the cell, the dsRNA is processed into short RNA sequences of approximately 21–23 nucleotides in length by the endogenous DICER protein and integrated into the RNA-induced silencing complex (RISC). In this complex, the antisense strand binds to mRNA with the complementary sequence, leading to mRNA cleavage and degradation, resulting in protein loss [202]. Given the critical role protein kinases play in disease progression, intense focus has been on the development of chemical inhibitors for the treatment of disease. The first kinase inhibitor was approved for use in 2001 and to date well over 20 have received approval, primarily for the treatment of cancer [203]. Chemical suppression of kinase function is elicited via two broad types of inhibitors: those which compete with ATP for active site binding, and non-ATP-based small molecule inhibitors, which target the site of substrate binding [203].

Part of the power of these two methods of kinase knockdown lies in the fact that they can be used in large screens: for the assessment of all kinases in a particular signaling pathway, cellular function, or disease state. The two methods are often used in parallel in order to overcome specific limitations associated with each. The use of genome/kinase wide screens requires careful determination of what read out will provide the greatest amount of information on kinase function. These screens can assess both negative and positive functionality for a given read-out, such as cell viability. Negative screens can identify kinases required for survival such that their suppression results in cell death, and therefore the causative shRNA, dropping out of the final cell population. Positive selection screens can

use cell viability to assess the development of therapeutic resistance with knock-down of kinases required for drug sensitivity resulting in their enrichment within the final cell population. Below we highlight the use of both RNAi- and inhibitor-based screens in the assessment of protein kinase function.

The readout for many screens is cell viability; however, altered cell state (differentiation) and pathway activation can also be used to assess kinase function in genome/kinase wide screens. In melanoma cancer cell lines, the role of various MAPK and AGC kinase isoforms was established *in vitro*, revealing that knock-down of ERK1/2, SAPK/JNK, and PKC $\beta$  resulted in cell death. Follow-up studies established that knockdown of PKC $\beta$  alone reduced cell proliferation, colony formation, and migratory capacity [204]. Genetic- and chemical inhibitor-based screens were combined to assess Notch pathway activation in breast cancer cells. In cells expressing a Hes1-firefly luciferase reporter, Notch activation was augmented in cells treated with an ERK inhibitor, while in a kinase-specific siRNA screen, knockdown of TRB3, a regulator of the MAPK-ERK pathway, significantly reduced cell proliferation and tumor growth [205]. Genetic and chemical inhibitor screens were also carried out in parallel to assess regulators of myeloid differentiation in cell lines of acute myeloid leukemia (AML). Two AML cell lines were treated with 84 kinases inhibitors, with pan GSK-3 inhibitors significantly affecting differentiation. Following this result, an independent genetic screen was carried out assessing the role of 1000 genes in myeloid differentiation with the  $\alpha$  homolog of GSK-3 scoring highly. Follow-up work knocked down GSK-3 $\alpha$  expression in AML cells, leading to decreased cell growth, increased apoptosis, and attenuated methylcellulose-induced colony formation [206]. Finally, Cronan *et al.* used siRNA to screen the metastatic human breast cancer cell line MDA-MD-231 to assess the role of seven MAP3Ks in tumor growth and metastasis. MDA-MB-231 cells with stable knockdown of one specific MAP3K were then injected into the fat pad of immune compromised mice. Results indicated roles for some MAP3Ks in both primary and metastatic tumor growth including MEKK2 and MLK3, while knockdown of MEKK1 had no effect on tumor growth but decreased metastatic tumor frequency. Tp12, MEKK3, and B-Raf inhibition was shown to decrease tumor growth with no effect on metastatic frequency, while TAK1 knockdown had no effect on primary or metastatic tumor growth [207].

Kinase screens can also be carried out to assess the effectiveness of drug combinations as well as synthetic lethal combinations. Liu-Sullivan *et al.* used human lung cancer cell lines with varying mutational profiles to screen for kinase knock-down that would influence the response to a polo-like kinase 1 (PLK1) inhibitor at both high and low doses. Results of this study revealed knockdown of 25 different kinases affected cellular response to PLK1 inhibition, with the majority increasing drug sensitivity including knockdown of the AGC kinases MAST4 and LATS2, while knockdown of another AGC kinase, SGK3, conferred resistance to the PLK1 inhibitor [208]. The development of drug resistance is a critical focus of research as it is an inevitable clinical occurrence for targeted therapies, and in many cases the mechanisms underlying resistance are unknown. The resistance

of ER $\alpha$  positive breast cancer to Tamoxifen has been well documented, thus an siRNA screen of known and putative kinases in MCF7 breast cancer cells was carried out to assess mediators of this acquired resistance. Results revealed a role for CDK10 in tumor cell sensitivity to Tamoxifen treatment, with follow-up studies attributing this resistance to CDK10 suppression-induced ETS2-driven transcription of C-Raf, leading to MAPK activation and loss of tumor growth reliance on estrogen signaling [209]. Another screen in human breast cancer cells sought to identify agents that modify resistance to PI3K inhibitor treatments. Over 40 agents affecting pathways involved in growth, metabolism, and apoptosis were applied to cell lines rendered insensitive to PI3K inhibitor treatment. The drug LEE011, a highly specific CDK4 and 6 inhibitor, emerged as the strongest sensitizer to PI3K inhibitor treatment, with sensitization also seen with two mTOR inhibitors and the Akt/PKB inhibitor MK2206. Following confirmation of results with siRNA-mediated knockdown, xenografts were treated with a PI3K inhibitor alone, or in combination with the CDK4 and CDK6 inhibitors. While PI3K inhibition alone slowed tumor growth, progression still occurred whereas combination treatment leads to a greater suppression of tumor growth [210].

An additional study examined the effect of suppressing GSK-3 activity on the functionality of oncology relevant drugs, with cell viability used as the readout. Results revealed that mTOR inhibitor function was perturbed to the greatest extent, with rapamycin and everolimus showing a complete loss of growth-inhibitory effects in combination with suppression of GSK-3 activity. In a second screen, a small molecule library of 367 kinase inhibitors was used to assess drug response rates and cell viability in combination with GSK-3 inhibition. The effectiveness of 89 separate compounds was altered upon GSK-3 inhibition, with mTOR inhibitor function again suppressed, while PLK1 inhibitor function was enhanced. Finally, the researchers performed a modifier RNAi screen of the entire human kinome in the presence/absence of GSK-3 inhibition. Their results revealed that approximately 35% of the kinome interacts with GSK-3 in the regulation of cell viability in human colonic epithelial cells, including previously described interactions with Akt, B-Raf, and CK1, as well as unidentified or underappreciated interactions with TNK2, LCK, ABL1, and PKC $\beta$  [211].

#### 1.3.4

##### **Functional Assessment of Kinase Function *in vivo*: Animal Models**

While knockdown of kinase activity through RNAi and chemical inhibitors allows for the assessment of kinase function, ultimately these models do not fully recapitulate the system as a whole with its many interacting cell types. The development of mouse models in which a particular kinase has been deleted provides a more informative method to ascertain their complete biological function. Taking advantage of endogenous homologous recombination, it is possible to engineer a mutation at a given locus in mouse embryonic stem cells that can then be used to generate a knockout mouse. While efforts are underway to generate knockout mice for the over 25 000 murine genes, the critical nature of some of these genes

results in embryonic lethality, hampering our ability to isolate the roles they play in adulthood.

The introduction of Cre-LoxP technology allowed for control over both the spatial and temporal activation/deletion of a given gene in the mouse. Cre recombinase is derived from the P1 bacteriophage and excises DNA placed between two LoxP sites, which are 34 base-pair nucleotide sequences [212–215]. Following inclusion of these sites into the genomic DNA and mouse generation, these animals are crossed to transgenic Cre expressing mice, where a tissue-specific promoter is used to drive Cre expression and limit gene excision to a particular cell type. This system can also be further modified to restrict when the Cre-mediated excision occurs, through the ligation of the ligand-binding domain of the estrogen receptor that restricts transcription until the mice are administered Tamoxifen [216, 217]. Two other methods to control temporal gene deletion include using the reverse-tetracycline-controlled transactivator, in which doxycycline must be administered in order to induce Cre expression [218] or Cre can be delivered to tissues through injection of a viral vector [219]. Both global- and tissue-specific knockouts have had immense impact on our understanding of kinase function in disease, below we highlight the significant gains in understanding of one particular kinase, GSK-3, through the use of animal models.

As described earlier, the significant sequence homology between the two homologs of GSK-3,  $\alpha$  and  $\beta$ , increased speculation that their functions would be highly similar with each being able to compensate for loss of the other. The use of knockout models revealed that the two homologs had critical nonredundant functions. Mice lacking GSK-3 $\alpha$  are viable with no overt abnormalities but, depending on background strain, have improved glucose tolerance, with increased ability to clear glucose as well as decreased total body fat mass and increased glycogen deposition in the liver. These effects on glucose clearance were not found on the C57BL/6 strain under normal conditions [44, 220] but were revealed under metabolic stress [220]. Other groups also have examined the consequence of global knockout of GSK-3 $\alpha$  expression in specific tissues, with global loss of this gene resulting in both decreased sperm motility and capacitation [221], increased cardiac hypertrophy and dysfunction following myocardial infarction [222, 223], and premature aging with the development of cardiac hypertrophy and dysfunction, impaired clearance of debris from skeletal muscle and loss of cartilage, and bone ossification from the knee joint [224]. In the brain, global loss of GSK-3 $\alpha$  altered cerebellar structure and function, with mice displaying decreased aggression, exploratory activity, loss of social motivation, and deficits in fear conditioning [225]. Unlike previous studies on the global GSK-3 $\alpha$  KO animals, a recent study described the compensatory upregulation in GSK-3 $\beta$  expression in the kidney medulla to which the lack of gross morphological changes was attributed. However, elevated water intake and urine output were seen in these mice, which was correlated with changes in aquaporin 2 expression [226].

The nonredundant functions in GSK-3 $\alpha$  and  $\beta$  were initially highlighted by the embryonic lethality associated with the GSK-3 $\beta$  knockout mouse [45]. This

lethality can occur at two different time points during gestation with mice dying mid-gestation due to significant liver degeneration, as a result of TNF $\alpha$ -mediated toxicity or death occurs later in gestation due to heart-patterning defects including the aorta and pulmonary artery arising from the right ventricle and ventricular septal defects [43]. GSK-3 $\beta$  knockout mice are embryonic lethal, while global GSK-3 $\beta$  heterozygotes are viable and allow study of the effects of partial knockdown of GSK-3 $\beta$  expression. These mice display impairments in long-term memory formation [227], as well as behavioral changes including decreased exploratory behavior without deficits in locomotion. These mice also have varying responses to injury in the heart with GSK-3 $\beta$  inhibition exacerbating ischemic injury but have a protective role in injury from ischemia/reperfusion [228].

Despite the information gathered from global GSK-3 $\alpha$  knockouts and GSK-3 $\beta$  heterozygotes, it was necessary to develop tissue-specific knockouts of both homologs to ascertain the effects of single cell type GSK-3 $\alpha$  loss, as well as to understand the role of complete GSK-3 $\beta$  loss in specific tissues. In the heart, the role of both  $\alpha$  and  $\beta$  has been investigated using Cre-targeted to cardiomyocytes. These studies revealed that GSK-3 $\alpha$  inactivation limits remodeling and preserves function following myocardial infarction, with similar responses seen in cardiomyocyte-specific GSK-3 $\beta$  KO. The effect of GSK-3 $\beta$  knockout was also assessed in response to pressure overload; however, these mice exhibited a normal hypertrophic response [229, 230]. Isoform-specific effects have also been closely examined in the brain, with deletion targeted to highly specific regions. Neuron-specific knockdown of GSK-3 $\alpha$  leads to alterations in basal synaptic transmission, as well as memory impairment, highlighting a role for the alpha homolog in synaptic plasticity [231]. Knockout of GSK-3 $\beta$  in D2-receptor-expressing neurons, but not D1-expressing neurons, was shown to mimic antipsychotic actions without affecting signaling pathways involved in mood-related behaviors [232]. Serotonin neuron-selective knockout of GSK-3 $\beta$  was also shown to decrease firing, cAMP production, and serotonin release affecting serotonin-regulated behaviors [233].

As GSK-3 has well-described roles in glucose handling, the knockout of either isoform has been studied in tissues linked to glucose metabolism. Liver-specific GSK-3 $\beta$  knockouts had normal glucose clearance and blood glucose levels, while skeletal muscle-specific knockout of GSK-3 $\beta$  resulted in improved glucose tolerance with no changes in glucose-stimulated insulin release [234]. Knockout of GSK-3 $\beta$  in skeletal muscle also enhanced recovery to muscle disuse with faster recovery of muscle mass following hind limb suspension and increased expression of myogenesis-associated genes [235]. Mice lacking GSK-3 $\beta$  in the  $\beta$ -islets of the pancreas showed mild improvement in glucose tolerance, as well as glucose-induced insulin secretion and increased  $\beta$ -cell mass. These mice were resistant to high-fat diet-induced diabetes as they showed improved glucose tolerance and expanded  $\beta$ -cell mass [236]. Breeding of  $\beta$ -cell-specific GSK-3 $\beta$  knockout mice with mice either lacking IRS1 expression (IRS1  $-/-$ ) or with reduced insulin receptor expression (IR  $+/-$  mice) prevented the development of insulin resistance and diabetes otherwise seen in these mice [237].

More recently, the consequence of tissue-specific loss of both GSK-3 $\alpha$  and  $\beta$  has been examined. Loss of both GSK-3 $\alpha$  and  $\beta$  was assessed in the brain through the use of nestin-Cre targeting neural progenitor cells. These mice displayed larger heads at every developmental stage until death at birth, with enlarged ventral brain areas and massive expansion of neuronal progenitor cell pools [238]. Targeted loss of both GSK-3 $\alpha$  and  $\beta$  in mammary gland epithelial cells leads to transdifferentiation of the mammary gland, resulting in dams unable to support nourishment of pups. These mice also developed large mammary ductal adenocarcinomas by 6–7 months of age, with tumors displaying areas of atypical squamous differentiation. The loss of both GSK-3 $\alpha$  and  $\beta$  leads to significant upregulation of  $\beta$ -catenin expression, and the development of these tumors closely resembled those seen in  $\beta$ -catenin mutant mice. To assess the importance of GSK-3 $\alpha$  and  $\beta$  outside of their effects on  $\beta$ -catenin, triple transgenic mice lacking mammary gland expression of GSK-3 $\alpha$  and  $\beta$  and  $\beta$ -catenin were established. While the lack of  $\beta$ -catenin corrected the gland transdifferentiation, these mice still developed tumors, though latency was increased by  $\sim$ 4 weeks and tumors displayed different histological characteristics [239].

These results highlight the complex, tissue- and isoform-specific functions of GSK-3 that could only be revealed in the context of a mouse model. Similar work has been undertaken to understand the role of other protein kinases in health and disease through the generation of global and conditional knockout animals. As a resource, several reviews are listed on the role of these other kinases [132, 240–242].

### 1.3.5

#### **CRISPR/Cas9 Genomic Recombineering**

Traditionally, as previously described, RNAi-mediated knockdown has been the technique most widely used to examine the importance of proteins to a given signaling pathway or disease model. However, given the reported issues with this technique including partial knockdown and off-target effects [241], improved approaches have been sought. Two promising techniques that allowed direct sequence-specific targeting of genes were Zinc Finger Nucleases (ZFNs) and Transcriptional Activator-Like Effector Nucleases (TALENs). Both of these techniques rely on DNA–protein interactions to guide the fused nuclease domain of the restriction enzyme FokI, to specific sites within the genome to activate nuclease activity. However, both of these techniques also have limitations/complications, including design issues limiting the regions of the genome that could be targeted, as well as protein engineering challenges [243, 244]. In essence, each target must be built from scratch by serial addition of nucleotide-specific protein domains. While these techniques were being refined, a further tool for genomic editing emerged from studies of prokaryotic immune systems.

The CRISPR (Clustered Regularly Interspaced Short Palindromic Repeats)/Cas system in prokaryotes functions as a defense mechanism against bacteriophage infection. In bacteria, CRISPR exploits sequence-specific homology pairing

between nucleic acids to direct nuclease activity to phage-induced DNA. Early studies revealed that bacteria possess three separate CRISPR/Cas systems, type I, II, and III, but type II was most readily engineered for genomic editing as it only requires a single protein, Cas9, to locate the target sequence and cleave DNA [245–247]. The type II CRISPR assay has three critical components required to engineer the system including Cas9, a tracrRNA, that facilitates DNA cleavage by Cas9, and a crRNA, which contains the target genomic sequence [244]. The system can then be targeted to a gene of interest through modification of the 20 nucleotide spacer sequence of the crRNA [247]. One critical aspect of this sequence is that it must contain the Protospacer-Adjacent Motif (PAM) immediately proximal to the region of homology to the DNA target, as the PAM sequence functions in the DNA target search mechanism of Cas9 and also elicits the transition between the two conformations of Cas9, from binding to cleavage [244, 247]. Further studies revealed that the tracrRNA and crRNA could be fused to create a single guide RNA (sgRNA) [248], further simplifying the system.

Two papers were published in 2013 that engineered the CRISPR/Cas system for use in mammalian cells [249, 250]. In mammals, Cas9-induced double-stranded breaks (DSBs) can be repaired by one of two pathways; Non-Homologous End Joining (NHEJ) or Homology-Directed Repair (HDR). NHEJ is more common as it does not require a homologous template; however, it is error prone, often resulting in insertions or deletions leading to frame-shift mutations and the absence of functional protein [244, 247]. This reliance on the error-prone NHEJ pathway allows this tool to be used in the generation of knockouts in a specific gene directed by the sgRNA, while introduction of an sgRNA and a single-stranded DNA template with homology to the target gene allows for introduction of transgenes, tags of interest or single base changes into the genome [247]. Since 2013, CRISPR/Cas technology has become widely used in animal models and human cancer cell lines for rapid generation of mutations and assessment of protein expression with greater specificity and fewer limitations to those carried out with RNAi. Some issues do remain such as off-target effects and the need sequence each change given the unpredictable nature of the repair, but the advantages and power of the approach outweigh these problems and have not yet hindered application.

Due to the recent adoption of this technique, its usefulness in the study of protein kinases in health and disease is in its infancy; however, genome-wide screens using sgRNA libraries have already been published. Three such studies have highlighted the role of proteins in proliferation and drug resistance, including a role for protein kinases [251–253]. In a study by Shalem *et al.*, two screens were performed to assess essential genes for survival of A375 human melanoma cell line and to identify genes that confer resistance to the B-Raf inhibitor vemurafenib. While the screen on cell survival showed that most of the deleted sgRNAs targeted essential genes such as ribosomal structural components, the resistance screen revealed a small group of previously reported and unreported genes including NF1, NF2, and the E3 ligase Cullin 3 [253]. A similar experiment was carried out to assess genes whose loss conferred resistance to etoposide, a widely

employed chemotherapeutic agent, in human leukemia cell lines. Results revealed a previously unreported role for the protein kinase CDK6 in etoposide-induced cytotoxicity [251].

Matano *et al.* employed CRISPR/Cas9 to examine the progression of colon cancer from benign adenoma to advanced adenocarcinoma by introducing mutations in normal human intestinal epithelial cells cultured into organoids with media supplemented with niche factors. Progressive genomic editing of organoids with sgRNAs targeting APC, SMAD4, and TP53 for knockout, as well as sgRNAs and templates for HDR-repair induced point mutations in KRas and PIK3CA, allowed for growth of organoids without niche factors in those containing alterations in all five genes. This approach also allowed the study of the effect of mutating each gene alone or in combination in progression from benign adenoma to adenocarcinoma, through injection of organoids into the kidney subcapsule. Off-target effects of CRISPR technology were also assessed through exome sequencing with no off-target mutations detected [254].

CRISPR/Cas9 technology has also been used to assess the role of mutations in PKC isoforms in cancer progression. This is a controversial area as controlled trials examining chemotherapy alone or in combination with PKC inhibitors, revealed significantly decreased response and disease control rates in nonsmall cell lung cancers treated with chemotherapeutic agents and PKC inhibitors [255]. These results may not be all that surprising given that 46 cancer-associated mutations have been detected in PKC, with the majority reducing or abolishing PKC activity and none leading to activation of the kinase [256]. HCT116 colon cancer cells contain a loss-of-function mutation in PKC $\beta$  and CRISPR/Cas9 was used to correct this mutation to the wild-type version of the sequence. Restoration of the wild-type PKC $\beta$  resulted in increased PKC activity as well as a more tumor-suppressive phenotype, in that HCT116 cells showed decreased anchorage-independent growth [256].

Aside from generating mutations in human cancer cell lines, the CRISPR approach can also be used *in vivo* to generate mutations in animal models. Delivery of CRISPR/Cas9 components targeting Pten to the mouse liver via hydrodynamic tail vein injections resulted in just over 3% of hepatocytes becoming null for Pten, while an additional 0.4% showed intermediate staining, which may represent heterozygous deletion. These results were duplicated in Pten floxed mice transduced with Cre via hydrodynamic injection. Coinjection of sgRNA targeting both Pten and p53 using the same method resulted in tumor development with similar histology to combined Pten and p53 floxed conditional knockout mice. This method also was used to establish point mutations in  $\beta$ -catenin through the injection of sgRNA targeting  $\beta$ -catenin, along with oligos containing four point mutations in  $\beta$ -catenin. Mutations in  $\beta$ -catenin, resulting in nuclear expression, were detected in approximately 0.5% of hepatocytes, demonstrating the reduced efficiency of HDR, but still resulting in increased target gene expression 7 days post injection [257]. Another method of adopting CRISPR/Cas9 technology became possible through generation of a Cre-dependent Rosa26 Cas9 knockin mouse, followed by viral or nonviral sgRNA delivery to induce genome

editing. This method has been used to assess mutations in the serine/threonine kinase LKB1, as well as KRas and p53, in lung adenocarcinoma, requiring introduction of a single vector containing the sgRNAs for required mutations, as well as Cre recombinase, delivered, via the trachea, into Cre-dependent Cas9 mice. Results showed lung nodule development with progression to advanced adenocarcinoma within 2 months and that LKB1/p53 double knockout cells were highly enriched over the entire lung [258].

CRISPR/Cas9 technology is rapidly changing the way the contributions of various proteins can be assessed in disease models as it is highly specific and allows for rapid generation of mouse models with multiple allelic changes, compared with techniques requiring introduction of targeted mutations into ES cells, via homologous recombination, to generate chimeric animals, followed by many months of breeding to generate experimental animals. Given the rapid uptake of this technology and advances in usability and application, CRISPR-mediated recombineering is likely to become the method of choice for both *in vitro* and *in vivo* analysis. Several excellent reviews have recently been published on the development of this technique, the mechanisms that underlie its specificity, as well as its use in various models of disease [244, 247, 259, 260].

#### 1.4

##### Final Thoughts

While there is little question that targeting members of the extensive kinome family has yielded a number of major and high profile successes, there is also a lesser appreciated but much greater volume of failures. Many highly selective and potent inhibitors have failed in preclinical and clinical trials for a variety of reasons. Should these targets be abandoned? The answer lies in understanding why these drugs failed and part of that derives from the ubiquity and dependencies of normal cellular regulation on this class of enzymes. Given the very finite number of potential targets, perhaps efforts should focus on precision targeting within tissues and even within cells. We know that signaling pathways are not freely diffusible and that a large fraction of signaling proteins are sequestered within macromolecular complexes, each tuned to specific functions, responses, and locations. This is how specificity and efficiency are maintained. Yet our clumsy techniques for modulating the functions of these proteins are indiscriminate in terms of subcellular location or context of complex. Whether it is a small molecule, siRNA or CRISPR recombination, all of the selected molecules in a given cell are inactivated. This may provide an opportunity for drug discovery whereby we target not the enzyme but its interaction with a specific set of complexes. This, coupled with more selective tissue targeting, may resuscitate targets that cause toxicity if universally inactivated. Clearly, an underappreciated subset of targeting proteins are the scaffolds that frame the macromolecular complexes, although, it should be pointed out that these scaffolds are often targeted by disease mutations, underscoring their importance.

Lastly, while the protein kinase family is large compared to most protein families, the research community has largely focused on a small subset. Of the 500+ members of this group, only 10–20% have been studied in any significant detail (<http://healthydebate.ca/opinions/canadian-biomedical-science-outstanding-but-also-redundant>). The emergence of genome-wide screens has started to change this picture, but the conservative, risk-averse nature of scientific funding acts as a disincentive for researchers to explore too far outside of the boundaries of current knowledge. For those who dare, there may be rich fruit to uncover.

### Acknowledgments

JLG is supported by a Canadian Breast Cancer Foundation fellowship (Ontario Division). JRW is supported by grants from the Terry Fox Foundation and Canadian Institutes of Health Research.

### References

- Hunter, T. (1987) A thousand and one protein kinases. *Cell*, **50** (6), 823–829.
- Manning, G., Whyte, D.B., Martinez, R., Hunter, T., and Sudarsanam, S. (2002) The protein kinase complement of the human genome. *Science*, **298** (5600), 1912–1934.
- Pearce, L.R., Komander, D., and Alessi, D.R. (2010) The nuts and bolts of AGC protein kinases. *Nat. Rev. Mol. Cell Biol.*, **11** (1), 9–22.
- Arencibia, J.M., Pastor-Flores, D., Bauer, A.F., Schulze, J.O., and Biondi, R.M. (2013) AGC protein kinases: from structural mechanism of regulation to allosteric drug development for the treatment of human diseases. *Biochim. Biophys. Acta*, **1834** (7), 1302–1321.
- Hanks, S.K. and Hunter, T. (1995) Protein kinases 6. The eukaryotic protein kinase superfamily: kinase (catalytic) domain structure and classification. *FASEB J.*, **9** (8), 576–596.
- Fyffe, C. and Falasca, M. (2013) 3-Phosphoinositide-dependent protein kinase-1 as an emerging target in the management of breast cancer. *Cancer Manage. Res.*, **5**, 271–280.
- Sato, S., Fujita, N., and Tsuruo, T. (2002) Interference with PDK1-akt survival signaling pathway by UCN-01 (7-hydroxystaurosporine). *Oncogene*, **21** (11), 1727–1738.
- Gonzalez, E. and McGraw, T.E. (2009) The akt kinases: isoform specificity in metabolism and cancer. *Cell Cycle*, **8** (16), 2502–2508.
- Chen, W.S., Xu, P.Z., Gottlob, K., Chen, M.L., Sokol, K., Shiyanova, T., Roninson, I., Weng, W., Suzuki, R., Tobe, K., Kadowaki, T., and Hay, N. (2001) Growth retardation and increased apoptosis in mice with homozygous disruption of the Akt1 gene. *Genes Dev.*, **15** (17), 2203–2208.
- Garofalo, R.S., Orena, S.J., Rafidi, K., Torchia, A.J., Stock, J.L., Hildebrandt, A.L., Coskran, T., Black, S.C., Brees, D.J., Wicks, J.R., McNeish, J.D., and Coleman, K.G. (2003) Severe diabetes, age-dependent loss of adipose tissue, and mild growth deficiency in mice lacking Akt2/PKB beta. *J. Clin. Invest.*, **112** (2), 197–208.
- Tschopp, O., Yang, Z.Z., Brodbeck, D., Dummler, B.A., Hemmings-Mieszczak, M., Watanabe, T., Michaelis, T., Frahm, J., and Hemmings, B.A. (2005) Essential role of protein kinase B gamma (PKB gamma/Akt3) in postnatal brain development but not in glucose

- homeostasis. *Development*, **132** (13), 2943–2954.
20. Tzivion, G., Dobson, M., and Ramakrishnan, G. (2011) FOXO transcription factors; regulation by AKT and 14-3-3 proteins. *Biochim. Biophys. Acta*, **1813** (11), 1938–1945.
  21. Xu, K., Liu, P., and Wei, W. (2014) mTOR signaling in tumorigenesis. *Biochim. Biophys. Acta*, **1846** (2), 638–654.
  22. Mochly-Rosen, D., Das, K., and Grimes, K.V. (2012) Protein kinase C, an elusive therapeutic target? *Nat. Rev. Drug Discovery*, **11** (12), 937–957.
  23. Urtreger, A.J., Kazanietz, M.G., and Bal de Kier Joffe, E.D. (2012) Contribution of individual PKC isoforms to breast cancer progression. *IUBMB Life*, **64** (1), 18–26.
  24. Lang, F., Perrotti, N., and Stournaras, C. (2010) Colorectal carcinoma cells--regulation of survival and growth by SGK1. *Int. J. Biochem. Cell Biol.*, **42** (10), 1571–1575.
  25. Tessier, M. and Woodgett, J.R. (2006) Role of the phox homology domain and phosphorylation in activation of serum and glucocorticoid-regulated kinase-3. *J. Biol. Chem.*, **281** (33), 23978–23989.
  26. Wayman, G.A., Lee, Y.S., Tokumitsu, H., Silva, A.J., and Soderling, T.R. (2008) Calmodulin-kinases: modulators of neuronal development and plasticity. *Neuron*, **59** (6), 914–931.
  27. Maier, L.S. and Bers, D.M. (2007) Role of Ca<sup>2+</sup>/calmodulin-dependent protein kinase (CaMK) in excitation-contraction coupling in the heart. *Cardiovasc. Res.*, **73** (4), 631–640.
  28. Stull, J.T., Kamm, K.E., and Vandenboom, R. (2011) Myosin light chain kinase and the role of myosin light chain phosphorylation in skeletal muscle. *Arch. Biochem. Biophys.*, **510** (2), 120–128.
  29. Gowans, G.J., Hawley, S.A., Ross, F.A., and Hardie, D.G. (2013) AMP is a true physiological regulator of AMP-activated protein kinase by both allosteric activation and enhancing net phosphorylation. *Cell Metab.*, **18** (4), 556–566.
  30. Grahame Hardie, D. (2014) AMP-activated protein kinase: a key regulator of energy balance with many roles in human disease. *J. Int. Med.*, **276** (6), 543–559.
  31. Varjosalo, M., Keskitalo, S., Van Droogen, A., Nurkkala, H., Vichalkovski, A., Aebersold, R., and Gstaiger, M. (2013) The protein interaction landscape of the human CMGC kinase group. *Cell Rep.*, **3** (4), 1306–1320.
  32. Malumbres, M., Harlow, E., Hunt, T., Hunter, T., Lahti, J.M., Manning, G., Morgan, D.O., Tsai, L.H., and Wolgemuth, D.J. (2009) Cyclin-dependent kinases: a family portrait. *Nat. Cell Biol.*, **11** (11), 1275–1276.
  33. Day, P.J., Cleasby, A., Tickle, I.J., O'Reilly, M., Coyle, J.E., Holding, F.P., McMenamin, R.L., Yon, J., Chopra, R., Lengauer, C., and Jhoti, H. (2009) Crystal structure of human CDK4 in complex with a D-type cyclin. *Proc. Natl. Acad. Sci. U.S.A.*, **106** (11), 4166–4170.
  34. Takaki, T., Echaliier, A., Brown, N.R., Hunt, T., Endicott, J.A., and Noble, M.E. (2009) The structure of CDK4/cyclin D3 has implications for models of CDK activation. *Proc. Natl. Acad. Sci. U.S.A.*, **106** (11), 4171–4176.
  35. Malumbres, M. (2014) Cyclin-dependent kinases. *Genome Biol.*, **15** (6), 122.
  36. Kawauchi, T. (2014) Cdk5 regulates multiple cellular events in neural development, function and disease. *Dev. Growth Differ.*, **56** (5), 335–348.
  37. Tsai, L.H., Takahashi, T., Caviness, V.S. Jr., and Harlow, E. (1993) Activity and expression pattern of cyclin-dependent kinase 5 in the embryonic mouse nervous system. *Development*, **119** (4), 1029–1040.
  38. Kim, E.K. and Choi, E.J. (2015) Compromised MAPK signaling in human diseases: an update. *Arch. Toxicol.*, **89** (6), 867–882.
  39. Burotto, M., Chiou, V.L., Lee, J.M., and Kohn, E.C. (2014) The MAPK pathway across different malignancies: a new perspective. *Cancer*, **120** (22), 3446–3456.

32. Roberts, P.J. and Der, C.J. (2007) Targeting the raf-MEK-ERK mitogen-activated protein kinase cascade for the treatment of cancer. *Oncogene*, **26** (22), 3291–3310.
33. Crews, C.M., Alessandrini, A., and Erikson, R.L. (1992) The primary structure of MEK, a protein kinase that phosphorylates the ERK gene product. *Science*, **258** (5081), 478–480.
34. Roskoski, R. Jr., (2010) RAF protein-serine/threonine kinases: structure and regulation. *Biochem. Biophys. Res. Commun.*, **399** (3), 313–317.
35. Lee, J.D., Ulevitch, R.J., and Han, J. (1995) Primary structure of BMK1: a new mammalian map kinase. *Biochem. Biophys. Res. Commun.*, **213** (2), 715–724.
36. Kato, Y., Chao, T.H., Hayashi, M., Tapping, R.I., and Lee, J.D. (2000) Role of BMK1 in regulation of growth factor-induced cellular responses. *Immunol. Res.*, **21** (2–3), 233–237.
37. Keshet, Y. and Seger, R. (2010) The MAP kinase signaling cascades: a system of hundreds of components regulates a diverse array of physiological functions. *Methods Mol. Biol.*, **661**, 3–38.
38. Sabio, G. and Davis, R.J. (2014) TNF and MAP kinase signalling pathways. *Semin. Immunol.*, **26** (3), 237–245.
39. Davis, R.J. (2000) Signal transduction by the JNK group of MAP kinases. *Cell*, **103** (2), 239–252.
40. Woodgett, J.R. (1990) Molecular cloning and expression of glycogen synthase kinase-3/factor A. *EMBO J.*, **9** (8), 2431–2438.
41. Ling, L.S., Voskas, D., and Woodgett, J.R. (2013) Activation of PDK-1 maintains mouse embryonic stem cell self-renewal in a PKB-dependent manner. *Oncogene*, **32** (47), 5397–5408.
42. Kockeritz, L., Doble, B., Patel, S., and Woodgett, J.R. (2006) Glycogen synthase kinase-3—an overview of an over-achieving protein kinase. *Curr. Drug Targets*, **7** (11), 1377–1388.
43. Kerkela, R., Kockeritz, L., Macaulay, K., Zhou, J., Doble, B.W., Beahm, C., Greytak, S., Woulfe, K., Trivedi, C.M., Woodgett, J.R., Epstein, J.A., Force, T., and Huggins, G.S. (2008) Deletion of GSK-3beta in mice leads to hypertrophic cardiomyopathy secondary to cardiomyoblast hyperproliferation. *J. Clin. Invest.*, **118** (11), 3609–3618.
44. MacAulay, K., Doble, B.W., Patel, S., Hansotia, T., Sinclair, E.M., Drucker, D.J., Nagy, A., and Woodgett, J.R. (2007) Glycogen synthase kinase 3alpha-specific regulation of murine hepatic glycogen metabolism. *Cell Metab.*, **6** (4), 329–337.
45. Hoeflich, K.P., Luo, J., Rubie, E.A., Tsao, M.S., Jin, O., and Woodgett, J.R. (2000) Requirement for glycogen synthase kinase-3beta in cell survival and NF-kappaB activation. *Nature*, **406** (6791), 86–90.
46. Xu, B.E., Lee, B.H., Min, X., Lenertz, L., Heise, C.J., Stippec, S., Goldsmith, E.J., and Cobb, M.H. (2005) WNK1: analysis of protein kinase structure, downstream targets, and potential roles in hypertension. *Cell Res.*, **15** (1), 6–10.
47. McCormick, J.A. and Ellison, D.H. (2011) The WNKs: atypical protein kinases with pleiotropic actions. *Physiol. Rev.*, **91** (1), 177–219.
48. Ubersax, J.A. and Ferrell, J.E. Jr., (2007) Mechanisms of specificity in protein phosphorylation. *Nat. Rev. Mol. Cell Biol.*, **8** (7), 530–541.
49. Lemmon, M.A. and Schlessinger, J. (2010) Cell signaling by receptor tyrosine kinases. *Cell*, **141** (7), 1117–1134.
50. Takeuchi, K. and Ito, F. (2011) Receptor tyrosine kinases and targeted cancer therapeutics. *Biol. Pharm. Bull.*, **34** (12), 1774–1780.
51. Schlessinger, J. and Lemmon, M.A. (2003) SH2 and PTB domains in tyrosine kinase signaling. *Sci. STKE*, **2003** (191), RE12.
52. Hunter, T. (2009) Tyrosine phosphorylation: thirty years and counting. *Curr. Opin. Cell Biol.*, **21** (2), 140–146.
53. Colicelli, J. (2010) ABL tyrosine kinases: evolution of function, regulation, and specificity. *Sci. Signal.*, **3** (139), re6.
54. Kim, L.C., Song, L., and Haura, E.B. (2009) Src kinases as therapeutic targets for cancer. *Nat. Rev. Clin. Oncol.*, **6** (10), 587–595.

55. Balabanov, S., Braig, M., and Brummendorf, T.H. (2014) Current aspects in resistance against tyrosine kinase inhibitors in chronic myelogenous leukemia. *Drug Discovery Today Technol.*, **11**, 89–99.
56. Meijer, L., Flajolet, M., and Greengard, P. (2004) Pharmacological inhibitors of glycogen synthase kinase 3. *Trends Pharmacol. Sci.*, **25** (9), 471–480.
57. Liu, F., Ma, X.H., Ule, J., Bibb, J.A., Nishi, A., DeMaggio, A.J., Yan, Z., Nairn, A.C., and Greengard, P. (2001) Regulation of cyclin-dependent kinase 5 and casein kinase 1 by metabotropic glutamate receptors. *Proc. Natl. Acad. Sci. U.S.A.*, **98** (20), 11062–11068.
58. Walter, J., Fluhrer, R., Hartung, B., Willem, M., Kaether, C., Capell, A., Lammich, S., Multhaup, G., and Haass, C. (2001) Phosphorylation regulates intracellular trafficking of beta-secretase. *J. Biol. Chem.*, **276** (18), 14634–14641.
59. Xiao, J., Tagliabracci, V.S., Wen, J., Kim, S.A., and Dixon, J.E. (2013) Crystal structure of the golgi casein kinase. *Proc. Natl. Acad. Sci. U.S.A.*, **110** (26), 10574–10579.
60. Gould, K.L. and Nurse, P. (1989) Tyrosine phosphorylation of the fission yeast *cdc2+* protein kinase regulates entry into mitosis. *Nature*, **342**, 39–45.
61. Potter, L.R. (2005) Domain analysis of human transmembrane guanylyl cyclase receptors: implications for regulation. *Front. Biosci.*, **10**, 1205–1220.
62. Saha, S., Biswas, K.H., Kondapalli, C., Isloor, N., and Visweswariah, S.S. (2009) The linker region in receptor guanylyl cyclases is a key regulatory module: mutational analysis of guanylyl cyclase C. *J. Biol. Chem.*, **284** (40), 27135–27145.
63. Oliver, P.M., Fox, J.E., Kim, R., Rockman, H.A., Kim, H.S., Reddick, R.L., Pandey, K.N., Milgram, S.L., Smithies, O., and Maeda, N. (1997) Hypertension, cardiac hypertrophy, and sudden death in mice lacking natriuretic peptide receptor A. *Proc. Natl. Acad. Sci. U.S.A.*, **94** (26), 14730–14735.
64. Lopez, M.J., Wong, S.K., Kishimoto, I., Dubois, S., Mach, V., Friesen, J., Garbers, D.L., and Beuve, A. (1995) Salt-resistant hypertension in mice lacking the guanylyl cyclase-A receptor for atrial natriuretic peptide. *Nature*, **378** (6552), 65–68.
65. Yang, R.B., Robinson, S.W., Xiong, W.H., Yau, K.W., Birch, D.G., and Garbers, D.L. (1999) Disruption of a retinal guanylyl cyclase gene leads to cone-specific dystrophy and paradoxical rod behavior. *J. Neurosci.*, **19** (14), 5889–5897.
66. Dowling, R.J., Topisirovic, I., Fonseca, B.D., and Sonenberg, N. (2010) Dissecting the role of mTOR: lessons from mTOR inhibitors. *Biochim. Biophys. Acta*, **1804** (3), 433–439.
67. Knuutila, S., Bjorkqvist, A.M., Autio, K., Tarkkanen, M., Wolf, M., Monni, O., Szymanska, J., Larramendy, M.L., Tapper, J., Pere, H., El-Rifai, W., Hemmer, S., Wasenius, V.M., Vidgren, V., and Zhu, Y. (1998) DNA copy number amplifications in human neoplasms: review of comparative genomic hybridization studies. *Am. J. Pathol.*, **152** (5), 1107–1123.
68. Tell, V. and Hilgeroth, A. (2013) Recent developments of protein kinase inhibitors as potential AD therapeutics. *Front. Cell. Neurosci.*, **7**, 189.
69. Giacobini, E. and Becker, R.E. (2007) One hundred years after the discovery of Alzheimer's disease. A turning point for therapy? *J. Alzheimers Dis.*, **12** (1), 37–52.
70. Benilova, I., Karran, E., and De Strooper, B. (2012) The toxic abeta oligomer and Alzheimer's disease: an emperor in need of clothes. *Nat. Neurosci.*, **15** (3), 349–357.
71. Kuperstein, I., Broersen, K., Benilova, I., Rozenski, J., Jonckheere, W., Debulpaep, M., Vandersteen, A., Segers-Nolten, I., Van Der Werf, K., Subramaniam, V., Braeken, D., Callewaert, G., Bartic, C., D'Hooge, R., Martins, I.C., Rousseau, F., Schymkowitz, J., and De Strooper, B. (2010) Neurotoxicity of Alzheimer's disease abeta peptides is induced by small

- changes in the Abeta42 to Abeta40 ratio. *EMBO J.*, **29** (19), 3408–3420.
72. Avila, J., Lucas, J.J., Perez, M., and Hernandez, F. (2004) Role of tau protein in both physiological and pathological conditions. *Physiol. Rev.*, **84** (2), 361–384.
  73. Avila, J. (2006) Tau phosphorylation and aggregation in Alzheimer's disease pathology. *FEBS Lett.*, **580** (12), 2922–2927.
  74. Castro-Alvarez, J.F., Uribe-Arias, S.A., Mejia-Raigosa, D., and Cardona-Gomez, G.P. (2014) Cyclin-dependent kinase 5, a node protein in diminished tauopathy: a systems biology approach. *Front. Aging Neurosci.*, **6**, 232.
  75. Hye, A., Kerr, F., Archer, N., Foy, C., Poppe, M., Brown, R., Hamilton, G., Powell, J., Anderton, B., and Lovestone, S. (2005) Glycogen synthase kinase-3 is increased in white cells early in Alzheimer's disease. *Neurosci. Lett.*, **373** (1), 1–4.
  76. Kwok, J.B., Loy, C.T., Hamilton, G., Lau, E., Hallupp, M., Williams, J., Owen, M.J., Broe, G.A., Tang, N., Lam, L., Powell, J.F., Lovestone, S., and Schofield, P.R. (2008) Glycogen synthase kinase-3beta and tau genes interact in Alzheimer's disease. *Ann. Neurol.*, **64** (4), 446–454.
  77. Leroy, K., Yilmaz, Z., and Brion, J.P. (2007) Increased level of active GSK-3beta in Alzheimer's disease and accumulation in argyrophilic grains and in neurones at different stages of neurofibrillary degeneration. *Neuropathol. Appl. Neurobiol.*, **33** (1), 43–55.
  78. Hooper, C., Killick, R., and Lovestone, S. (2008) The GSK3 hypothesis of Alzheimer's disease. *J. Neurochem.*, **104** (6), 1433–1439.
  79. Huang, H.C. and Klein, P.S. (2006) Multiple roles for glycogen synthase kinase-3 as a drug target in Alzheimer's disease. *Curr. Drug Targets*, **7** (11), 1389–1397.
  80. Chico, L.K., Van Eldik, L.J., and Watterson, D.M. (2009) Targeting protein kinases in central nervous system disorders. *Nat. Rev. Drug Discovery*, **8** (11), 892–909.
  81. Takashima, A., Noguchi, K., Michel, G., Mercken, M., Hoshi, M., Ishiguro, K., and Imahori, K. (1996) Exposure of rat hippocampal neurons to amyloid beta peptide (25-35) induces the inactivation of phosphatidylinositol-3 kinase and the activation of tau protein kinase I/glycogen synthase kinase-3 beta. *Neurosci. Lett.*, **203** (1), 33–36.
  82. Ly, P.T., Wu, Y., Zou, H., Wang, R., Zhou, W., Kinoshita, A., Zhang, M., Yang, Y., Cai, F., Woodgett, J., and Song, W. (2013) Inhibition of GSK3beta-mediated BACE1 expression reduces Alzheimer-associated phenotypes. *J. Clin. Invest.*, **123** (1), 224–235.
  83. Martin, L., Latypova, X., Wilson, C.M., Magnaudeix, A., Perrin, M.L., Yardin, C., and Terro, F. (2013) Tau protein kinases: involvement in Alzheimer's disease. *Ageing Res. Rev.*, **12** (1), 289–309.
  84. Hernandez, F., Borrell, J., Guaza, C., Avila, J., and Lucas, J.J. (2002) Spatial learning deficit in transgenic mice that conditionally over-express GSK-3beta in the brain but do not form tau filaments. *J. Neurochem.*, **83** (6), 1529–1533.
  85. Gomez de Barreda, E., Perez, M., Gomez Ramos, P., de Cristobal, J., Martin-Maestro, P., Moran, A., Dawson, H.N., Vitek, M.P., Lucas, J.J., Hernandez, F., and Avila, J. (2010) Tau-knockout mice show reduced GSK3-induced hippocampal degeneration and learning deficits. *Neurobiol. Dis.*, **37** (3), 622–629.
  86. Engel, T., Hernandez, F., Avila, J., and Lucas, J.J. (2006) Full reversal of Alzheimer's disease-like phenotype in a mouse model with conditional overexpression of glycogen synthase kinase-3. *J. Neurosci.*, **26** (19), 5083–5090.
  87. Rockenstein, E., Torrance, M., Adame, A., Mante, M., Bar-on, P., Rose, J.B., Crews, L., and Masliah, E. (2007) Neuroprotective effects of regulators of the glycogen synthase kinase-3beta signaling pathway in a transgenic model of Alzheimer's disease are associated with reduced amyloid

- precursor protein phosphorylation. *J. Neurosci.*, **27** (8), 1981–1991.
88. Stambolic, V., Ruel, L., and Woodgett, J.R. (1996) Lithium inhibits glycogen synthase kinase-3 activity and mimics wingless signalling in intact cells. *Curr. Biol.*, **6** (12), 1664–1668.
  89. Oruch, R., Elderbi, M.A., Khattab, H.A., Pryme, I.F., and Lund, A. (2014) Lithium: a review of pharmacology, clinical uses, and toxicity. *Eur. J. Pharmacol.*, **740**, 464–473.
  90. Liu, S.J., Zhang, A.H., Li, H.L., Wang, Q., Deng, H.M., Netzer, W.J., Xu, H., and Wang, J.Z. (2003) Overactivation of glycogen synthase kinase-3 by inhibition of phosphoinositol-3 kinase and protein kinase C leads to hyperphosphorylation of tau and impairment of spatial memory. *J. Neurochem.*, **87** (6), 1333–1344.
  91. De Ferrari, G.V., Chacon, M.A., Barria, M.I., Garrido, J.L., Godoy, J.A., Olivares, G., Reyes, A.E., Alvarez, A., Bronfman, M., and Inestrosa, N.C. (2003) Activation of wnt signaling rescues neurodegeneration and behavioral impairments induced by beta-amyloid fibrils. *Mol. Psychiatry*, **8** (2), 195–208.
  92. King, M.K., Pardo, M., Cheng, Y., Downey, K., Jope, R.S., and Beurel, E. (2014) Glycogen synthase kinase-3 inhibitors: rescuers of cognitive impairments. *Pharmacol. Ther.*, **141** (1), 1–12.
  93. Fiorentini, A., Rosi, M.C., Grossi, C., Luccarini, I., and Casamenti, F. (2010) Lithium improves hippocampal neurogenesis, neuropathology and cognitive functions in APP mutant mice. *PLoS One*, **5** (12), e14382.
  94. Hampel, H., Ewers, M., Burger, K., Annas, P., Mortberg, A., Bogstedt, A., Frolich, L., Schroder, J., Schonknecht, P., Riepe, M.W., Kraft, I., Gasser, T., Leyhe, T., Moller, H.J., Kurz, A., and Basun, H. (2009) Lithium trial in Alzheimer's disease: a randomized, single-blind, placebo-controlled, multicenter 10-week study. *J. Clin. Psychiatry*, **70** (6), 922–931.
  95. Sereno, L., Coma, M., Rodriguez, M., Sanchez-Ferrer, P., Sanchez, M.B., Gich, I., Agullo, J.M., Perez, M., Avila, J., Guardia-Laguarta, C., Clarimon, J., Lleó, A., and Gomez-Isla, T. (2009) A novel GSK-3beta inhibitor reduces Alzheimer's pathology and rescues neuronal loss in vivo. *Neurobiol. Dis.*, **35** (3), 359–367.
  96. Hu, S., Begum, A.N., Jones, M.R., Oh, M.S., Beech, W.K., Beech, B.H., Yang, F., Chen, P., Ubeda, O.J., Kim, P.C., Davies, P., Ma, Q., Cole, G.M., and Frautschy, S.A. (2009) GSK3 inhibitors show benefits in an Alzheimer's disease (AD) model of neurodegeneration but adverse effects in control animals. *Neurobiol. Dis.*, **33** (2), 193–206.
  97. Shah, K. and Lahiri, D.K. (2014) Cdk5 activity in the brain – multiple paths of regulation. *J. Cell Sci.*, **127**(Pt. 11), 2391–2400.
  98. Patrick, G.N., Zukerberg, L., Nikolic, M., de la Monte, S., Dikkes, P., and Tsai, L.H. (1999) Conversion of p35 to p25 deregulates Cdk5 activity and promotes neurodegeneration. *Nature*, **402** (6762), 615–622.
  99. Sengupta, A., Kabat, J., Novak, M., Wu, Q., Grundke-Iqbal, I., and Iqbal, K. (1998) Phosphorylation of tau at both thr 231 and ser 262 is required for maximal inhibition of its binding to microtubules. *Arch. Biochem. Biophys.*, **357** (2), 299–309.
  100. Cruz, J.C., Tseng, H.C., Goldman, J.A., Shih, H., and Tsai, L.H. (2003) Aberrant Cdk5 activation by p25 triggers pathological events leading to neurodegeneration and neurofibrillary tangles. *Neuron*, **40** (3), 471–483.
  101. Piedrahita, D., Hernandez, I., Lopez-Tobon, A., Fedorov, D., Obara, B., Manjunath, B.S., Boudreau, R.L., Davidson, B., Laferla, F., Gallego-Gomez, J.C., Kosik, K.S., and Cardona-Gomez, G.P. (2010) Silencing of CDK5 reduces neurofibrillary tangles in transgenic Alzheimer's mice. *J. Neurosci.*, **30** (42), 13966–13976.
  102. Wen, Y., Planel, E., Herman, M., Figueroa, H.Y., Wang, L., Liu, L., Lau, L.F., Yu, W.H., and Duff, K.E. (2008) Interplay between cyclin-dependent kinase 5 and glycogen

- synthase kinase 3 beta mediated by neuregulin signaling leads to differential effects on tau phosphorylation and amyloid precursor protein processing. *J. Neurosci.*, **28** (10), 2624–2632.
103. Perry, G., Roder, H., Nunomura, A., Takeda, A., Friedlich, A.L., Zhu, X., Raina, A.K., Holbrook, N., Siedlak, S.L., Harris, P.L., and Smith, M.A. (1999) Activation of neuronal extracellular receptor kinase (ERK) in Alzheimer disease links oxidative stress to abnormal phosphorylation. *Neuroreport*, **10** (11), 2411–2415.
  104. Ploia, C., Antoniou, X., Sclip, A., Grande, V., Cardinetti, D., Colombo, A., Canu, N., Benussi, L., Ghidoni, R., Forloni, G., and Borsello, T. (2011) JNK plays a key role in tau hyperphosphorylation in Alzheimer's disease models. *J. Alzheimers Dis.*, **26** (2), 315–329.
  105. Chong, Y.H., Shin, Y.J., Lee, E.O., Kaye, R., Glabe, C.G., and Tenner, A.J. (2006) ERK1/2 activation mediates abeta oligomer-induced neurotoxicity via caspase-3 activation and tau cleavage in rat organotypic hippocampal slice cultures. *J. Biol. Chem.*, **281** (29), 20315–20325.
  106. Morishima, Y., Gotoh, Y., Zieg, J., Barrett, T., Takano, H., Flavell, R., Davis, R.J., Shirasaki, Y., and Greenberg, M.E. (2001) Beta-amyloid induces neuronal apoptosis via a mechanism that involves the c-jun N-terminal kinase pathway and the induction of fas ligand. *J. Neurosci.*, **21** (19), 7551–7560.
  107. Feld, M., Krawczyk, M.C., Sol Fustinana, M., Blake, M.G., Baratti, C.M., Romano, A., and Boccia, M.M. (2014) Decrease of ERK/MAPK overactivation in prefrontal cortex reverses early memory deficit in a mouse model of Alzheimer's disease. *J. Alzheimers Dis.*, **40** (1), 69–82.
  108. Quiroz-Baez, R., Rojas, E., and Arias, C. (2009) Oxidative stress promotes JNK-dependent amyloidogenic processing of normally expressed human APP by differential modification of alpha-, beta- and gamma-secretase expression. *Neurochem. Int.*, **55** (7), 662–670.
  109. Guglielmotto, M., Monteleone, D., Giliberto, L., Fornaro, M., Borghi, R., Tamagno, E., and Tabaton, M. (2011) Amyloid-beta(4)(2) activates the expression of BACE1 through the JNK pathway. *J. Alzheimers Dis.*, **27** (4), 871–883.
  110. Mazzitelli, S., Xu, P., Ferrer, I., Davis, R.J., and Tournier, C. (2011) The loss of c-jun N-terminal protein kinase activity prevents the amyloidogenic cleavage of amyloid precursor protein and the formation of amyloid plaques in vivo. *J. Neurosci.*, **31** (47), 16969–16976.
  111. Sclip, A., Tozzi, A., Abaza, A., Cardinetti, D., Colombo, I., Calabresi, P., Salmona, M., Welker, E., and Borsello, T. (2014) c-jun N-terminal kinase has a key role in Alzheimer disease synaptic dysfunction in vivo. *Cell Death Dis.*, **5**, e1019.
  112. Hensley, K., Floyd, R.A., Zheng, N.Y., Nael, R., Robinson, K.A., Nguyen, X., Pye, Q.N., Stewart, C.A., Geddes, J., Markesbery, W.R., Patel, E., Johnson, G.V., and Bing, G. (1999) p38 kinase is activated in the Alzheimer's disease brain. *J. Neurochem.*, **72** (5), 2053–2058.
  113. Sheng, J.G., Jones, R.A., Zhou, X.Q., McGinness, J.M., Van Eldik, L.J., Mrak, R.E., and Griffin, W.S. (2001) Interleukin-1 promotion of MAPK-p38 overexpression in experimental animals and in Alzheimer's disease: potential significance for tau protein phosphorylation. *Neurochem. Int.*, **39** (5–6), 341–348.
  114. Sun, A., Liu, M., Nguyen, X.V., and Bing, G. (2003) P38 MAP kinase is activated at early stages in Alzheimer's disease brain. *Exp. Neurol.*, **183** (2), 394–405.
  115. Swatton, J.E., Sellers, L.A., Faull, R.L., Holland, A., Iritani, S., and Bahn, S. (2004) Increased MAP kinase activity in Alzheimer's and down syndrome but not in schizophrenia human brain. *Eur. J. Neurosci.*, **19** (10), 2711–2719.
  116. Giovannini, M.G., Scali, C., Prosperi, C., Bellucci, A., Vannucchi, M.G., Rosi, S., Pepeu, G.,

- and Casamenti, F. (2002) Beta-amyloid-induced inflammation and cholinergic hypofunction in the rat brain in vivo: involvement of the p38MAPK pathway. *Neurobiol. Dis.*, **11** (2), 257–274.
117. Giraldo, E., Lloret, A., Fuchsberger, T., and Vina, J. (2014) Abeta and tau toxicities in Alzheimer's are linked via oxidative stress-induced p38 activation: protective role of vitamin E. *Redox Biol.*, **2**, 873–877.
118. Bach, J.P., Mengel, D., Wahle, T., Kautz, A., Balzer-Geldsetzer, M., Al-Abed, Y., Dodel, R., and Bacher, M. (2011) The role of CN1-1493 in the function of primary microglia with respect to amyloid-beta. *J. Alzheimers Dis.*, **26** (1), 69–80.
119. Roy, S.M., Grum-Tokars, V.L., Schavocky, J.P., Saeed, F., Staniszewski, A., Teich, A.F., Arancio, O., Bachstetter, A.D., Webster, S.J., Van Eldik, L.J., Minasov, G., Anderson, W.F., Pelletier, J.C., and Watterson, D.M. (2015) Targeting human central nervous system protein kinases: an isoform selective p38alpha MAPK inhibitor that attenuates disease progression in Alzheimer's disease mouse models. *ACS Chem. Neurosci.*, **6** (4), 666–680.
120. Pei, J.J., An, W.L., Zhou, X.W., Nishimura, T., Norberg, J., Benedikz, E., Gotz, J., and Winblad, B. (2006) P70 S6 kinase mediates tau phosphorylation and synthesis. *FEBS Lett.*, **580** (1), 107–114.
121. Ma, T., Hoeffler, C.A., Capetillo-Zarate, E., Yu, F., Wong, H., Lin, M.T., Tampellini, D., Klann, E., Blitzer, R.D., and Gouras, G.K. (2010) Dysregulation of the mTOR pathway mediates impairment of synaptic plasticity in a mouse model of Alzheimer's disease. *PLoS One*, **5** (9). doi: 10.1371/journal.pone.0012845
122. Avrahami, L., Farfara, D., Shaham-Kol, M., Vassar, R., Frenkel, D., and Eldar-Finkelman, H. (2013) Inhibition of glycogen synthase kinase-3 ameliorates beta-amyloid pathology and restores lysosomal acidification and mammalian target of rapamycin activity in the Alzheimer disease mouse model: in vivo and in vitro studies. *J. Biol. Chem.*, **288** (2), 1295–1306.
123. Lafay-Chebassier, C., Perault-Pochat, M.C., Page, G., Rioux Bilan, A., Damjanac, M., Pain, S., Houeto, J.L., Gil, R., and Hugon, J. (2006) The immunosuppressant rapamycin exacerbates neurotoxicity of abeta peptide. *J. Neurosci. Res.*, **84** (6), 1323–1334.
124. Sun, M.K. and Alkon, D.L. (2014) The "memory kinases": roles of PKC isoforms in signal processing and memory formation. *Prog. Mol. Biol. Transl. Sci.*, **122**, 31–59.
125. Skovronsky, D.M., Moore, D.B., Milla, M.E., Doms, R.W., and Lee, V.M. (2000) Protein kinase C-dependent alpha-secretase competes with beta-secretase for cleavage of amyloid-beta precursor protein in the trans-golgi network. *J. Biol. Chem.*, **275** (4), 2568–2575.
126. Etcheberrigaray, R., Tan, M., Dewachter, I., Kuiperi, C., Van der Auwera, I., Wera, S., Qiao, L., Bank, B., Nelson, T.J., Kozikowski, A.P., Van Leuven, F., and Alkon, D.L. (2004) Therapeutic effects of PKC activators in Alzheimer's disease transgenic mice. *Proc. Natl. Acad. Sci. U.S.A.*, **101** (30), 11141–11146.
127. Lee, W., Boo, J.H., Jung, M.W., Park, S.D., Kim, Y.H., Kim, S.U., and Mook-Jung, I. (2004) Amyloid beta peptide directly inhibits PKC activation. *Mol. Cell. Neurosci.*, **26** (2), 222–231.
128. Pakaski, M., Balaspiri, L., Checler, F., and Kasa, P. (2002) Human amyloid-beta causes changes in the levels of endothelial protein kinase C and its alpha isoform in vitro. *Neurochem. Int.*, **41** (6), 409–414.
129. Kim, T., Hinton, D.J., and Choi, D.S. (2011) Protein kinase C-regulated abeta production and clearance. *Int. J. Alzheimers Dis.*, **2011**, 857368.
130. Choi, D.S., Wang, D., Yu, G.Q., Zhu, G., Kharazia, V.N., Paredes, J.P., Chang, W.S., Deitchman, J.K., Mucke, L., and Messing, R.O. (2006) PKCepsilon increases endothelin converting enzyme activity and reduces amyloid plaque pathology in transgenic

- mice. *Proc. Natl. Acad. Sci. U.S.A.*, **103** (21), 8215–8220.
131. Hanahan, D. and Weinberg, R.A. (2011) Hallmarks of cancer: the next generation. *Cell*, **144** (5), 646–674.
  132. Carnero, A. and Paramio, J.M. (2014) The PTEN/PI3K/AKT pathway in vivo, cancer mouse models. *Front. Oncol.*, **4**, 252.
  133. Kang, S., Bader, A.G., and Vogt, P.K. (2005) Phosphatidylinositol 3-kinase mutations identified in human cancer are oncogenic. *Proc. Natl. Acad. Sci. U.S.A.*, **102** (3), 802–807.
  134. Adams, J.R., Xu, K., Liu, J.C., Agamez, N.M., Loch, A.J., Wong, R.G., Wang, W., Wright, K.L., Lane, T.F., Zacksenhaus, E., and Egan, S.E. (2011) Cooperation between Pik3ca and p53 mutations in mouse mammary tumor formation. *Cancer Res.*, **71** (7), 2706–2717.
  135. Meyer, D.S., Brinkhaus, H., Muller, U., Muller, M., Cardiff, R.D., and Bentires-Alj, M. (2011) Luminal expression of PIK3CA mutant H1047R in the mammary gland induces heterogeneous tumors. *Cancer Res.*, **71** (13), 4344–4351.
  136. Yuan, W., Stawiski, E., Janakiraman, V., Chan, E., Durinck, S., Edgar, K.A., Kljavin, N.M., Rivers, C.S., Gnad, F., Roose-Girma, M., Haverty, P.M., Fedorowicz, G., Heldens, S., Soriano, R.H., Zhang, Z., Wallin, J.J., Johnson, L., Merchant, M., Modrusan, Z., Stern, H.M., and Seshagiri, S. (2013) Conditional activation of Pik3ca(H1047R) in a knock-in mouse model promotes mammary tumorigenesis and emergence of mutations. *Oncogene*, **32** (3), 318–326.
  137. Meyer, D.S., Koren, S., Leroy, C., Brinkhaus, H., Muller, U., Klebba, I., Muller, M., Cardiff, R.D., and Bentires-Alj, M. (2013) Expression of PIK3CA mutant E545K in the mammary gland induces heterogeneous tumors but is less potent than mutant H1047R. *Oncogenesis*, **2**, e74.
  138. Trejo, C.L., Green, S., Marsh, V., Collisson, E.A., Iezza, G., Phillips, W.A., and McMahon, M. (2013) Mutationally activated PIK3CA(H1047R) cooperates with BRAF(V600E) to promote lung cancer progression. *Cancer Res.*, **73** (21), 6448–6461.
  139. Kinross, K.M., Montgomery, K.G., Kleinschmidt, M., Waring, P., Ivetac, I., Tikoo, A., Saad, M., Hare, L., Roh, V., Mantamadiotis, T., Sheppard, K.E., Ryland, G.L., Campbell, I.G., Gorringer, K.L., Christensen, J.G., Cullinane, C., Hicks, R.J., Pearson, R.B., Johnstone, R.W., McArthur, G.A., and Phillips, W.A. (2012) An activating Pik3ca mutation coupled with pten loss is sufficient to initiate ovarian tumorigenesis in mice. *J. Clin. Invest.*, **122** (2), 553–557.
  140. Hare, L.M., Phesse, T.J., Waring, P.M., Montgomery, K.G., Kinross, K.M., Mills, K., Roh, V., Heath, J.K., Ramsay, R.G., Ernst, M., and Phillips, W.A. (2014) Physiological expression of the PI3K-activating mutation Pik3ca(H1047R) combines with apc loss to promote development of invasive intestinal adenocarcinomas in mice. *Biochem. J.*, **458** (2), 251–258.
  141. Joshi, A., Miller, C. Jr., Baker, S.J., and Ellenson, L.H. (2015) Activated mutant p110alpha causes endometrial carcinoma in the setting of biallelic pten deletion. *Am. J. Pathol.*, **185** (4), 1104–1113.
  142. Blanco-Aparicio, C., Renner, O., Leal, J.F., and Carnero, A. (2007) PTEN, more than the AKT pathway. *Carcinogenesis*, **28** (7), 1379–1386.
  143. Blanco-Aparicio, C., Canamero, M., Cecilia, Y., Pequeno, B., Renner, O., Ferrer, I., and Carnero, A. (2010) Exploring the gain of function contribution of AKT to mammary tumorigenesis in mouse models. *PLoS One*, **5** (2), e9305.
  144. Hutchinson, J.N., Jin, J., Cardiff, R.D., Woodgett, J.R., and Muller, W.J. (2004) Activation of akt-1 (PKB-alpha) can accelerate ErbB-2-mediated mammary tumorigenesis but suppresses tumor invasion. *Cancer Res.*, **64** (9), 3171–3178.
  145. Dillon, R.L., Marcotte, R., Hennessy, B.T., Woodgett, J.R., Mills, G.B., and Muller, W.J. (2009) Akt1 and akt2 play distinct roles in the initiation and metastatic phases of

- mammary tumor progression. *Cancer Res.*, **69** (12), 5057–5064.
146. Maroulakou, I.G., Oemler, W., Naber, S.P., and Tschlis, P.N. (2007) Akt1 ablation inhibits, whereas Akt2 ablation accelerates, the development of mammary adenocarcinomas in mouse mammary tumor virus (MMTV)-ErbB2/neu and MMTV-polyoma middle T transgenic mice. *Cancer Res.*, **67** (1), 167–177.
  147. Martini, M., Ciraolo, E., Gulluni, F., and Hirsch, E. (2013) Targeting PI3K in cancer: any good news? *Front. Oncol.*, **3**, 108.
  148. Miller, B.W., Przepiora, D., de Claro, R.A., Lee, K., Nie, L., Simpson, N., Gudi, R., Saber, H., Shord, S., Bullock, J., Marathe, D., Mehrotra, N., Hsieh, L.S., Ghosh, D., Brown, J., Kane, R.C., Justice, R., Kaminskas, E., Farrell, A.T., and Pazdur, R. (2015) FDA approval: idelalisib monotherapy for the treatment of patients with follicular lymphoma and small lymphocytic lymphoma. *Clin. Cancer Res.*, **21** (77), 1525–1529.
  149. Keating, G.M. (2015) Idelalisib: a review of its use in chronic lymphocytic leukaemia and indolent non-hodgkin's lymphoma. *Target. Oncol.*, **10** (1), 141–151.
  150. Menon, S. and Manning, B.D. (2008) Common corruption of the mTOR signaling network in human tumors. *Oncogene*, **27** (Suppl. 2), S43–S51.
  151. Cargnello, M., Tcherkezian, J., and Roux, P.P. (2015) The expanding role of mTOR in cancer cell growth and proliferation. *Mutagenesis*, **30** (2), 169–176.
  152. Wander, S.A., Hennessy, B.T., and Slingerland, J.M. (2011) Next-generation mTOR inhibitors in clinical oncology: how pathway complexity informs therapeutic strategy. *J. Clin. Invest.*, **121** (4), 1231–1241.
  153. Bauer, T.M., Patel, M.R., and Infante, J.R. (2015) Targeting PI3 kinase in cancer. *Pharmacol. Ther.*, **146**, 53–60.
  154. Owonikoko, T.K. and Khuri, F.R. (2013) Targeting the PI3K/AKT/mTOR pathway: biomarkers of success and tribulation. *Am. Soc. Clin. Oncol. Educ. Book*.
  155. O'Reilly, K.E., Rojo, F., She, Q.B., Solit, D., Mills, G.B., Smith, D., Lane, H., Hofmann, F., Hicklin, D.J., Ludwig, D.L., Baselga, J., and Rosen, N. (2006) mTOR inhibition induces upstream receptor tyrosine kinase signaling and activates akt. *Cancer Res.*, **66** (3), 1500–1508.
  156. Kim, J., Thorne, S.H., Sun, L., Huang, B., and Mochly-Rosen, D. (2011) Sustained inhibition of PKC $\alpha$  reduces intravasation and lung seeding during mammary tumor metastasis in an in vivo mouse model. *Oncogene*, **30** (3), 323–333.
  157. Jr. Sledge, G.W. and Gokmen-Polar, Y. (2006) Protein kinase C- $\beta$  as a therapeutic target in breast cancer. *Semin. Oncol.*, **33** (3, Suppl. 9), S15–S18.
  158. Grossoni, V.C., Todaro, L.B., Kazanietz, M.G., de Kier Joffe, E.D., and Urtreger, A.J. (2009) Opposite effects of protein kinase C  $\beta$ 1 (PKC $\beta$ 1) and PKC $\epsilon$  in the metastatic potential of a breast cancer murine model. *Breast Cancer Res. Treat.*, **118** (3), 469–480.
  159. Mischak, H., Goodnight, J.A., Kolch, W., Martiny-Baron, G., Schaechtle, C., Kazanietz, M.G., Blumberg, P.M., Pierce, J.H., and Mushinski, J.F. (1993) Overexpression of protein kinase C- $\delta$  and - $\epsilon$  in NIH 3T3 cells induces opposite effects on growth, morphology, anchorage dependence, and tumorigenicity. *J. Biol. Chem.*, **268** (9), 6090–6096.
  160. Gorin, M.A. and Pan, Q. (2009) Protein kinase C  $\epsilon$ : an oncogene and emerging tumor biomarker. *Mol. Cancer*, **8**, 9. doi: 10.1186/1476-4598-8-9
  161. Pan, Q., Bao, L.W., Kleer, C.G., Sabel, M.S., Griffith, K.A., Teknos, T.N., and Merajver, S.D. (2005) Protein kinase C  $\epsilon$  is a predictive biomarker of aggressive breast cancer and a validated target for RNA interference anticancer therapy. *Cancer Res.*, **65** (18), 8366–8371.
  162. Parker, P.J., Justilien, V., Riou, P., Linch, M., and Fields, A.P. (2014) Atypical protein kinase  $\epsilon$  as a

- human oncogene and therapeutic target. *Biochem. Pharmacol.*, **88** (1), 1–11.
163. Liu, S.G., Wang, B.S., Jiang, Y.Y., Zhang, T.T., Shi, Z.Z., Yang, Y., Yang, Y.L., Wang, X.C., Lin, D.C., Zhang, Y., Yang, H., Cai, Y., Zhan, Q.M., and Wang, M.R. (2011) Atypical protein kinase ciota (PKCiota) promotes metastasis of esophageal squamous cell carcinoma by enhancing resistance to anoikis via PKCiota-SKP2-AKT pathway. *Mol. Cancer Res.*, **9** (4), 390–402.
  164. Martin-Liberal, J., Lagares-Tena, L., and Larkin, J. (2014) Prospects for MEK inhibitors for treating cancer. *Expert Opin. Drug Saf.*, **13** (4), 483–495.
  165. Santarpia, L., Lippman, S.M., and El-Naggar, A.K. (2012) Targeting the MAPK-RAS-RAF signaling pathway in cancer therapy. *Expert Opin. Ther. Targets*, **16** (1), 103–119.
  166. Robert, C., Karaszewska, B., Schachter, J., Rutkowski, P., Mackiewicz, A., Stroiakovski, D., Lichinitser, M., Dummer, R., Grange, F., Mortier, L., Chiarion-Sileni, V., Drucis, K., Krajsova, I., Hauschild, A., Lorigan, P., Wolter, P., Long, G.V., Flaherty, K., Nathan, P., Ribas, A., Martin, A.M., Sun, P., Crist, W., Legos, J., Rubin, S.D., Little, S.M., and Schadendorf, D. (2015) Improved overall survival in melanoma with combined dabrafenib and trametinib. *N. Engl. J. Med.*, **372** (1), 30–39.
  167. Grossi, V., Peserico, A., Tezil, T., and Simone, C. (2014) p38alpha MAPK pathway: a key factor in colorectal cancer therapy and chemoresistance. *World J. Gastroenterol.*, **20** (29), 9744–9758.
  168. Rudalska, R., Dauch, D., Longerich, T., McJunkin, K., Wuestefeld, T., Kang, T.W., Hohmeyer, A., Pesic, M., Leibold, J., von Thun, A., Schirmacher, P., Zuber, J., Weiss, K.H., Powers, S., Malek, N.P., Eilers, M., Sipos, B., Lowe, S.W., Geffers, R., Laufer, S., and Zender, L. (2014) In vivo RNAi screening identifies a mechanism of sorafenib resistance in liver cancer. *Nat. Med.*, **20** (10), 1138–1146.
  169. Bubici, C. and Papa, S. (2014) JNK signalling in cancer: in need of new, smarter therapeutic targets. *Br. J. Pharmacol.*, **171** (1), 24–37.
  170. Hui, L., Zatloukal, K., Scheuch, H., Stepniak, E., and Wagner, E.F. (2008) Proliferation of human HCC cells and chemically induced mouse liver cancers requires JNK1-dependent p21 down-regulation. *J. Clin. Invest.*, **118** (12), 3943–3953.
  171. Chen, N., Nomura, M., She, Q.B., Ma, W.Y., Bode, A.M., Wang, L., Flavell, R.A., and Dong, Z. (2001) Suppression of skin tumorigenesis in c-jun NH(2)-terminal kinase-2-deficient mice. *Cancer Res.*, **61** (10), 3908–3912.
  172. Wang, H., Wang, H.S., Zhou, B.H., Li, C.L., Zhang, F., Wang, X.F., Zhang, G., Bu, X.Z., Cai, S.H., and Du, J. (2013) Epithelial-mesenchymal transition (EMT) induced by TNF-alpha requires AKT/GSK-3beta-mediated stabilization of snail in colorectal cancer. *PLoS One*, **8** (2), e56664.
  173. Zheng, H., Li, W., Wang, Y., Liu, Z., Cai, Y., Xie, T., Shi, M., Wang, Z., and Jiang, B. (2013) Glycogen synthase kinase-3 beta regulates snail and beta-catenin expression during fas-induced epithelial-mesenchymal transition in gastrointestinal cancer. *Eur. J. Cancer*, **49** (12), 2734–2746.
  174. Wen, W., Ding, J., Sun, W., Fu, J., Chen, Y., Wu, K., Ning, B., Han, T., Huang, L., Chen, C., Xie, D., Li, Z., Feng, G., Wu, M., Xie, W., and Wang, H. (2012) Cyclin G1-mediated epithelial-mesenchymal transition via phosphoinositide 3-kinase/Akt signaling facilitates liver cancer progression. *Hepatology*, **55** (6), 1787–1798.
  175. McCubrey, J.A., Steelman, L.S., Bertrand, F.E., Davis, N.M., Sokolosky, M., Abrams, S.L., Montalto, G., D'Assoro, A.B., Libra, M., Nicoletti, F., Maestro, R., Basecke, J., Rakus, D., Gizak, A., Demidenko, Z.N., Cocco, L., Martelli, A.M., and Cervello, M. (2014) GSK-3 as potential target for therapeutic intervention in cancer. *Oncotarget*, **5** (10), 2881–2911.

176. Desbois-Mouthon, C., Blivet-Van Eggelpoel, M.J., Beurel, E., Boissan, M., Delelo, R., Cadoret, A., and Capeau, J. (2002) Dysregulation of glycogen synthase kinase-3 $\beta$  signaling in hepatocellular carcinoma cells. *Hepatology*, **36** (6), 1528–1536.
177. Farago, M., Dominguez, I., Landesman-Bollag, E., Xu, X., Rosner, A., Cardiff, R.D., and Seldin, D.C. (2005) Kinase-inactive glycogen synthase kinase 3 $\beta$  promotes wnt signaling and mammary tumorigenesis. *Cancer Res.*, **65** (13), 5792–5801.
178. Zhou, W., Wang, L., Gou, S.M., Wang, T.L., Zhang, M., Liu, T., and Wang, C.Y. (2012) ShRNA silencing glycogen synthase kinase-3  $\beta$  inhibits tumor growth and angiogenesis in pancreatic cancer. *Cancer Lett.*, **316** (2), 178–186.
179. Mann, K.M., Ward, J.M., Yew, C.C., Kovochich, A., Dawson, D.W., Black, M.A., Brett, B.T., Sheetz, T.E., Dupuy, A.J., Australian Pancreatic Cancer Genome Initiative, Chang, D.K., Biankin, A.V., Waddell, N., Kassahn, K.S., Grimmond, S.M., Rust, A.G., Adams, D.J., Jenkins, N.A., and Copeland, N.G. (2012) Sleeping beauty mutagenesis reveals cooperating mutations and pathways in pancreatic adenocarcinoma. *Proc. Natl. Acad. Sci. U.S.A.*, **109** (16), 5934–5941.
180. Rahrman, E.P., Watson, A.L., Keng, V.W., Choi, K., Moriarity, B.S., Beckmann, D.A., Wolf, N.K., Sarver, A., Collins, M.H., Moertel, C.L., Wallace, M.R., Gel, B., Serra, E., Ratner, N., and Largaespada, D.A. (2013) Forward genetic screen for malignant peripheral nerve sheath tumor formation identifies new genes and pathways driving tumorigenesis. *Nat. Genet.*, **45** (7), 756–766.
181. Kean, M.J., Couzens, A.L., and Gingras, A.C. (2012) Mass spectrometry approaches to study mammalian kinase and phosphatase associated proteins. *Methods*, **57** (4), 400–408.
182. Zhang, L., Holmes, I.P., Hochgrafe, F., Walker, S.R., Ali, N.A., Humphrey, E.S., Wu, J., de Silva, M., Kersten, W.J., Connor, T., Falk, H., Allan, L., Street, I.P., Bentley, J.D., Pilling, P.A., Monahan, B.J., Peat, T.S., and Daly, R.J. (2013) Characterization of the novel broad-spectrum kinase inhibitor CTx-0294885 as an affinity reagent for mass spectrometry-based kinome profiling. *J. Proteome Res.*, **12** (7), 3104–3116.
183. Dunham, W.H., Mullin, M., and Gingras, A.C. (2012) Affinity-purification coupled to mass spectrometry: basic principles and strategies. *Proteomics*, **12** (10), 1576–1590.
184. Song, J., Hao, Y., Du, Z., Wang, Z., and Ewing, R.M. (2012) Identifying novel protein complexes in cancer cells using epitope-tagging of endogenous human genes and affinity-purification mass spectrometry. *J. Proteome Res.*, **11** (12), 5630–5641.
185. Dalglish, G.L., Furge, K., Greenman, C., Chen, L., Bignell, G., Butler, A., Davies, H., Edkins, S., Hardy, C., Latimer, C., Teague, J., Andrews, J., Barthorpe, S., Beare, D., Buck, G., Campbell, P.J., Forbes, S., Jia, M., Jones, D., Knott, H., Kok, C.Y., Lau, K.W., Leroy, C., Lin, M.L., McBride, D.J., Maddison, M., Maguire, S., McLay, K., Menzies, A., Mironenko, T., Muldergill, L., Mudie, L., O'Meara, S., Pleasance, E., Rajasingham, A., Shepherd, R., Smith, R., Stebbings, L., Stephens, P., Tang, G., Tarpey, P.S., Turrell, K., Dykema, K.J., Khoo, S.K., Petillo, D., Wondergem, B., Anema, J., Kahnoski, R.J., Teh, B.T., Stratton, M.R., and Futreal, P.A. (2010) Systematic sequencing of renal carcinoma reveals inactivation of histone modifying genes. *Nature*, **463** (7279), 360–363.
186. Song, X.M., Fiedler, M., Galuska, D., Ryder, J.W., Fernstrom, M., Chibalin, A.V., Wallberg-Henriksson, H., and Zierath, J.R. (2002) 5-aminoimidazole-4-carboxamide ribonucleoside treatment improves glucose homeostasis in insulin-resistant diabetic (ob/ob) mice. *Diabetologia*, **45** (1), 56–65.
187. Liu, J., Bang, A.G., Kintner, C., Orth, A.P., Chanda, S.K., Ding, S., and Schultz, P.G. (2005) Identification of

- the wnt signaling activator leucine-rich repeat in flightless interaction protein 2 by a genome-wide functional analysis. *Proc. Natl. Acad. Sci. U.S.A.*, **102** (6), 1927–1932.
188. Kim, Y.H., Lee, E.K., Park, S.A., Kim, N.H., and Kim, C.W. (2012) Proteomic analysis of plasma from a tau transgenic mouse. *Int. J. Dev. Neurosci.*, **30** (4), 277–283.
  189. Stuhlmiller, T.J., Earp, H.S., and Johnson, G.L. (2014) Adaptive reprogramming of the breast cancer kinome. *Clin. Pharmacol. Ther.*, **95** (4), 413–415.
  190. Duncan, J.S., Whittle, M.C., Nakamura, K., Abell, A.N., Midland, A.A., Zawistowski, J.S., Johnson, N.L., Granger, D.A., Jordan, N.V., Darr, D.B., Usary, J., Kuan, P.F., Smalley, D.M., Major, B., He, X., Hoadley, K.A., Zhou, B., Sharpless, N.E., Perou, C.M., Kim, W.Y., Gomez, S.M., Chen, X., Jin, J., Frye, S.V., Earp, H.S., Graves, L.M., and Johnson, G.L. (2012) Dynamic reprogramming of the kinome in response to targeted MEK inhibition in triple-negative breast cancer. *Cell*, **149** (2), 307–321.
  191. Cooper, M.J., Cox, N.J., Zimmerman, E.I., Dewar, B.J., Duncan, J.S., Whittle, M.C., Nguyen, T.A., Jones, L.S., Ghose Roy, S., Smalley, D.M., Kuan, P.F., Richards, K.L., Christopherson, R.I., Jin, J., Frye, S.V., Johnson, G.L., Baldwin, A.S., and Graves, L.M. (2013) Application of multiplexed kinase inhibitor beads to study kinome adaptations in drug-resistant leukemia. *PLoS One*, **8** (6), e66755.
  192. Roszik, J., Toth, G., Szollosi, J., and Vereb, G. (2013) Validating pharmacological disruption of protein-protein interactions by acceptor photobleaching FRET imaging. *Methods Mol. Biol.*, **986**, 165–178.
  193. Ng, T., Squire, A., Hansra, G., Bornancin, F., Prevostel, C., Hanby, A., Harris, W., Barnes, D., Schmidt, S., Mellor, H., Bastiaens, P.I., and Parker, P.J. (1999) Imaging protein kinase calpha activation in cells. *Science*, **283** (5410), 2085–2089.
  194. Yadav, R.B., Burgos, P., Parker, A.W., Iadevaia, V., Proud, C.G., Allen, R.A., O'Connell, J.P., Jeshtadi, A., Stubbs, C.D., and Botchway, S.W. (2013) mTOR direct interactions with rheb-GTPase and raptor: sub-cellular localization using fluorescence lifetime imaging. *BMC Cell. Biol.*, **14**, 3. doi: 10.1186/1471-2121-14-3
  195. Veeriah, S., Leboucher, P., de Naurois, J., Jethwa, N., Nye, E., Bunting, T., Stone, R., Stamp, G., Calleja, V., Jeffrey, S.S., Parker, P.J., and Larjani, B. (2014) High-throughput time-resolved FRET reveals Akt/PKB activation as a poor prognostic marker in breast cancer. *Cancer Res.*, **74** (18), 4983–4995.
  196. Shea, K.F., Wells, C.M., Garner, A.P., and Jones, G.E. (2008) ROCK1 and LIMK2 interact in spread but not blebbing cancer cells. *PLoS One*, **3** (10), e3398.
  197. Fernandez-Duenas, V., Llorente, J., Gandia, J., Borroto-Escuela, D.O., Agnati, L.F., Tasca, C.I., Fuxe, K., and Ciruela, F. (2012) Fluorescence resonance energy transfer-based technologies in the study of protein-protein interactions at the cell surface. *Methods*, **57** (4), 467–472.
  198. Larbouret, C., Gaborit, N., Chardes, T., Coelho, M., Campigna, E., Bascoul-Mollevis, C., Mach, J.P., Azria, D., Robert, B., and Pelegrin, A. (2012) In pancreatic carcinoma, dual EGFR/HER2 targeting with cetuximab/trastuzumab is more effective than treatment with trastuzumab/erlotinib or lapatinib alone: implication of receptors' down-regulation and dimers' disruption. *Neoplasia*, **14** (2), 121–130.
  199. Mizutani, T., Kondo, T., Darmanin, S., Tsuda, M., Tanaka, S., Tobiume, M., Asaka, M., and Ohba, Y. (2010) A novel FRET-based biosensor for the measurement of BCR-ABL activity and its response to drugs in living cells. *Clin. Cancer Res.*, **16** (15), 3964–3975.
  200. Morris, M.C. (2013) Fluorescent biosensors – probing protein kinase

- function in cancer and drug discovery. *Biochim. Biophys. Acta*, **1834** (7), 1387–1395.
201. Fire, A., Xu, S., Montgomery, M.K., Kostas, S.A., Driver, S.E., and Mello, C.C. (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*, **391** (6669), 806–811.
  202. Dykxhoorn, D.M., Novina, C.D., and Sharp, P.A. (2003) Killing the messenger: short RNAs that silence gene expression. *Nat. Rev. Mol. Cell Biol.*, **4** (6), 457–467.
  203. Breen, M.E. and Soellner, M.B. (2015) Small molecule substrate phosphorylation site inhibitors of protein kinases: approaches and challenges. *ACS Chem. Biol.*, **10** (1), 175–189.
  204. Schonherr, M., Bhattacharya, A., Kottek, T., Szymczak, S., Koberle, M., Wickenhauser, C., Siebols, U., Saalbach, A., Koczan, D., Magin, T.M., Simon, J.C., and Kunz, M. (2014) Genomewide RNAi screen identifies protein kinase *cb* and new members of mitogen-activated protein kinase pathway as regulators of melanoma cell growth and metastasis. *Pigm. Cell Melanoma Res.*, **27** (3), 418–430.
  205. Izrailit, J., Berman, H.K., Datti, A., Wrana, J.L., and Reedijk, M. (2013) High throughput kinase inhibitor screens reveal TRB3 and MAPK-ERK/TGFbeta pathways as fundamental notch regulators in breast cancer. *Proc. Natl. Acad. Sci. U.S.A.*, **110** (5), 1714–1719.
  206. Banerji, V., Frumm, S.M., Ross, K.N., Li, L.S., Schinzel, A.C., Hahn, C.K., Kakoza, R.M., Chow, K.T., Ross, L., Alexe, G., Tolliday, N., Ingulizian, H., Galinsky, I., Stone, R.M., DeAngelo, D.J., Roti, G., Aster, J.C., Hahn, W.C., Kung, A.L., and Stegmaier, K. (2012) The intersection of genetic and chemical genomic screens identifies GSK-3alpha as a target in human acute myeloid leukemia. *J. Clin. Invest.*, **122** (3), 935–947.
  207. Cronan, M.R., Nakamura, K., Johnson, N.L., Granger, D.A., Cuevas, B.D., Wang, J.G., Mackman, N., Scott, J.E., Dohman, H.G., and Johnson, G.L. (2012) Defining MAP3 kinases required for MDA-MB-231 cell tumor growth and metastasis. *Oncogene*, **31** (34), 3889–3900.
  208. Liu-Sullivan, N., Zhang, J., Bakleh, A., Marchica, J., Li, J., Siolas, D., Laquerre, S., Degenhardt, Y.Y., Wooster, R., Chang, K., Hannon, G.F., and Powers, S. (2011) Pooled shRNA screen for sensitizers to inhibition of the mitotic regulator polo-like kinase (PLK1). *Oncotarget*, **2** (12), 1254–1264.
  209. Iorns, E., Turner, N.C., Elliott, R., Syed, N., Garrone, O., Gasco, M., Tutt, A.N., Crook, T., Lord, C.J., and Ashworth, A. (2008) Identification of CDK10 as an important determinant of resistance to endocrine therapy for breast cancer. *Cancer Cell*, **13** (2), 91–104.
  210. Vora, S.R., Juric, D., Kim, N., Mino-Kenudson, M., Huynh, T., Costa, C., Lockerman, E.L., Pollack, S.F., Liu, M., Li, X., Lehar, J., Wiesmann, M., Wartmann, M., Chen, Y., Cao, Z.A., Pinzon-Ortiz, M., Kim, S., Schlegel, R., Huang, A., and Engelman, J.A. (2014) CDK 4/6 inhibitors sensitize PI3CA mutant breast cancer to PI3K inhibitors. *Cancer Cell*, **26** (1), 136–149.
  211. Thorne, C.A., Wichaidit, C., Coster, A.D., Posner, B.A., Wu, L.F., and Altschuler, S.J. (2015) GSK-3 modulates cellular responses to a broad spectrum of kinase inhibitors. *Nat. Chem. Biol.*, **11** (1), 58–63.
  212. Sauer, B. and Henderson, N. (1988) Site-specific DNA recombination in mammalian cells by the cre recombinase of bacteriophage P1. *Proc. Natl. Acad. Sci. U.S.A.*, **85** (14), 5166–5170.
  213. Sternberg, N. and Hamilton, D. (1981) Bacteriophage P1 site-specific recombination. I. Recombination between loxP sites. *J. Mol. Biol.*, **150** (4), 467–486.
  214. Hall, B., Limaye, A., and Kulkarni, A.B. (2009) Overview: generation of gene knockout mice. *Curr. Protoc. Cell Biol.*, Chapter 19 Unit 19.12 19.12.1-17.
  215. Gu, H., Marth, J.D., Orban, P.C., Mossman, H., and Rajewsky, K. (1994) Deletion of a DNA polymerase beta gene segment in T cells using cell

- type-specific gene targeting. *Science*, **265** (5168), 103–106.
216. Feil, R., Wagner, J., Metzger, D., and Chambon, P. (1997) Regulation of cre recombinase activity by mutated estrogen receptor ligand-binding domains. *Biochem. Biophys. Res. Commun.*, **237** (3), 752–757.
  217. Hayashi, S. and McMahon, A.P. (2002) Efficient recombination in diverse tissues by a tamoxifen-inducible form of cre: a tool for temporally regulated gene activation/inactivation in the mouse. *Dev. Biol.*, **244** (2), 305–318.
  218. Utomo, A.R., Nikitin, A.Y., and Lee, W.H. (1999) Temporal, spatial, and cell type-specific control of cre-mediated DNA recombination in transgenic mice. *Nat. Biotechnol.*, **17** (11), 1091–1096.
  219. Anton, M. and Graham, F.L. (1995) Site-specific recombination mediated by an adenovirus vector expressing the cre recombinase protein: a molecular switch for control of gene expression. *J. Virol.*, **69** (8), 4600–4606.
  220. Patel, S., MacAulay, K., and Woodgett, J.R. (2011) Tissue-specific analysis of glycogen synthase kinase-3alpha (GSK-3alpha) in glucose metabolism: effect of strain variation. *PLoS One*, **6** (1), e15845.
  221. Bhattacharjee, R., Goswami, S., Dudiki, T., Popkie, A.P., Phiel, C.J., Kline, D., and Vijayaraghavan, S. (2015) Targeted disruption of glycogen synthase kinase 3a (gsk3a) in mice affects sperm motility resulting in male infertility. *Biol. Reprod.*, **92** (3), 65.
  222. Lal, H., Zhou, J., Ahmad, F., Zaka, R., Vagnozzi, R.J., Decaul, M., Woodgett, J., Gao, E., and Force, T. (2012) Glycogen synthase kinase-3alpha limits ischemic injury, cardiac rupture, post-myocardial infarction remodeling and death. *Circulation*, **125** (1), 65–75.
  223. Zhou, J., Lal, H., Chen, X., Shang, X., Song, J., Li, Y., Kerkela, R., Doble, B.W., MacAulay, K., DeCaul, M., Koch, W.J., Farber, J., Woodgett, J., Gao, E., and Force, T. (2010) GSK-3alpha directly regulates beta-adrenergic signaling and the response of the heart to hemodynamic stress in mice. *J. Clin. Invest.*, **120** (7), 2280–2291.
  224. Zhou, J., Freeman, T.A., Ahmad, F., Shang, X., Mangano, E., Gao, E., Farber, J., Wang, Y., Ma, X.L., Woodgett, J., Vagnozzi, R.J., Lal, H., and Force, T. (2013) GSK-3alpha is a central regulator of age-related pathologies in mice. *J. Clin. Invest.*, **123** (4), 1821–1832.
  225. Kaidanovich-Beilin, O., Lipina, T.V., Takao, K., van Eede, M., Hattori, S., Laliberte, C., Khan, M., Okamoto, K., Chambers, J.W., Fletcher, P.J., MacAulay, K., Doble, B.W., Henkelman, M., Miyakawa, T., Roder, J., and Woodgett, J.R. (2009) Abnormalities in brain structure and behavior in GSK-3alpha mutant mice. *Mol. Brain*, **2**, 35. doi: 10.1186/1756-6606-2-35
  226. Norregaard, R., Tao, S., Nilsson, L., Woodgett, J.R., Kakade, V., Yu, A.S., Howard, C., and Rao, R. (2015) Glycogen synthase kinase 3alpha regulates urine concentrating mechanism in mice. *Am. J. Physiol. Renal. Physiol.*, **308** (6), F650–F660.
  227. Kimura, T., Yamashita, S., Nakao, S., Park, J.M., Murayama, M., Mizoroki, T., Yoshiike, Y., Sahara, N., and Takashima, A. (2008) GSK-3beta is required for memory reconsolidation in adult brain. *PLoS One*, **3** (10), e3540.
  228. Zhai, P., Sciarretta, S., Galeotti, J., Volpe, M., and Sadoshima, J. (2011) Differential roles of GSK-3beta during myocardial ischemia and ischemia/reperfusion. *Circ. Res.*, **109** (5), 502–511.
  229. Woulfe, K.C., Gao, E., Lal, H., Harris, D., Fan, Q., Vagnozzi, R., DeCaul, M., Shang, X., Patel, S., Woodgett, J.R., Force, T., and Zhou, J. (2010) Glycogen synthase kinase-3beta regulates post-myocardial infarction remodeling and stress-induced cardiomyocyte proliferation in vivo. *Circ. Res.*, **106** (10), 1635–1645.
  230. Ahmad, F., Lal, H., Zhou, J., Vagnozzi, R.J., Yu, J.E., Shang, X., Woodgett, J.R., Gao, E., and Force, T. (2014) Cardiomyocyte-specific deletion

- of Gsk3alpha mitigates post-myocardial infarction remodeling, contractile dysfunction, and heart failure. *J. Am. Coll. Cardiol.*, **64** (7), 696–706.
231. Maurin, H., Lechat, B., Dewachter, I., Ris, L., Louis, J.V., Borghgraef, P., Devijver, H., Jaworski, T., and Van Leuven, F. (2013) Neurological characterization of mice deficient in GSK3alpha highlight pleiotropic physiological functions in cognition and pathological activity as tau kinase. *Mol. Brain*, **6**, 27. doi: 10.1186/1756-6606-6-27
232. Urs, N.M., Snyder, J.C., Jacobsen, J.P., Peterson, S.M., and Caron, M.G. (2012) Deletion of GSK3beta in D2R-expressing neurons reveals distinct roles for beta-arrestin signaling in antipsychotic and lithium action. *Proc. Natl. Acad. Sci. U.S.A.*, **109** (50), 20732–20737.
233. Zhou, W., Chen, L., Paul, J., Yang, S., Li, F., Sampson, K., Woodgett, J.R., Beaulieu, J.M., Gamble, K.L., and Li, X. (2012) The effects of glycogen synthase kinase-3beta in serotonin neurons. *PLoS One*, **7** (8), e43262.
234. Patel, S., Doble, B.W., MacAulay, K., Sinclair, E.M., Drucker, D.J., and Woodgett, J.R. (2008) Tissue-specific role of glycogen synthase kinase 3beta in glucose homeostasis and insulin action. *Mol. Cell. Biol.*, **28** (20), 6314–6328.
235. Pansters, N.A., Schols, A.M., Verhees, K.J., de Theije, C.C., Snepvangers, F.J., Kelders, M.C., Ubags, N.D., Haegens, A., and Langen, R.C. (2015) Muscle-specific GSK-3beta ablation accelerates regeneration of disuse-atrophied skeletal muscle. *Biochim. Biophys. Acta*, **1852** (3), 490–506.
236. Liu, Y., Tanabe, K., Baronnier, D., Patel, S., Woodgett, J., Cras-Meneur, C., and Permutt, M.A. (2010) Conditional ablation of gsk-3beta in islet beta cells results in expanded mass and resistance to fat feeding-induced diabetes in mice. *Diabetologia*, **53** (12), 2600–2610.
237. Tanabe, K., Liu, Z., Patel, S., Doble, B.W., Li, L., Cras-Meneur, C., Martinez, S.C., Welling, C.M., White, M.F., Bernal-Mizrachi, E., Woodgett, J.R., and Permutt, M.A. (2008) Genetic deficiency of glycogen synthase kinase-3beta corrects diabetes in mouse models of insulin resistance. *PLoS Biol.*, **6** (2), e37.
238. Kim, W.Y., Wang, X., Wu, Y., Doble, B.W., Patel, S., Woodgett, J.R., and Snider, W.D. (2009) GSK-3 is a master regulator of neural progenitor homeostasis. *Nat. Neurosci.*, **12** (11), 1390–1397.
239. Dembowy, J., Adissu, H.A., Liu, J.C., Zacksenhaus, E., and Woodgett, J.R. (2014) Effect of glycogen synthase kinase-3 inactivation on mouse mammary gland development and oncogenesis. *Oncogene*, doi: 10.1038/onc.2014.279
240. Kyriakis, J.M. and Avruch, J. (2012) Mammalian MAPK signal transduction pathways activated by stress and inflammation: a 10-year update. *Physiol. Rev.*, **92** (2), 689–737.
241. Wang, X., Destrument, A., and Tournier, C. (2007) Physiological roles of MKK4 and MKK7: insights from animal models. *Biochim. Biophys. Acta*, **1773** (8), 1349–1357.
242. Lim, S. and Kaldis, P. (2013) Cdks, cyclins and CKIs: roles beyond cell cycle regulation. *Development*, **140** (15), 3079–3093.
243. Moore, J.D. (2015) The impact of CRISPR-Cas9 on target identification and validation. *Drug Discovery Today*, **20** (4), 450–457.
244. Hsu, P.D., Lander, E.S., and Zhang, F. (2014) Development and applications of CRISPR-Cas9 for genome engineering. *Cell*, **157** (6), 1262–1278.
245. Mojica, F.J., Diez-Villasenor, C., Garcia-Martinez, J., and Soria, E. (2005) Intervening sequences of regularly spaced prokaryotic repeats derive from foreign genetic elements. *J. Mol. Evol.*, **60** (2), 174–182.
246. Pourcel, C., Salvignol, G., and Vergnaud, G. (2005) CRISPR elements in yersinia pestis acquire new repeats by preferential uptake of bacteriophage DNA, and provide additional tools for evolutionary studies. *Microbiology*, **151**(Pt. 3), 653–663.

247. Hartenian, E. and Doench, J.G. (2015) Genetic screens and functional genomics using CRISPR/Cas9 technology. *FEBS J.*, **282** (8), 1383–1393.
248. Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J.A., and Charpentier, E. (2012) A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science*, **337** (6096), 816–821.
249. Cong, L., Ran, F.A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P.D., Wu, X., Jiang, W., Marraffini, L.A., and Zhang, F. (2013) Multiplex genome engineering using CRISPR/Cas systems. *Science*, **339** (6121), 819–823.
250. Mali, P., Yang, L., Esvelt, K.M., Aach, J., Guell, M., DiCarlo, J.E., Norville, J.E., and Church, G.M. (2013) RNA-guided human genome engineering via Cas9. *Science*, **339** (6121), 823–826.
251. Wang, T., Wei, J.J., Sabatini, D.M., and Lander, E.S. (2014) Genetic screens in human cells using the CRISPR-Cas9 system. *Science*, **343** (6166), 80–84.
252. Chen, S., Sanjana, N.E., Zheng, K., Shalem, O., Lee, K., Shi, X., Scott, D.A., Song, J., Pan, J.Q., Weissleder, R., Lee, H., Zhang, F., and Sharp, P.A. (2015) Genome-wide CRISPR screen in a mouse model of tumor growth and metastasis. *Cell*, **160** (6), 1246–1260.
253. Shalem, O., Sanjana, N.E., Hartenian, E., Shi, X., Scott, D.A., Mikkelsen, T.S., Heckl, D., Ebert, B.L., Root, D.E., Doench, J.G., and Zhang, F. (2014) Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science*, **343** (6166), 84–87.
254. Matano, M., Date, S., Shimokawa, M., Takano, A., Fujii, M., Ohta, Y., Watanabe, T., Kanai, T., and Sato, T. (2015) Modeling colorectal cancer using CRISPR-Cas9-mediated engineering of human intestinal organoids. *Nat. Med.*, **21** (3), 256–262.
255. Zhang, L.L., Cao, F.F., Wang, Y., Meng, F.L., Zhang, Y., Zhong, D.S., and Zhou, Q.H. (2014) The protein kinase C (PKC) inhibitors combined with chemotherapy in the treatment of advanced non-small cell lung cancer: meta-analysis of randomized controlled trials. *Clin. Transl. Oncol.*, **17** (5), 371–377.
256. Antal, C.E., Hudson, A.M., Kang, E., Zanca, C., Wirth, C., Stephenson, N.L., Trotter, E.W., Gallegos, L.L., Miller, C.J., Furnari, F.B., Hunter, T., Brognard, J., and Newton, A.C. (2015) Cancer-associated protein kinase C mutations reveal kinase's role as tumor suppressor. *Cell*, **160** (3), 489–502.
257. Xue, W., Chen, S., Yin, H., Tammela, T., Papagiannakopoulos, T., Joshi, N.S., Cai, W., Yang, G., Bronson, R., Crowley, D.G., Zhang, F., Anderson, D.G., Sharp, P.A., and Jacks, T. (2014) CRISPR-mediated direct mutation of cancer genes in the mouse liver. *Nature*, **514** (7522), 380–384.
258. Platt, R.J., Chen, S., Zhou, Y., Yim, M.J., Swiech, L., Kempton, H.R., Dahlman, J.E., Parnas, O., Eisenhaure, T.M., Jovanovic, M., Graham, D.B., Jhunjhunwala, S., Heidenreich, M., Xavier, R.J., Langer, R., Anderson, D.G., Hacohen, N., Regev, A., Feng, G., Sharp, P.A., and Zhang, F. (2014) CRISPR-Cas9 knockin mice for genome editing and cancer modeling. *Cell*, **159** (2), 440–455.
259. Doudna, J.A. and Charpentier, E. (2014) Genome editing. The new frontier of genome engineering with CRISPR-Cas9. *Science*, **346** (6213), 1258096.
260. Sander, J.D. and Joung, J.K. (2014) CRISPR-cas systems for editing, regulating and targeting genomes. *Nat. Biotechnol.*, **32** (4), 347–355.

